

FORM PTO-1390 (Modified)  
(REV 11-2000)

U.S. DEPARTMENT OF COMMERCE PATENT AND TRADEMARK OFFICE

ATTORNEY'S DOCKET NUMBER

**TRANSMITTAL LETTER TO THE UNITED STATES**  
**DESIGNATED/ELECTED OFFICE (DO/EO/US)**  
**CONCERNING A FILING UNDER 35 U.S.C. 371**

**3631-0112P**

U.S. APPLICATION NO (IF KNOWN, SEE 37 CFR

**NEW 09/980916**

INTERNATIONAL APPLICATION NO.  
**PCT/DK00/00205**

INTERNATIONAL FILING DATE  
**19 April 2000**

PRIORITY DATE CLAIMED  
**23 April 1999**

TITLE OF INVENTION

**METHOD FOR DOWN-REGULATING IL5 ACTIVITY**

APPLICANT(S) FOR DO/EO/US

**KLYSNER, Steen**

Applicant herewith submits to the United States Designated/Elected Office (DO/EO/US) the following items and other information:

1. ☒ This is a **FIRST** submission of items concerning a filing under 35 U.S.C. 371.
2. ☐ This is a **SECOND** or **SUBSEQUENT** submission of items concerning a filing under 35 U.S.C. 371.
3. ☒ This is an express request to begin national examination procedures (35 U.S.C. 371(f)). The submission must include items (5), (6), (9) and (24) indicated below.
4. ☒ The US has been elected by the expiration of 19 months from the priority date (Article 31).
5. ☒ A copy of the International Application as filed (35 U.S.C. 371 (c) (2))
  - a. ☐ is attached hereto (required only if not communicated by the International Bureau).
  - b. ☒ has been communicated by the International Bureau.
  - c. ☐ is not required, as the application was filed in the United States Receiving Office (RO/US).
6. ☐ An English language translation of the International Application as filed (35 U.S.C. 371(c)(2)).
  - a. ☐ is attached hereto.
  - b. ☐ has been previously submitted under 35 U.S.C. 154(d)(4).
7. ☒ Amendments to the claims of the International Application under PCT Article 19 (35 U.S.C. 371 (c)(3))
  - a. ☐ are attached hereto (required only if not communicated by the International Bureau).
  - b. ☐ have been communicated by the International Bureau.
  - c. ☐ have not been made; however, the time limit for making such amendments has NOT expired.
  - d. ☒ have not been made and will not be made.
8. ☐ An English language translation of the amendments to the claims under PCT Article 19 (35 U.S.C. 371(c)(3)).
9. ☐ An oath or declaration of the inventor(s) (35 U.S.C. 371 (c)(4)).
10. ☐ An English language translation of the annexes to the International Preliminary Examination Report under PCT Article 36 (35 U.S.C. 371 (c)(5)).
11. ☒ A copy of the International Preliminary Examination Report (PCT/IPEA/409).
12. ☒ A copy of the International Search Report (PCT/ISA/210).

**Items 13 to 20 below concern document(s) or information included:**

13. ☒ An Information Disclosure Statement under 37 CFR 1.97 and 1.98.
14. ☐ An assignment document for recording. A separate cover sheet in compliance with 37 CFR 3.28 and 3.31 is included.
15. ☒ A **FIRST** preliminary amendment.
16. ☐ A **SECOND** or **SUBSEQUENT** preliminary amendment.
17. ☐ A substitute specification.
18. ☐ A change of power of attorney and/or address letter.
19. ☒ A computer-readable form of the sequence listing in accordance with PCT Rule 13ter.2 and 35 U.S.C. 1.821 - 1.825.
20. ☐ A second copy of the published international application under 35 U.S.C. 154(d)(4).
21. ☐ A second copy of the English language translation of the international application under 35 U.S.C. 154(d)(4).
22. ☒ Certificate of Mailing by Express Mail
23. ☒ Other items or information:

**Application Data Sheet**

**Paper copy of sequence listing**

**Seven (7) sheets of formal drawings**

U.S. APPLICATION NO. (IF KNOWN, SEE 37 CFR

INTERNATIONAL APPLICATION NO

ATTORNEY'S DOCKET NUMBER

09/2980916

PCT/DK00/00205

3631-0112P

24. The following fees are submitted:

**BASIC NATIONAL FEE ( 37 CFR 1.492 (a) (1) - (5) ) :**

- ☒ Neither international preliminary examination fee (37 CFR 1.482) nor international search fee (37 CFR 1.445(a)(2)) paid to USPTO and International Search Report not prepared by the EPO or JPO ..... **\$1040.00**
- ☐ International preliminary examination fee (37 CFR 1.482) not paid to USPTO but International Search Report prepared by the EPO or JPO ..... **\$890.00**
- ☐ International preliminary examination fee (37 CFR 1.482) not paid to USPTO but international search fee (37 CFR 1.445(a)(2)) paid to USPTO ..... **\$740.00**
- ☐ International preliminary examination fee (37 CFR 1.482) paid to USPTO but all claims did not satisfy provisions of PCT Article 33(1)-(4) ..... **\$710.00**
- ☐ International preliminary examination fee (37 CFR 1.482) paid to USPTO and all claims satisfied provisions of PCT Article 33(1)-(4) ..... **\$100.00**

**ENTER APPROPRIATE BASIC FEE AMOUNT =****CALCULATIONS PTO USE ONLY****\$1,040.00**Surcharge of **\$130.00** for furnishing the oath or declaration later than ☐ 20 ☒ 30 months from the earliest claimed priority date (37 CFR 1.492 (e)).**\$130.00**

CLAIMS	NUMBER FILED	NUMBER EXTRA	RATE
Total claims	65 - 20 =	45	x \$18.00
Independent claims	7 - 3 =	4	x \$84.00

**\$810.00****\$336.00**Multiple Dependent Claims (check if applicable). ☒**\$280.00****TOTAL OF ABOVE CALCULATIONS =****\$2,596.00**

- ☐ Applicant claims small entity status. See 37 CFR 1.27). The fees indicated above are reduced by 1/2.

**\$0.00****SUBTOTAL =****\$2,596.00**Processing fee of **\$130.00** for furnishing the English translation later than ☐ 20 ☐ 30 months from the earliest claimed priority date (37 CFR 1.492 (f)).**\$0.00****TOTAL NATIONAL FEE =****\$2,596.00**Fee for recording the enclosed assignment (37 CFR 1.21(h)). The assignment must be accompanied by an appropriate cover sheet (37 CFR 3.28, 3.31) (check if applicable). ☐**\$0.00****TOTAL FEES ENCLOSED =****\$2,596.00**Amount to be:  
refunded

\$

charged

\$

- a. ☒ A check in the amount of **\$2,596.00** to cover the above fees is enclosed.
- b. ☐ Please charge my Deposit Account No. \_\_\_\_\_ in the amount of \_\_\_\_\_ to cover the above fees. A duplicate copy of this sheet is enclosed.
- c. ☒ The Commissioner is hereby authorized to charge any additional fees which may be required, or credit any overpayment to Deposit Account No. **02-2448**. A duplicate copy of this sheet is enclosed.
- d. ☐ Fees are to be charged to a credit card. **WARNING:** Information on this form may become public. **Credit card information should not be included on this form.** Provide credit card information and authorization on PTO-2038.

**NOTE: Where an appropriate time limit under 37 CFR 1.494 or 1.495 has not been met, a petition to revive (37 CFR 1.137(a) or (b)) must be filed and granted to restore the application to pending status.**

SEND ALL CORRESPONDENCE TO:

Birch, Stewart, Kolasch & Birch, LLP or Customer No. 2292  
P.O. Box 747  
Falls Church, VA 22040-0747  
(714) 708-8555

SIGNATURE

for Leonard R. Svensson

NAME

**30,330**

REGISTRATION NUMBER

**23 October 2001**

DATE

/lmt

PATENT  
3631-0112P

IN THE U.S. PATENT AND TRADEMARK OFFICE

Applicant: KLYSNER, Steen Conf.:  
Int'l. Appl. No.: PCT/DK00/00205  
Appl. No.: NEW Group:  
Filed: October 19, 2001 Examiner:  
For: METHOD FOR DOWN-REGULATING IL5  
ACTIVITY

PRELIMINARY AMENDMENT

**BOX PATENT APPLICATION**

Assistant Commissioner for Patents  
Washington, DC 20231

October 23, 2001

Sir:

The following Preliminary Amendments and Remarks are respectfully submitted in connection with the above-identified application.

AMENDMENTS

IN THE SPECIFICATION:

Please amend the specification as follows:

Before line 1, insert --This application is the national phase under 35 U.S.C. § 371 of PCT International Application No. PCT/DK00/00205 which has an International filing date of April 19, 2000, which designated the United States of America and was published in English.- -

Please replace the Sequence Listing located after the drawings with the Substitute Sequence Listing enclosed herewith.

20061209 09150360

In the Claims:

Please cancel claims 1-68 without prejudice or disclaimer of the subject matter contained therein.

Please add the following claims:

--69. A method for *in vivo* down-regulation of interleukin 5 (IL5) activity in an animal, including a human being, the method comprising effecting presentation to the animal's immune system of an immunogenically effective amount of

- at least one IL5 polypeptide autologous in the animal or a subsequence thereof which has been formulated so that immunization of the animal with the autologous IL5 polypeptide or subsequence thereof induces production by the animal of antibodies against the IL5 polypeptide, and/or
- at least one IL5 analogue wherein is introduced at least one modification in the amino acid sequence of the animal's autologous IL5 polypeptide which has as a result that immunization of the animal with the analogue induces production of antibodies in the animal against the animal's autologous IL5 polypeptide.

70. The method according to claim 69, wherein is presented an IL5 analogue with at least one modification of the IL5 amino acid sequence.

71. The method according to claim 70, wherein the modification has as a result that a substantial fraction of IL5 B-cell epitopes are preserved and that

- at least one foreign T helper lymphocyte epitope ( $T_H$  epitope) is introduced, and/or
- at least one first moiety is introduced which effects targeting of the modified molecule to an antigen presenting cell (APC) or a B-lymphocyte, and/or



- at least one second moiety is introduced which stimulates the immune system, and/or
- at least one third moiety is introduced which optimizes presentation of the modified IL5 polypeptide to the immune system.

72. The method according to claim 71, wherein the modification includes introduction as side groups, by covalent or non-covalent binding to suitable chemical groups in IL5 or a subsequence thereof, of the foreign T<sub>H</sub> epitope and/or of the first and/or of the second and/or of the third moiety.

73. The method according to claim 71, wherein the modification includes amino acid substitution and/or deletion and/or insertion and/or addition.

74. The method according to claim 73, wherein the modification results in the provision of a fusion polypeptide.

75. The method according to claim 73, wherein introduction of the amino acid substitution and/or deletion and/or insertion and/or addition results in a substantial preservation of the overall tertiary structure of IL5.

76. The method according to claim 70, wherein the modification includes duplication of at least one IL5 B-cell epitope and/or introduction of a hapten.

77. The method according to claim 71, wherein the foreign T-cell epitope is immunodominant in the animal.

78. The method according to claim 71, wherein the foreign T-cell epitope is promiscuous.

79. The method according to claim 78, wherein the at least one foreign T-cell epitope is selected from a natural promiscuous T-cell epitope and an artificial MHC-II binding peptide sequence.

80. The method according to claim 79, wherein the natural T-cell epitope is selected from a Tetanus toxoid epitope such as P2 or P30, a diphtheria toxoid epitope, an influenza virus hemagglutinin epitope, and a *P. falciparum* CS epitope.

81. The method according to claim 71, wherein the first moiety is a substantially specific binding partner for a B-lymphocyte specific surface antigen or for an APC specific surface antigen, such as a hapten or a carbohydrate for which there is a receptor on the B-lymphocyte or the APC.

82. The method according to claim 71, wherein the second moiety is selected from a cytokine, a hormone, and a heat-shock protein.

83. The method according to claim 74, wherein the cytokine is selected from, or is an effective part of, interferon  $\gamma$  (IFN- $\gamma$ ), Flt3L, interleukin 1 (IL-1), interleukin 2 (IL-2), interleukin 4 (IL-4), interleukin 6 (IL-6), interleukin 12 (IL-12), interleukin 13 (IL-13), interleukin 15 (IL-15), and granulocyte-macrophage colony stimulating factor (GM-CSF), and the heat-shock protein is selected from, or is an effective part of any of, HSP70, HSP90, HSC70, GRP94, and calreticulin (CRT).

84. The method according to claim 71, wherein the third moiety is of lipid nature, such as a palmitoyl group, a myristyl group, a farnesyl group, a geranyl-geranyl group, a GPI-anchor, and an N-acyl diglyceride group.

85. The method according to claim 69, wherein the IL5 polypeptide has been modified in at least one of loops 1-3 or in the amino

acid residues C-terminal to helix D, said loops and said helix D corresponding to those shown in Fig. 3 for human and murine IL5.

86. The method according to claim 85, wherein the IL5 polypeptide is a human IL5 polypeptide.

87. The method according to claim 86, wherein the human IL5 polypeptide has been modified by substituting at least one amino acid sequence in SEQ ID NO: 1 with at least one amino acid sequence of equal or different length thereby giving rise to a foreign T<sub>H</sub> epitope, wherein substituted amino acid residues are selected from the group consisting of residues 87-90, residues 88-91, residues 32-43, residues 33-43, residues 59-64, residues 86-91, and residues 110-113.

88. The method according to claim 69, wherein presentation to the immune system is effected by having at least two copies of the IL5 polypeptide, the subsequence thereof or the modified IL5 polypeptide covalently or non-covalently linked to a carrier molecule capable of effecting presentation of multiple copies of antigenic determinants.

89. The method according to claim 69, wherein the IL5 polypeptide, the subsequence thereof, or the modified IL5 polypeptide has been formulated with an adjuvant which facilitates breaking of autotolerance to autoantigens.

90. The method according to claim 89, wherein the adjuvant is selected from the group consisting of an immune targeting adjuvant; an immune modulating adjuvant such as a toxin, a cytokine and a mycobacterial derivative; an oil formulation; a polymer; a micelle forming adjuvant; a saponin; an immunostimulating complex matrix (an ISCOM matrix); a particle; DDA; aluminium adjuvants; DNA adjuvants;  $\gamma$ -inulin; and an encapsulating adjuvant.

91. The method according to claim 69, wherein an effective amount of the IL5 polypeptide or the IL5 analogue is administered to the animal via a route selected from the parenteral route such as the intradermal, the subdermal, the intracutaneous, the subcutaneous, and the intramuscular routes; the peritoneal route; the oral route; the buccal route; the sublingual route; the epidural route; the spinal route; the anal route; and the intracranial route.

92. The method according to claim 91, wherein the effective amount is between 0.5 µg and 2,000 µg of the IL5 polypeptide, the subsequence thereof or the analogue thereof.

93. The method according to claim 91, which includes at least one administration of the IL5 polypeptide or analogue per year, such as at least 2, at least 3, at least 4, at least 6, and at least 12 administrations per year.

94. The method according to claim 91, wherein the IL5 polypeptide or analogue is contained in a virtual lymph node (VLN) device.

95. The method according to claim 69, wherein presentation of modified IL5 to the immune system is effected by introducing nucleic acid(s) encoding the modified IL5 into the animal's cells and thereby obtaining *in vivo* expression by the cells of the nucleic acid(s) introduced.

96. The method according to claim 95, wherein the nucleic acid(s) introduced is/are selected from naked DNA, DNA formulated with charged or uncharged lipids, DNA formulated in liposomes, DNA included in a viral vector, DNA formulated with a transfection-facilitating protein or polypeptide, DNA formulated with a targeting protein or polypeptide, DNA formulated with Calcium precipitating agents, DNA coupled to an inert carrier molecule,

DNA encapsulated in chitin or chitosan, and DNA formulated with an adjuvant.

97. The method according to claim 95, wherein the nucleic acids are administered intraarterially, intravenously, or by the routes defined in claim 91.

98. The method according to claim 96, wherein the nucleic acid(s) is/are contained in a VLN device.

99. The method according to claim 96, which includes at least one administration of the nucleic acids per year, such as at least 2, at least 3, at least 4, at least 6, and at least 12 administrations per year.

100. A method for treating and/or preventing and/or ameliorating asthma or other chronic allergic conditions characterized by eosinophilia, the method comprising down-regulating IL5 activity according to the method according to claim 69 to such an extent that the number of eosinophil cells, either systemically or locally at the disease focus, is significantly reduced, such as a reduction of at least 20%.

101. An IL5 analogue which is derived from an animal IL5 polypeptide wherein is introduced a modification which has as a result that immunization of the animal with the analogue induces production of antibodies against the IL5 polypeptide, and wherein the modification involves amino acid substitution and/or insertion and/or deletion to any one of loops 1-3 or C-terminally to helix D in IL5.

102. An IL5 analogue according to claim 101, wherein the modification has as a result that a substantial fraction of IL5 B-cell epitopes are preserved and that

- at least one foreign T helper lymphocyte epitope ( $T_H$  epitope) is introduced, and/or
- at least one first moiety is introduced which effects targeting of the modified molecule to an antigen presenting cell (APC) or a B-lymphocyte, and/or
- at least one second moiety is introduced which stimulates the immune system, and/or
- at least one third moiety is introduced which optimizes presentation of the modified IL5 polypeptide to the immune system. .

103. An immunogenic composition comprising an immunogenically effective amount of an IL5 polypeptide autologous in an animal, said IL5 polypeptide being formulated together with an immunologically acceptable adjuvant so as to break the animal's autotolerance towards the IL5 polypeptide, the composition further comprising a pharmaceutically and immunologically acceptable carrier and/or vehicle.

104. An immunogenic composition comprising an immunogenically effective amount of an IL5 analogue according to claim 101, the composition further comprising a pharmaceutically and immunologically acceptable carrier and/or vehicle and optionally an adjuvant.

105. An immunogenic composition according to claim 103, wherein the adjuvant is selected from the group consisting of an immune targeting adjuvant; an immune modulating adjuvant such as a toxin, a cytokine and a mycobacterial derivative; an oil formulation; a polymer; a micelle forming adjuvant; a saponin; an immunostimulating complex matrix (an ISCOM matrix); a particle; DDA; aluminium adjuvants; DNA adjuvants;  $\gamma$ -inulin; and an encapsulating adjuvant.

106. A nucleic acid fragment which encodes an IL5 analogue according to claim 101.

107. A vector carrying the nucleic acid fragment according to claim 106.

108. The vector according to claim 107 which is capable of autonomous replication.

109. The vector according to claim 107 which is selected from the group consisting of a plasmid, a phage, a cosmid, a mini-chromosome, and a virus.

110. The vector according to claim 107, comprising, in the 5'→3' direction and in operable linkage, a promoter for driving expression of a nucleic acid fragment which encodes an IL5 analogue derived from an animal IL5 polypeptide wherein is introduced a modification which has as a result that immunization of the animal with the analogue induces production of antibodies against the IL5 polypeptide, and wherein the modification involves amino acid substitution and/or insertion and/or deletion to any one of loops 1-3 or C-terminally to helix D in IL5, optionally a nucleic acid sequence encoding a leader peptide enabling secretion of or integration into the membrane of the polypeptide fragment, said nucleic acid fragment, and optionally a terminator.

111. The vector according to claim 107 which, when introduced into a host cell, is integrated in the host cell genome.

112. The vector according to claim 107 which, when introduced into a host cell, is not capable of being integrated in the host cell genome.

113. The vector according to claim 107, wherein the promoter drives expression in a eukaryotic cell and/or in a prokaryotic cell.

114. A transformed cell carrying the vector according to claim 107.

115. The transformed cell according to claim 114 which is capable of replicating said nucleic acid fragment.

116. The transformed cell according to claim 115, which is a microorganism selected from a bacterium, a yeast, a protozoan, or a cell derived from a multicellular organism selected from a fungus, an insect cell such as an S<sub>2</sub> or an SF cell, a plant cell, and a mammalian cell.

117. The transformed cell according to claim 116 which is a bacterium of the genus *Escherichia*, *Bacillus*, *Salmonella*, or *Mycobacterium*.

118. The transformed cell according to claim 120, which is selected from the group consisting of an *E. coli* cell, and a non-pathogenic *Mycobacterium* cell such as *M. bovis* BCG.

119. The transformed cell according to claim 114, which expresses said nucleic acid fragment.

120. The transformed cell according to claim 123, which secretes or carries on its surface, an IL5 analogue derived from an animal IL5 polypeptide wherein is introduced a modification which has as a result that immunization of the animal with the analogue induces production of antibodies against the IL5 polypeptide, and wherein the modification involves amino acid substitution and/or insertion and/or deletion to any one of loops 1-3 or C-terminally to helix D in IL5.



121. The method according to claim 69, wherein presentation to the immune system is effected by administering a non-pathogenic microorganism or virus which is carrying a nucleic acid fragment which encodes and expresses the IL5 polypeptide or analogue.

122. The method according to claim 121, wherein the virus is a non-virulent pox virus such as a vaccinia virus.

123. The method according to claim 122, wherein the microorganism is a bacterium.

124. The method according to claim 121, wherein the non-pathogenic microorganism or virus is administered one single time to the animal.

125. A composition for inducing production of antibodies against IL5, the composition comprising

- a nucleic acid fragment which encodes an IL5 analogue derived from an animal IL5 polypeptide wherein is introduced a modification which has as a result that immunization of the animal with the analogue induces production of antibodies against the IL5 polypeptide, and wherein the modification involves amino acid substitution and/or insertion and/or deletion to any one of loops 1-3 or C-terminally to helix D in IL5 or a vector carrying said nucleic acid fragment, and
- a pharmaceutically and immunologically acceptable carrier and/or vehicle and/or adjuvant.

126. The composition according to claim 125, wherein the nucleic acid fragment is selected from naked DNA, DNA formulated with charged or uncharged lipids, DNA formulated in liposomes, DNA included in a viral vector, DNA formulated with a transfection-facilitating protein or polypeptide, DNA formulated with a

targeting protein or polypeptide, DNA formulated with Calcium precipitating agents, DNA coupled to an inert carrier molecule, DNA encapsulated in chitin or chitosan, and DNA formulated with an adjuvant.

127. A stable cell line which carries a vector carrying a nucleic acid fragment encoding an IL5 analogue derived from an animal IL5 polypeptide wherein is introduced a modification which has as a result that immunization of the animal with the analogue induces production of antibodies against the IL5 polypeptide, and wherein the modification involves amino acid substitution and/or insertion and/or deletion to any one of loops 1-3 or C-terminally to helix D in IL5, and which expresses said nucleic acid fragment, and which optionally secretes or carries said IL5 analogue on its surface.

128. A method for the preparation of the cell according to claim 114, the method comprising transforming a host cell with the nucleic acid fragment which encodes an IL5 analogue derived from an animal IL5 polypeptide wherein is introduced a modification which has as a result that immunization of the animal with the analogue induces production of antibodies against the IL5 polypeptide, and wherein the modification involves amino acid substitution and/or insertion and/or deletion to any one of loops 1-3 or C-terminally to helix D in IL5 or with a vector carrying said nucleic acid fragment.

129. A method for the identification of a modified IL5 polypeptide which is capable of inducing antibodies against unmodified IL5 in an animal species where the unmodified IL5 polypeptide is a self-protein, the method comprising

- preparing, by means of peptide synthesis or genetic engineering techniques, a set of mutually distinct modified IL5 polypeptides wherein amino acids have been added to,

inserted in, deleted from, or substituted into the amino acid sequence of an IL5 polypeptide of the animal species thereby giving rise to amino acid sequences in the set which comprise T-cell epitopes which are foreign to the animal species, or preparing a set of nucleic acid fragments encoding the set of mutually distinct modified IL5 polypeptides,

- testing members of the set for their ability to induce production of antibodies by the animal species against the unmodified IL5, and
- identifying and optionally isolating the member(s) of the set which significantly induces antibody production against unmodified IL5 in the animal species, or identifying and optionally isolating the polypeptide expression products encoded by members of the set of nucleic acid fragments which significantly induces antibody production against unmodified IL5 in the animal species.

130. A method for the preparation of an immunogenic composition comprising at least one modified IL5 polypeptide which is capable of inducing antibodies against unmodified IL5 in an animal species where the unmodified IL5 polypeptide is a self-protein, the method comprising

- preparing, by means of peptide synthesis or genetic engineering techniques, a set of mutually distinct modified IL5 polypeptides wherein amino acids have been added to, inserted in, deleted from, or substituted into the amino acid sequence of an IL5 polypeptide of the animal species thereby giving rise to amino acid sequences in the set comprising T-cell epitopes which are foreign to the animal,
- testing members of the set for their ability to induce production of antibodies by the animal species against the unmodified IL5, and
- admixing the member(s) of the set which significantly

induces production of antibodies in the animal species which are reactive with IL5 with a pharmaceutically and immunologically acceptable carrier and/or vehicle, optionally in combination with at least one pharmaceutically and immunologically acceptable adjuvant.

131. The method according to claim 129, wherein preparation of the members of the set comprises preparation of mutually distinct nucleic acid sequences, each sequence being a nucleic acid sequence according to claim 106, insertion of the nucleic acid sequences into appropriate expression vectors, transformation of suitable host cells with the vectors, and expression of the nucleic acid sequences, optionally followed by isolation of the expression products.

132. The method according to claim 131, wherein the preparation of the nucleic acid sequences and/or the vectors is achieved by the aid of a molecular amplification technique such as PCR, or by the aid of nucleic acid synthesis.--

REMARKS

Claims 1-68 have been cancelled, and claims 69-132 have been added.

The specification has been amended to provide a cross-reference to the previously filed International Application.

The specification has also been amended to insert the sequence listing.

Enclosed herewith in full compliance to 37 C.F.R. §§1.821-1.825 is a Substitute Sequence Listing to be inserted into the

specification as indicated above. The Substitute Sequence Listing in no way introduces new matter into the specification.

Also submitted herewith in full compliance to 37 C.F.R. §§1.821-1.825 is a disk copy of the Substitute Sequence Listing. The disk copy of the Substitute Sequence Listing, file "3631-0112P.txt", is identical to the paper copy, except that it lacks formatting.

Entry of the above amendments is earnestly solicited. An early and favorable first action on the merits is earnestly solicited.


Should there be any outstanding matters that need to be resolved in the present application, the Examiner is respectfully requested to contact Leonard R. Svensson (Reg. 30,330) at the telephone number of the undersigned below.

If necessary, the Commissioner is hereby authorized in this, concurrent, and future replies, to charge payment or credit any overpayment to Deposit Account No. 02-2448 for any additional fees required under 37 C.F.R. § 1.16 or under 37 C.F.R. § 1.17; particularly, extension of time fees.

Respectfully submitted,

BIRCH, STEWART, KOLASCH & BIRCH, LLP

By

for  # 47,604  
Leonard R. Svensson, #30,330

P.O. Box 747  
Falls Church, VA 22040-0747  
(714) 708-8555

LRS/lmt  
3631-0112P

METHOD FOR DOWN-REGULATING IL5 ACTIVITY

## FIELD OF THE INVENTION

5 The present invention relates to improvements in therapy and prevention of conditions characterized by an elevated level of eosinophil leukocytes, i.e. conditions such as asthma and other chronic allergic diseases. More specifically, the present invention provides a method for down-regulating inter-

10 leukin 5 (IL5) by enabling the production of antibodies against IL5 thereby reducing the level of activity of eosinophils. The invention also provides for methods of producing modified IL5 useful in this method as well as for the modified IL5 as such. Also encompassed by the present invention are nu-

15 cleic acid fragments encoding modified IL5 as well as vectors incorporating these nucleic acid fragments and host cells and cell lines transformed therewith. The invention also provides for a method for the identification of IL5 analogues which are useful in the method of the invention as well as for compositions comprising modified IL5 or comprising nucleic acids en-

20 coding the IL5 analogues.

## BACKGROUND OF THE INVENTION

25

Asthma is a common disease of the airways, affecting about 10% of the population. The present treatments is primarily based on the administration of steroids and represents a market value exceeding well over a billion dollars. For yet unknown

30 reasons the incidence and morbidity of asthmatics have increased worldwide over the past two decades. Today, an improved understanding of the immunological mechanisms involved in asthmatic conditions combined with an explosive development in biotechnology provides a new basis for the development of

35 alternative and perhaps better strategies for treatment.

A general feature in the pathogenesis of asthma and other chronic allergic diseases has proven to be elevated numbers of

eosinophils, especially in the bronchial mucosa of the lungs. Upon activation eosinophils secrete a number of mediators that are actively involved in the inflammatory airway response. In the activation of eosinophils, interleukin 5 (IL5) plays an important role.

IL5 is a cytokine found in many mammalian species and among others both the human and the murine gene for IL5 have been cloned (Tanabe et al., 1987, Campbell et al., 1988). The human gene consists of four exons with three introns positioned at chromosome 5 and codes for a 134 amino acid residue precursor, including a 19 amino acid N-terminal leader sequence which has the amino acid sequence set forth in SEQ ID NO: 62. Posttranslational cleavage generates the mature 115 amino acid residue protein (SEQ ID NO: 1). The murine IL5 (mIL5) gene similarly codes for a 133 amino acid residue pre-cursor with a 20 amino acid leader sequence which has the amino acid sequence set forth in SEQ ID NO: 64. The processed mature mIL5 is thus 113 amino acid residues long (SEQ ID NO: 12), missing two N-terminal amino acid residues by alignment with human IL5. The amino acid sequences of hIL5 and mIL5 are 70% identical compared to 77% at nucleotide level of the coding regions (Azuma et al., 1986). Higher similarity was reported within human primates; 99% identity is reported for the coding regions of the human and the Rhesus monkey nucleotide sequences (Villinger et al., 1995).

The human amino acid sequence has two potential N-glycosylation sites and the murine three. Human IL5 has been shown to be both N-glycosylated as well as O-glycosylated at Thr 3. Studies of hIL5 has demonstrated that the glycosylation is not necessary for the biological activity even though the stability seems to be affected by de-glycosylation (Tominaga et al., 1990; Kodama et al., 1993).

## Structure of IL5

The active IL5 is a homo-dimer and the 3-dimensional structure of recombinant hIL5 has been determined by X-ray crystallography (Milburn et al., 1993). The 2 monomers are organised in an antiparallel manner and covalently bound by two interchain disulfide bridges (44-87' and 87-44'), thus engaging all 4 cysteines of the 2 monomers.

10 The secondary structure of the monomers consists of 4  $\alpha$ -helices (A-D) intermitted by 3 linking regions (loops) including two short stretches of  $\beta$ -sheets. This 4 $\alpha$  helix bundle is known as the "common cytokine fold", which has also been reported for IL-2, IL-4, GM-CSF, and M-CSF. But all these are monomers and the homodimer-structure in which the D-helix completes the 4 $\alpha$  helix motif of the opposite monomer is unique to IL5.

The native monomers alone has been shown to be biologically inactive (for reviews see Callard & Gearing, 1994; Takutsu et al., 1997). It is nevertheless possible to produce a modified recombinant biologically active monomer by inserting 8 additional amino acid residues in loop 3, connecting the helices C and D. This enables helix D to complete the 4 helix structure within one polypeptide chain and thus enable the monomer to interact with its receptor (Dickason & Huston, 1996; Dickason et al., 1996).

The IL5 receptor is primarily present on eosinophils and it is composed of an  $\alpha$ -chain and a  $\beta$ -chain. The  $\alpha$ -chain of the receptor is specific for IL5 and the  $\beta$ -chain, which assure high-affinity binding and signal transduction, is shared with the hetero-dimer receptors for IL-3 and GM-CSF. The sharing of a receptor component could be the reason for the cross-competition seen between IL5, IL-3 and GM-CSF (for review, see Lopez et al., 1992). However, it was recently demonstrated that the regulation of the IL5R is distinct from the regulation of the IL-3R and the GM-CSFR, further indicating a



highly specialised role of IL5 in the regulation of the eosinophilic response (Wang et al., 1998).

The C-terminal part of IL5 seems to be important in both binding to the IL5R and for the biological activity, since removal of more than two C-terminal amino acid residues results in a decline in both the binding affinity to the IL5 R and in the biological activity in an IL5 bioassay (Proudfoot et al., 1996). Other residues have also been found to be important for binding to the receptor, such as Glu12, which is involved in binding to the  $\beta$ -chain, while the Arg90 and Glu109 residues are involved in the binding to the  $\alpha$ -chain of the receptor. In general, binding to the IL5R seems to occur in regions overlapping helices A and D, where helix D is primarily responsible for the binding to the specific IL5R  $\alpha$ -chain (Graber et al., 1995; Takastu et al., 1997).

#### IL5's homology to other proteins

The two 4-helix domain motifs seen in the homodimer has strikingly similar secondary and tertiary structure as compared to the cytokine fold found in GM-CSF and M-CSF, IL-2, IL-4 and human and porcine growth hormone (Milburn et al., 1993). However, even though striking similarities are also observed in the intron/exon organisation and position of cysteines (Tanabe et al., 1987; Cambell et al., 1988) suggesting a phylogenetic relationship with IL-2, IL-4 and GM-CSF, no significant homology with any of these or other cytokines is observed from the amino acid sequence.

#### Biological activity of IL5

IL5 is mainly secreted by fully differentiated Th2 cells, mast cells and eosinophils (Cousins et al., 1994; Takutsu et al., 1997). It has been shown to act on eosinophils, basophiles, cytotoxic T lymphocytes and on murine B cells (Callard & Gearing, 1994; Takutsu et al., 1997). The effects of IL5 on human B cells are still a matter of controversy. Augmentation of im-

munoglobulin synthesis under certain circumstances and binding to a variety of human B cell lines have been demonstrated. Even though mRNA for the hIL5R has been found in human B-cells, the actual presence of the receptor on these cells has  
5 still to be verified (Baumann & Paul, 1997; Huston et al., 1996).

The actions of IL5 on eosinophils include chemotaxis, enhanced adhesion to endothelial cells, activation and terminal differentiation of the cells. Furthermore it has been demonstrated  
10 that IL5 prevents mature eosinophils from apoptosis (Yamaguchi et al., 1991). These findings have contributed to the present concept of IL5 as being the most important cytokine for eosinophil differentiation (Corrigan & Kay, 1996; Karlen et al.,  
15 1998).

Physiologically, IL5 and its associated eosinophil activation is considered to serve a protective role against helminthic infections and possibly against certain tumours, since these  
20 diseases are typically accompanied by peripheral blood eosinophilia (Takutsu et al., 1997; Sanderson et al., 1992). It is, however, somewhat speculative as in two studies the authors failed to show any effect beside eosinophil down-regulation following administration of antibodies against IL5  
25 on the immunity (e.g. IgE levels) against *Nippostrongylus braziliensis* or *Schistosoma mansoni* in mice infected with these parasites (Sher et al., 1990; Coffman et al., 1989).

#### IL5 transgenic and "knock-out" animals

30

Studies of transgenic mice expressing IL5 or knock-out mice deficient for IL5 have given further knowledge of the physiological role of IL5.

35 Several IL5 transgenic mice have been reported:

A transgenic mouse expressing the IL5 gene in T cells was reported to have an increased white blood cell level character-

5  
rised by expansion of B220+ B lymphocytes and profound eosinophilia. This was accompanied by a massive peritoneal cavity cell exudate dominated by eosinophils and infiltration of eosinophils in nearly all organ systems (Lee et al., 1997a).

5

Another transgenic mouse, expressing the IL5 gene under control of a metallothionin promoter was characterised by an increase in the serum levels of IgM and IgA, a massive eosinophilia in peripheral blood and many other organs accompanied  
10 by the expansion of a distinctive CD5+ B cell population, which produce auto-antibodies (Tominaga et al., 1991).

15 A third study involved a transgenic mouse constitutively expressing IL5 in the lungs. These animals developed pathophysiological changes resembling those of human asthma, including eosinophil invasion of peribronchial spaces, epithelial hypertrophy and increased mucus production. Furthermore, development of airway hyper responsiveness was seen in the absence of antigens (Lee et al., 1997b).

20

IL5-deficient mice ('knock-out' mice) have also been studied. These mice (C57BL/6) have no obvious signs of disease and are fertile. The immunoglobulin levels and the specific antibody responses to DNP-OVA were normal. Basal levels of eosinophils  
25 are produced, but are 2-3 times lower than in control animals, indicating that eosinophils can be produced in the complete absence of IL5. When these mice were infected with *Mesocostoides corti* the eosinophilia normally seen was abolished and this absence of eosinophilia did not affect the worm burden  
30 produced by this parasite (Kopf et al., 1996).

In a study by Foster et al. (1996), the effect of IL5 knock-out on a common model of atopic airway inflammation was investigated. Sensitisation and aerosol challenge of mice with  
35 ovalbumin normally result in airway eosinophilia, airway hyperreactivity to  $\beta$ -methacholin and extensive lung damage analogous to that seen in asthma. In the IL5 deficient mice the eosinophilia, airway hyperreactivity and lung damage were

abolished. When IL5 expression in these mice was reconstituted, the aero-allergen induced eosinophilia and airway dysfunction were restored.

## 5 Pathophysiologic role of IL5

Asthma affect about 10% of the population worldwide and for yet unknown reasons the incidence and morbidity have increased over the past two decades (Ortega & Busse, 1997). It is a chronic airway disease characterised by recurrent and usually reversible air flow obstruction, inflammation and hyper responsiveness (Moxam and Costello, 1990). This produces symptoms of wheezing and breathlessness, which in severe cases can be fatal.

The animal experiments referred to above using transgenic mice constitutively expressing IL5 in the lungs (Lee et al., 1997a) and the IL5 deficient "knock-out" mice (Foster et al., 1996) strongly implicate a crucial role of IL5 in the pathogenesis of asthma. Further evidence supporting this can be deduced from several studies including asthmatic individuals.

Eosinophilia has been identified in bronchoalveolar lavage (BAL) fluid and in bronchial mucosal biopsies of subjects with asthma and correlates with disease severity. Several eosinophil products have been identified in the BAL fluid of patients with asthma and numbers of peripheral blood eosinophils correlate with asthma severity (Ortega & Busse 1997).

IL5 serum concentration was found to be elevated (median concentration 150 pg/ml) in 15 out of 29 patients with chronic severe asthma as compared to control subjects (Alexander et al., 1994).

In another study involving both non-atopic and atopic asthmatics, it was found that an enhanced IL5 production by helper T cells seems to cause the eosinophilic inflammation of both atopic and non-atopic asthma (Mori et al., 1997).

Other results also indicate that IL5 has a distinct role in other atopic diseases. Allergen induced systemic episodes in individuals with allergic rhinitis has recently been shown to correlate to allergen induced IL5 synthesis rather than IgE (Ohashi et al., 1998). The correlation of atopic reactions is also demonstrated in a study by Barata et al. (1998) in which a significant expression of IL5 by T-cells in a cutaneous late phase reaction is demonstrated.

10

These and other results have led several authors as Corrigan & Kay (1996), Danzig & Cuss (1997) to identify and recommend IL5 as a primary target in the development of a better treatment for asthma and atopic diseases involving eosinophilic inflammation. Chronic tissue damaging hypereosinophilia induced by parasitic infection, topical pulmonary eosinophilia and hypereosinophilic syndrome are examples of other pathogenic conditions that could be addressed by IL5 down regulation.

## 20 *In vivo* demonstration of the role of IL5

In several studies with rodent models of asthma it has been shown that treatment with monoclonal antibodies against IL5 (anti-IL5 mAb) results in dose-related inhibition of eosinophilia, as compared to non-treated controls (Nagai et al., 1993a & b; Chand et al., 1992; Coeffier et al., 1994; Kung et al., 1995; Underwood et al., 1996). In the study by Nagai et al. (1993a) the effect was also observed by treating the sensitised Balb/c mice with soluble IL5 receptor  $\alpha$ .

30

In one study with Balb/c mice (Hamelmann et al., 1997) and four studies with guinea pigs it was additionally shown that anti-IL5 mAb could inhibit airway hyperreactivity elicited with various substances in antigen sensitised animals (Mauser et al., 1993; Akutsu et al., 1995; van Oosterhout et al., 1995 & 1993). In some of the studies beneficial effects (cf. table 1) of the anti-IL5 mAb treatment were also observed microscopically (Mauser et al., 1993; Akutsu et al., 1995; Kung et

al., 1995). Importantly, in the study by Kung et al. (1995) a reduction of pulmonary inflammation in B6D2F1 mice was seen both when anti-IL5 mAb was administered hours before antigen challenge and also when administered up to five days after antigen challenge, indicating that the effect of anti-IL5 mAb may be both prophylactic and therapeutic for airway inflammation. This effect, however, was not observed by Underwood et al. when guinea pigs were given anti-IL5 mAb two hours after antigen challenge (Underwood et al., 1996).

In a study using a monkey model of asthma, Mauser et al. (1995) reported an inhibition of airway hyper reactivity after antigen challenge, when rat anti mouse-IL5 mAb was given 1 hour before antigen challenge. In addition, there was 75% reduction in the number of eosinophils in bronchoalveolar lavage (BAL) of antibody treated animals, as compared to non-treated controls. The effects on eosinophilia and hyperresponsiveness of anti-IL5 mAb was seen for up to three months after treatment (Mauser et al., 1995). Regarding allergic hyperresponsiveness, the results from studies by Nagai et al. (1993a and 1993b) document no reduction in hyperresponsiveness in conjunction to a reduction of eosinophil numbers in BAL.

All anti-IL5 mAb *in vivo* experiments mentioned so far have been done with rat-anti-mouse monoclonal antibodies. Egan et al. (1995) have reported experiments using humanised rat-anti-human IL5 monoclonal antibodies, called Sch 55700. These mAbs, inhibited lung lavage eosinophilia by 75% at a dose of 0,3 mg/kg when administered to sensitised monkeys. When Sch 55700 was given at 1 mg/kg in allergic mice, inhibition of airway eosinophilia was also observed.

#### Treatment of asthma at present and in the future

The current treatment of asthma is, as mentioned, corticosteroids which, by their anti-inflammatory action, are the most powerful drugs. Besides this,  $\beta_2$  agonists and methyl xanthine derivatives which all cause bronchodilation, and disodium

chromoglycate which 'stabilises' mast cells, thereby preventing mediator release, all have proven beneficial in asthma patients (Ortega & Busse 1997).

- 5 Future treatment of asthma may as discussed above include anti-IL5 mAbs. Celltech in corporation with Schering Plough have anti-IL5 mAb in phase I clinical trial for treatment of asthma. However, treatment with monoclonal antibodies entails a number of drawbacks. First of all, the development and pro-  
10 duction costs for a safe mAB (e.g. a humanised mAB) are very high, resulting in an expensive therapeutic product for the end user. Second, mABs have the disadvantageous characteristic seen from a patient point of view that they have to be administered with relatively short intervals. Third, by nature  
15 mABs exhibit a narrow specificity against one single epitope of the antigen. And, finally, mABs (even humanised) are immunogenic, leading to an increasingly fast inactivation of administered antibodies as treatment progresses over time.
- 20 Also use of antisense IL5 oligonucleotides for antisense therapy has been suggested by the company Hybridon for the treatment of asthma, allergies and inflammation. However, the antisense technology has proven to be technically difficult and, in fact, conclusive evidence of the feasibility of antisense  
25 therapy in humans has not yet been established.

Finally, WO 97/45448 (Bresagen Limited / Medvet Science) proposes the use of "modified and variant forms of IL5 molecules capable of antagonising the activity of IL5" in ameliorating,  
30 abating or otherwise reducing the aberrant effects caused by native or mutant forms of IL5. The antagonizing effect is reported to be the result of the variant forms of IL5 binding to the low affinity  $\alpha$  chain of IL5R but not to the high affinity receptors; in this way the variants compete with IL5 for binding to its receptors without exerting the physiological effects of IL5.  
35

Other atopic diseases involving eosinophilic inflammation are treated with either the symptomatica mentioned for asthma or immune therapy (IT) using hyposensitization with allergen extracts. The latter type of treatment is known to be effective  
5 against allergies against one or a few antigens, whereas IT is not feasible in the treatment of multiple allergies. Furthermore, the time scale for obtaining clinical improvement in patients susceptible to treatment is very long for conventional IT.

10 Thus, in spite of existing and possible future therapies for chronic allergic diseases such as asthma, there is a definite need for alternative ways of treating and ameliorating this and other chronic allergic diseases.

#### 15 OBJECT OF THE INVENTION

The object of the present invention is to provide novel therapies  
20 against chronic allergic conditions (such as asthma) characterized by eosinophilia. A further object is to develop an autovaccine against IL5, in order to obtain a novel treatment for asthma and for other pathological disorders involving chronic airway inflammation.

#### 25 SUMMARY OF THE INVENTION

The T-cell derived cytokine IL5 has, as mentioned above, a  
30 crucial role in orchestrating the eosinophilic response, affecting both the production, the localisation and the activation of eosinophils. As IL5 has not otherwise been reported to have a central role in the development of a protective immune response, this particular cytokine is in the opinion of the  
35 inventors an attractive therapeutic target for the treatment of asthma.



The general aim according to the present invention is to decrease the pathogenic levels of eosinophils in the airways of the asthma patient by down-regulating of the IL5 levels, since eosinophils depend on IL5 for attraction and activation. The result of a decreased eosinophil number in the airway mucosa would be a concomitant decrease in the airway inflammation, corresponding to a clinical improvement in the asthmatic patient.

10 The potential effect of such an approach has already been demonstrated in studies using anti IL5 monoclonal antibodies in animal models of airway inflammation, cf. the "PREAMBLE TO EX-AMPLES".

15 This current invention, however, takes the results obtained through passive immunisation one step further by using the approach of generating an active immune response through the concept of autovaccination. To the best of the inventor's knowledge, such an approach has never been suggested before.

20 The advantage of treating asthmatics with an IL5 autovaccine, as compared to current treatment with corticosteroids etc., is a reduction and/or elimination of side effects and most likely a better effect in terms of duration. When compared to anti-IL5 mAbs, the effect of an induced polyclonal Ab response is expected to be superior to passively injected monoclonal immunoglobulins since the polyclonal response has a broader specificity. Improvements with respect to administration regimens are also expected (since effective autovaccines described herein typically would require a maximum of 2-6 administrations per year).

When compared to hyposensitization, the present invention offers the attractive aspect of being non-specific; this is especially relevant when dealing with multi-allergic patients.

Thus, in its broadest and most general scope, the present invention relates to a method for *in vivo* down-regulation of in-

terleukin 5 (IL5) activity in an animal, including a human being, the method comprising effecting presentation to the animal's immune system of an immunologically effective amount of

- at least one IL5 polypeptide or subsequence thereof which has been formulated so that immunization of the animal with the IL5 polypeptide or subsequence thereof induces production of antibodies against the IL5 polypeptide, and/or

- at least one IL5 analogue wherein is introduced at least one modification in the IL5 amino acid sequence which has as a result that immunization of the animal with the analogue induces production of antibodies against the IL5 polypeptide.

The most attractive aspect of this approach is that e.g. asthma can be controlled by periodic but not very frequent immunizations, in contrast to a therapeutic approach which involves administration of anti-IL5 or molecules having a binding affinity to IL5 analogous therewith. It is expected that 1-4 annual injections with an immunogenic composition according to the invention will be sufficient to obtain the desired effect, whereas administration of other inhibitors of IL5 activity does or will require daily, or at least weekly, administrations.

The invention also relates to IL5 analogues as well as to nucleic acid fragments encoding a subset of these. Also immunogenic compositions comprising the analogues or the nucleic acid fragments are part of the invention.

The invention also relates to a method of identifying analogues of IL5 as well as a method for preparing a composition comprising the IL5 analogues.

## LEGENDS TO THE FIGURES

Fig. 1: The amino acid sequence of the mature human IL5 (SEQ ID NO: 1). The aligned murine sequence is included (SEQ ID NO: 12), but only positions that differ from the human sequence are displayed. The two "\*"s indicate the missing N-terminal residues of the murine IL5. The N-glycosylation positions are marked with double underlining, the O-glycosylated threonines of human IL5 are given in italics, and the cysteines in bold.

Fig. 2: The dimer and monomer structures of human IL5. A: Dimer structure of hIL5. The structure has only been obtained for residues 5-112, which means that the O-glycosylation site at Thr3 is not included. B: The same structure as in A, with the assignment of the helices (A-D and A'-D'). C: The monomer hIL5 with the amino acid residues differing from the mIL5 shown in light grey.

Fig. 3: The aligned mature human IL5 (hIL5) and murine IL5 (mIL5) amino acid sequences (SEQ ID NOs: 1 and 12) with indications of suitable substitution regions. The 4  $\alpha$ -helices A-D are surrounded by solid-line boxes, the  $\beta$ -sheets are double underlined and the positions of the two cysteines are marked with "▼". Identical residues in the two sequences are marked with "-" and non-identical residues with "\*". Loop 1 spans between helices A and B, Loop 2 spans between helices B and C, and loop 3 spans between loops C and D. Amino acid sequences to be substituted with foreign T<sub>H</sub> epitope containing peptides are marked in bold; one such sequence is surrounded by a dot-lined box because of residues overlapping with those substituted in a different construct. The amino acid sequences of 10 constructs (5 derived from human and

5 derived from murine IL5) are set forth in SEQ ID  
NOs: 2-11 and 13-22.

Fig. 4: ELISA results of DNA immunization testing two mIL5

autovaccine DNA vaccines.

Mice were DNA vaccinated with naked plasmid DNA encoding either ovalbumin, mIL5wt, mIL5.1 or mIL5.5. Sera obtained at day 77 were tested for reactivity against ovalbumin and murine IL5. Polystyrene micro-titer plates (Maxisorp, Nunc) were coated with ovalbumin (1 µg/well, Sigma) or purified recombinant murine IL5 (0.1 µg/well, E1320). The reactivities of diluted sera added to the wells were visualised using a goat anti-mouse secondary antibody. OD490 readings of the pre-bleeds were subtracted from the OD490 readings of the test samples, and the resulting values were presented for each individual mouse as bars. The OD490 readings of the pre-bleeds (in 1:25 dilution) were ranging from 0.025-0.034. Crucifixes indicate dead animals.

Fig. 5: Schematic representation of murine IL5 based autovaccine constructs.

The top figure represents murine wild-type IL5 monomer with helices A-C, loops 1-3 and the flexible C-terminal region. Remaining figures represent different autovaccine constructs having in-substitutions of the tetanus toxoid epitopes P2 and P30 in various positions. Specific constructs are detailed in the Examples.

## DETAILED DISCLOSURE OF THE INVENTION

Definitions

- 5 In the following, a number of terms used in the present specification and claims will be defined and explained in detail in order to clarify the metes and bounds of the invention.

The terms "T-lymphocyte" and "T-cell" will be used inter-  
10 changeably for lymphocytes of thymic origin which are responsible for various cell mediated immune responses as well as for helper activity in the humeral immune response. Likewise, the terms "B-lymphocyte" and "B-cell" will be used interchangeably for antibody-producing lymphocytes.

15 An "IL5 polypeptide" is herein intended to denote polypeptides having the amino acid sequence of the above-discussed IL5 proteins derived from humans and mice (or truncates thereof sharing a substantial amount of B-cell epitopes with intact IL5),  
20 but also polypeptides having the amino acid sequence identical to xeno-analogues of these two proteins isolated from other species are embraced by the term. Also unglycosylated forms of IL5 which are prepared in prokaryotic system are included within the boundaries of the term as are forms having varying  
25 glycosylation patterns due to the use of e.g. yeasts or other non-mammalian eukaryotic expression systems. It should, however, be noted that when using the term "an IL5 polypeptide" it is intended that the polypeptide in question is normally non-immunogenic when presented to the animal to be treated. In  
30 other words, the IL5 polypeptide is a self-protein or is a xeno-analogue of such a self-protein which will not normally give rise to an immune response against IL5 of the animal in question.

- 35 An "IL5 analogue" is an IL5 polypeptide which has been subjected to changes in its primary structure. Such a change can e.g. be in the form of fusion of an IL5 polypeptide to a suitable fusion partner (i.e. a change in primary structure exclu-

sively involving C- and/or N-terminal additions of amino acid residues) and/or it can be in the form of insertions and/or deletions and/or substitutions in the IL5 polypeptide's amino acid sequence. Also encompassed by the term are derivatized  
5 IL5 molecules, cf. the discussion below of modifications of IL5.

It should be noted that the use as a vaccine in a human of e.g. a canine analogue of human IL5 can be imagined to produce  
10 the desired immunity against IL5. Such use of an xeno-analogue for immunization is also considered to be an "IL5 analogue" as defined above.

When using the abbreviation "IL5" herein, this is intended as  
15 a reference to the amino acid sequence of mature, wildtype IL5 (also denoted "IL5m" and "IL5wt" herein). Mature human IL5 is denoted hIL5, hIL5m or hIL5wt, and murine mature IL5 is denoted mIL5, mIL5m, or mIL5wt. In cases where a DNA construct includes information encoding a leader sequence or other mate-  
20 rial, this will normally be clear from the context.

The term "polypeptide" is in the present context intended to mean both short peptides of from 2 to 10 amino acid residues, oligopeptides of from 11 to 100 amino acid residues, and poly-  
25 peptides of more than 100 amino acid residues. Furthermore, the term is also intended to include proteins, i.e. functional biomolecules comprising at least one polypeptide; when comprising at least two polypeptides, these may form complexes, be covalently linked, or may be non-covalently linked. The  
30 polypeptide(s) in a protein can be glycosylated and/or lipi- dated and/or comprise prosthetic groups.

The term "subsequence" means any consecutive stretch of at least 3 amino acids or, when relevant, of at least 3 nucleo-  
35 tides, derived directly from a naturally occurring IL5 amino acid sequence or nucleic acid sequence, respectively.

The term "animal" is in the present context in general intended to denote an animal species (preferably mammalian), such as *Homo sapiens*, *Canis domesticus*, etc. and not just one single animal. However, the term also denotes a population of such an animal species, since it is important that the individuals immunized according to the method of the invention all harbour substantially the same IL5 allowing for immunization of the animals with the same immunogen(s). If, for instance, genetic variants of IL5 exists in different human population it may be necessary to use different immunogens in these different populations in order to be able to break the autotolerance towards IL5 in each population. It will be clear to the skilled person that an animal in the present context is a living being which has an immune system. It is preferred that the animal is a vertebrate, such as a mammal.

By the term "in vivo down-regulation of IL5 activity" is herein meant reduction in the living organism of the number of interactions between IL5 and its receptors (or between IL5 and other possible biologically important binding partners for this molecule). The down-regulation can be obtained by means of several mechanisms: Of these, simple interference with the active site in IL5 by antibody binding is the most simple. However, it is also within the scope of the present invention that the antibody binding results in removal of IL5 by scavenger cells (such as macrophages and other phagocytic cells).

The expression "effecting presentation ... to the immune system" is intended to denote that the animal's immune system is subjected to an immunogenic challenge in a controlled manner. As will appear from the disclosure below, such challenge of the immune system can be effected in a number of ways of which the most important are vaccination with polypeptide containing "pharmaccines" (i.e. a vaccine which is administered to treat or ameliorate ongoing disease) or nucleic acid "pharmaccine" vaccination. The important result to achieve is that immune competent cells in the animal are confronted with the antigen in an immunologically effective manner, whereas the precise

mode of achieving this result is of less importance to the inventive idea underlying the present invention.

The term "immunogenically effective amount" has its usual meaning in the art, i.e. an amount of an immunogen which is capable of inducing an immune response which significantly engages pathogenic agents which share immunological features with the immunogen.

10 When using the expression that the IL5 has been "modified" is herein meant a chemical modification of the polypeptide which constitutes the backbone of IL5. Such a modification can e.g. be derivatization (e.g. alkylation, acylation, esterification etc.) of certain amino acid residues in the IL5 sequence, but  
15 as will be appreciated from the disclosure below, the preferred modifications comprise changes of (or additions to) the primary structure of the IL5 amino acid sequence.

When discussing "autotolerance towards IL5" it is understood  
20 that since IL5 is a self-protein in the population to be vaccinated, normal individuals in the population do not mount an immune response against IL5; it cannot be excluded, though, that occasional individuals in an animal population might be able to produce antibodies against native IL5, e.g. as part of  
25 an autoimmune disorder. At any rate, an animal will normally only be autotolerant towards its own IL5, but it cannot be excluded that IL5 analogues derived from other animal species or from a population having a different IL5 phenotype would also be tolerated by said animal.

30

A "foreign T-cell epitope" (or: "foreign T-lymphocyte epitope") is a peptide which is able to bind to an MHC molecule and which stimulates T-cells in an animal species. Preferred foreign T-cell epitopes in the invention are "promiscuous"  
35 epitopes, i.e. epitopes which bind to a substantial fraction of a particular class of MHC molecules in an animal species or population. Only a very limited number of such promiscuous T-cell epitopes are known, and they will be discussed in detail



below. It should be noted that in order for the immunogens which are used according to the present invention to be effective in as large a fraction of an animal population as possible, it may be necessary to 1) insert several foreign T-cell epitopes in the same IL5 analogue or 2) prepare several IL5 analogues wherein each analogue has a different promiscuous epitope inserted. It should be noted also that the concept of foreign T-cell epitopes also encompasses use of cryptic T-cell epitopes, i.e. epitopes which are derived from a self-protein and which only exerts immunogenic behaviour when existing in isolated form without being part of the self-protein in question.

A "foreign T helper lymphocyte epitope" (a foreign  $T_H$  epitope) is a foreign T cell epitope which binds an MHC Class II molecule and can be presented on the surface of an antigen presenting cell (APC) bound to the MHC Class II molecule.

A "functional part" of a (bio)molecule is in the present context intended to mean the part of the molecule which is responsible for at least one of the biochemical or physiological effects exerted by the molecule. It is well-known in the art that many enzymes and other effector molecules have an active site which is responsible for the effects exerted by the molecule in question. Other parts of the molecule may serve a stabilizing or solubility enhancing purpose and can therefore be left out if these purposes are not of relevance in the context of a certain embodiment of the present invention. For instance it is possible to use certain other cytokines as a modifying moiety in IL5 (cf. the detailed discussion below), and in such a case, the issue of stability may be irrelevant since the coupling to IL5 provides the stability necessary.

The term "adjuvant" has its usual meaning in the art of vaccine technology, i.e. a substance or a composition of matter which is 1) not in itself capable of mounting a specific immune response against the immunogen of the vaccine, but which is 2) nevertheless capable of enhancing the immune response

against the immunogen. Or, in other words, vaccination with the adjuvant alone does not provide an immune response against the immunogen, vaccination with the immunogen may or may not give rise to an immune response against the immunogen, but the combination of vaccination with immunogen and adjuvant induces an immune response against the immunogen which is stronger than that induced by the immunogen alone.

"Targeting" of a molecule is in the present context intended to denote the situation where a molecule upon introduction in the animal will appear preferentially in certain tissue(s) or will be preferentially associated with certain cells or cell types. The effect can be accomplished in a number of ways including formulation of the molecule in composition facilitating targeting or by introduction in the molecule of groups which facilitates targeting. These issues will be discussed in detail below.

"Stimulation of the immune system" means that a substance or composition of matter exhibits a general, non-specific immunostimulatory effect. A number of adjuvants and putative adjuvants (such as certain cytokines) share the ability to stimulate the immune system. The result of using an immunostimulating agent is an increased "alertness" of the immune system meaning that simultaneous or subsequent immunization with an immunogen induces a significantly more effective immune response compared to isolated use of the immunogen

#### Preferred embodiments of IL5 activity down-regulation

30

It is preferred that the IL5 polypeptide used as an immunogen in the method of the invention is a modified molecule wherein at least one change is present in the IL5 amino acid sequence, since the chances of obtaining the all-important breaking of autotolerance towards IL5 is greatly facilitated that way. It should be noted that this does not exclude the possibility of using such a modified IL5 in formulations which further facilitate the breaking of autotolerance against IL5, e.g. for-

mulations containing certain adjuvants discussed in detail below.

It has been shown (in Dalum I et al., 1996, J. Immunol. 157: 4796-4804) that potentially self-reactive B-lymphocytes recognizing self-proteins are physiologically present in normal individuals. However, in order for these B-lymphocytes to be induced to actually produce antibodies reactive with the relevant self-proteins, assistance is needed from cytokine producing T-helper lymphocytes ( $T_H$ -cells or  $T_H$ -lymphocytes). Normally this help is not provided because T-lymphocytes in general do not recognize T-cell epitopes derived from self-proteins when presented by antigen presenting cells (APCs). However, by providing an element of "foreignness" in a self-protein (i.e. by introducing an immunologically significant modification), T-cells recognizing the foreign element are activated upon recognizing the foreign epitope on an APC (such as, initially, a mononuclear cell). Polyclonal B-lymphocytes (which are also specialised APCs) capable of recognising self-epitopes on the modified self-protein also internalise the antigen and subsequently presents the foreign T-cell epitope(s) thereof, and the activated T-lymphocytes subsequently provide cytokine help to these self-reactive polyclonal B-lymphocytes. Since the antibodies produced by these polyclonal B-lymphocytes are reactive with different epitopes on the modified polypeptide, including those which are also present in the native polypeptide, an antibody cross-reactive with the non-modified self-protein is induced. In conclusion, the T-lymphocytes can be led to act as if the population of polyclonal B-lymphocytes have recognised an entirely foreign antigen, whereas in fact only the inserted epitope(s) is/are foreign to the host. In this way, antibodies capable of cross-reacting with non-modified self-antigens are induced.

Several ways of modifying a peptide self-antigen in order to obtain breaking of autotolerance are known in the art. Hence, according to the invention, the modification can include that

- at least one foreign T-cell epitope is introduced, and/or
- at least one first moiety is introduced which effects targeting of the modified molecule to an antigen presenting cell (APC), and/or
- 5 - at least one second moiety is introduced which stimulates the immune system, and/or
- at least one third moiety is introduced which optimises presentation of the modified IL5 polypeptide to the immune system.

10

However, all these modifications should be carried out while maintaining a substantial fraction of the original B-lymphocyte epitopes in IL5, since the B-lymphocyte recognition of the native molecule is thereby enhanced.

15

In one preferred embodiment, side groups (in the form of foreign T-cell epitopes or the above-mentioned first, second and third moieties) are covalently or non-covalently introduced. This is intended to mean that stretches of amino acid residues derived from IL5 are derivatized without altering the primary amino acid sequence, or at least without introducing changes in the peptide bonds between the individual amino acids in the chain.

20

- 25 An alternative, and preferred, embodiment utilises amino acid substitution and/or deletion and/or insertion and/or addition (which may be effected by recombinant means or by means of peptide synthesis; modifications which involves longer stretches of amino acids can give rise to fusion polypep-
- 30 tides). One especially preferred version of this embodiment is the technique described in WO 95/05849, which discloses a method for down-regulating self-proteins by immunising with analogues of the self-proteins wherein a number of amino acid sequence(s) has been substituted with a corresponding number
- 35 of amino acid sequence(s) which each comprise a foreign immunodominant T-cell epitope, while at the same time maintaining the overall tertiary structure of the self-protein in the analogue. For the purposes of the present invention, it is how-

ever sufficient if the modification (be it an amino acid insertion, addition, deletion or substitution) gives rise to a foreign T-cell epitope and at the same time preserves a substantial number of the B-cell epitopes in IL5. However, in order to obtain maximum efficacy of the immune response induced, it is preferred that the overall tertiary structure of IL5 is maintained in the modified molecule.

The following formula describes the IL5 constructs generally covered by the invention:

$$(\text{MOD}_1)_{s_1} (\text{IL5}_{e_1})_{n_1} (\text{MOD}_2)_{s_2} (\text{IL5}_{e_2})_{n_2} \dots (\text{MOD}_x)_{s_x} (\text{IL5}_{e_x})_{n_x} \quad (\text{I})$$

-where IL5<sub>e1</sub>-IL5<sub>ex</sub> are x B-cell epitope containing subsequences of IL5 which independently are identical or non-identical and which may contain or not contain foreign side groups, x is an integer  $\geq 3$ , n<sub>1</sub>-n<sub>x</sub> are x integers  $\geq 0$  (at least one is  $\geq 1$ ), MOD<sub>1</sub>-MOD<sub>x</sub> are x modifications introduced between the preserved B-cell epitopes, and s<sub>1</sub>-s<sub>x</sub> are x integers  $\geq 0$  (at least one is  $\geq 1$  if no side groups are introduced in the IL5<sub>e</sub> sequences). Thus, given the general functional restraints on the immunogenicity of the constructs, the invention allows for all kinds of permutations of the original IL5 sequence, and all kinds of modifications therein. Thus, included in the invention are modified IL5 obtained by omission of parts of the IL5 sequence which e.g. exhibit adverse effects *in vivo* or omission of parts which could give rise to undesired immunological reactions.

Maintenance of a substantial fraction of B-cell epitopes or even the overall tertiary structure of a protein which is subjected to modification as described herein can be achieved in several ways. One is simply to prepare a polyclonal antiserum directed against IL5 (e.g. an antiserum prepared in a rabbit) and thereafter use this antiserum as a test reagent (e.g. in a competitive ELISA) against the modified proteins which are produced. Modified versions (analogues) which react to the same extent with the antiserum as does IL5 must be regarded as

having the same overall tertiary structure as IL5 whereas analogues exhibiting a limited (but still significant and specific) reactivity with such an antiserum are regarded as having maintained a substantial fraction of the original B-cell epitopes.

Alternatively, a selection of monoclonal antibodies reactive with distinct epitopes on IL5 can be prepared and used as a test panel. This approach has the advantage of allowing 1) an epitope mapping of IL5 and 2) a mapping of the epitopes which are maintained in the analogues prepared.

Of course, a third approach would be to resolve the 3-dimensional structure of IL5 or of a biologically active truncate thereof (cf. above) and compare this to the resolved three-dimensional structure of the analogues prepared. Three-dimensional structure can be resolved by the aid of X-ray diffraction studies and NMR-spectroscopy. Further information relating to the tertiary structure can to some extent be obtained from circular dichroism studies which have the advantage of merely requiring the polypeptide in pure form (whereas X-ray diffraction requires the provision of crystallized polypeptide and NMR requires the provision of isotopic variants of the polypeptide) in order to provide useful information about the tertiary structure of a given molecule. However, ultimately X-ray diffraction and/or NMR are necessary to obtain conclusive data since circular dichroism can only provide indirect evidence of correct 3-dimensional structure via information of secondary structure elements.

30

One preferred embodiment of the invention utilises multiple presentations of B-lymphocyte epitopes of IL5 (i.e. formula I wherein at least one B-cell epitope is present in two positions). This effect can be achieved in various ways, e.g. by simply preparing fusion polypeptides comprising the structure  $(IL5)_m$ , where  $m$  is an integer  $\geq 2$  and then introduce the modifications discussed herein in at least one of the IL5 sequences, or alternatively, inserted between at least two of

the IL5 amino acid sequences. It is preferred that the modifications introduced includes at least one duplication of a B-lymphocyte epitope and/or the introduction of a hapten.

5 As mentioned above, the introduction of a foreign T-cell epitope can be accomplished by introduction of at least one amino acid insertion, addition, deletion, or substitution. Of course, the normal situation will be the introduction of more than one change in the amino acid sequence (e.g. insertion of  
10 or substitution by a complete T-cell epitope) but the important goal to reach is that the IL5 analogue, when processed by an antigen presenting cell (APC), will give rise to such a foreign immunodominant T-cell epitope being presented in context of an MCH Class II molecule on the  
15 surface of the APC. Thus, if the IL5 amino acid sequence in appropriate positions comprises a number of amino acid residues which can also be found in a foreign  $T_H$  epitope then the introduction of a foreign  $T_H$  epitope can be accomplished by providing the remaining amino acids of the foreign epitope by  
20 means of amino acid insertion, addition, deletion and substitution. In other words, it is not necessary to introduce a complete  $T_H$  epitope by insertion or substitution.

It is preferred that the number of amino acid insertions, deletions, substitutions or additions is at least 2, such as 3,  
25 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, and 25 insertions, substitutions, additions or deletions. It is furthermore preferred that the number of amino acid insertions, substitutions, additions or deletions is not in excess  
30 of 150, such as at most 100, at most 90, at most 80, and at most 70. It is especially preferred that the number of substitutions, insertions, deletions, or additions does not exceed 60, and in particular the number should not exceed 50 or even 40. Most preferred is a number of not more than 30. With re-  
35 spect to amino acid additions, it should be noted that these, when the resulting construct is in the form of a fusion polypeptide, is often considerably higher than 150.

Preferred embodiments of the invention includes modification by introducing at least one foreign immunodominant T<sub>H</sub> epitope. It will be understood that the question of immune dominance of a T<sub>H</sub> epitope depends on the animal species in question. As used  
 5 herein, the term "immunodominance" simply refers to epitopes which in the vaccinated individual gives rise to a significant immune response, but it is a well-known fact that a T<sub>H</sub> epitope which is immunodominant in one individual is not necessarily immunodominant in another individual of the same species, even  
 10 though it may be capable of binding MHC-II molecules in the latter individual.

Another important point is the issue of MHC restriction of T<sub>H</sub> epitopes. In general, naturally occurring T<sub>H</sub> epitopes are MHC  
 15 restricted, i.e. a certain peptide constituting a T<sub>H</sub> epitope will only bind effectively to a subset of MHC Class II molecules. This in turn has the effect that in most cases the use of one specific T<sub>H</sub> epitope will result in a vaccine component which is effective in a fraction of the population only, and  
 20 depending on the size of that fraction, it can be necessary to include more T<sub>H</sub> epitopes in the same molecule, or alternatively prepare a multi-component vaccine wherein the components are IL5 variants which are distinguished from each other by the nature of the T<sub>H</sub> epitope introduced.

25

If the MHC restriction of the T-cells used is completely unknown (for instance in a situation where the vaccinated animal has a poorly defined MHC composition), the fraction of the animal population covered by a specific vaccine composition  
 30 can be determined by means of the following formula:

$$f_{\text{population}} = 1 - \prod_{i=1}^n (1 - p_i) \quad (\text{II})$$

-where  $p_i$  is the frequency in the population of responders to  
 35 the  $i^{\text{th}}$  foreign T-cell epitope present in the vaccine composition, and  $n$  is the total number of foreign T-cell epitopes in the vaccine composition. Thus, a vaccine composition contain-



ing 3 foreign T-cell epitopes having response frequencies in the population of 0.8, 0.7, and 0.6, respectively, would give

$$1 - 0.2 \times 0.3 \times 0.4 = 0.976$$

-i.e. 97.6 percent of the population will statistically mount an MHC-II mediated response to the vaccine.

The above formula does not apply in situations where a more or less precise MHC restriction pattern of the peptides used is known. If, for instance a certain peptide only binds the human MHC-II molecules encoded by HLA-DR alleles DR1, DR3, DR5, and DR7, then the use of this peptide together with another peptide which binds the remaining MHC-II molecules encoded by HLA-DR alleles will accomplish 100% coverage in the population in question. Likewise, if the second peptide only binds DR3 and DR5, the addition of this peptide will not increase the coverage at all. If one bases the calculation of population response purely on MHC restriction of T-cell epitopes in the vaccine, the fraction of the population covered by a specific vaccine composition can be determined by means of the following formula:

$$f_{\text{population}} = 1 - \prod_{j=1}^3 (1 - \phi_j)^2 \quad (\text{III})$$

-wherein  $\phi_j$  is the sum of frequencies in the population of allelic haplotypes encoding MHC molecules which bind any one of the T-cell epitopes in the vaccine and which belong to the  $j^{\text{th}}$  of the 3 known HLA loci (DP, DR and DQ); in practice, it is first determined which MHC molecules will recognize each T-cell epitope in the vaccine and thereafter these MHC molecules are listed by type (DP, DR and DQ) - then, the individual frequencies of the different listed allelic haplotypes are summed for each type, thereby yielding  $\phi_1$ ,  $\phi_2$ , and  $\phi_3$ .

It may occur that the value  $p_i$  in formula II exceeds the corresponding theoretical value  $\pi_i$ :

$$\pi_i = 1 - \prod_{j=1}^3 (1 - v_j)^2 \quad (IV)$$

-wherein  $v_j$  is the sum of frequencies in the population of allelic haplotypes encoding MHC molecules which bind the  $i^{\text{th}}$  T-cell epitope in the vaccine and which belong to the  $j^{\text{th}}$  of the 3 known HLA loci (DP, DR and DQ). This means that in  $1 - \pi_i$  of the population there is a frequency of responders of  $f_{\text{residual}_i} = (p_i - \pi_i) / (1 - \pi_i)$ . Therefore, formula III can be adjusted so as to yield formula V:

$$f_{\text{population}} = 1 - \prod_{j=1}^3 (1 - \phi_j)^2 + \left( 1 - \prod_{i=1}^n (1 - f_{\text{residual}_i}) \right) \quad (V)$$

-where the term  $1 - f_{\text{residual}_i}$  is set to zero if negative. It should be noted that formula V requires that all epitopes have been haplotype mapped against identical sets of haplotypes.

Therefore, when selecting T-cell epitopes to be introduced in the IL5 analogue, it is important to include all knowledge of the epitopes which is available: 1) The frequency of responders in the population to each epitope, 2) MHC restriction data, and 3) frequency in the population of the relevant haplotypes.

There exists a number of naturally occurring "promiscuous" T-cell epitopes which are active in a large proportion of individuals of an animal species or an animal population and these are preferably introduced in the vaccine, thereby reducing the need for a very large number of different IL5 analogues in the same vaccine.

The promiscuous epitope can according to the invention be a naturally occurring human T-cell epitope such as epitopes from tetanus toxoid (e.g. the P2 and P30 epitopes), diphtheria toxoid, Influenza virus hemagglutinin (HA), and *P. falciparum* CS antigen.

Over the years a number of other promiscuous T-cell epitopes have been identified. Especially peptides capable of binding a

large proportion of HLA-DR molecules encoded by the different HLA-DR alleles have been identified and these are all possible T-cell epitopes to be introduced in the IL5 analogues used according to the present invention. Cf. also the epitopes discussed in the following references which are hereby all incorporated by reference herein: WO 98/23635 (Frazer IH et al., assigned to The University of Queensland); Southwood S et al., 1998, J. Immunol. **160**: 3363-3373; Sinigaglia F et al., 1988, Nature **336**: 778-780; Chicz RM et al., 1993, J. Exp. Med **178**: 27-47; Hammer J et al., 1993, Cell **74**: 197-203; and Falk K et al., 1994, Immunogenetics **39**: 230-242. The latter reference also deals with HLA-DQ and -DP ligands. All epitopes listed in these 5 references are relevant as candidate natural epitopes to be used in the present invention, as are epitopes which share common motifs with these.

Alternatively, the epitope can be any artificial T-cell epitope which is capable of binding a large proportion of MHC Class II molecules. In this context the pan DR epitope peptides ("PADRE") described in WO 95/07707 and in the corresponding paper Alexander J et al., 1994, Immunity **1**: 751-761 (both disclosures are incorporated by reference herein) are interesting candidates for epitopes to be used according to the present invention. It should be noted that the most effective PADRE peptides disclosed in these papers carry D-amino acids in the C- and N-termini in order to improve stability when administered. However, the present invention primarily aims at incorporating the relevant epitopes as part of the modified IL5 which should then subsequently be broken down enzymatically inside the lysosomal compartment of APCs to allow subsequent presentation in the context of an MHC-II molecule and therefore it is not expedient to incorporate D-amino acids in the epitopes used in the present invention.

One especially preferred PADRE peptide is the one having the amino acid sequence AKFVAAWTLKAAA (SEQ ID NO: 65) or an immunologically effective subsequence thereof. This, and other epitopes having the same lack of MHC restriction are preferred

T-cell epitopes which should be present in the IL5 analogues used in the inventive method. Such super-promiscuous epitopes will allow for the most simple embodiments of the invention wherein only one single modified IL5 is presented to the vaccinated animal's immune system.

As mentioned above, the modification of IL5 can also include the introduction of a first moiety which targets the modified IL5 to an APC or a B-lymphocyte. For instance, the first moiety can be a specific binding partner for a B-lymphocyte specific surface antigen or for an APC specific surface antigen. Many such specific surface antigens are known in the art. For instance, the moiety can be a carbohydrate for which there is a receptor on the B-lymphocyte or on the APC (e.g. mannan or mannose). Alternatively, the second moiety can be a haptan. Also an antibody fragment which specifically recognizes a surface molecule on APCs or lymphocytes can be used as a first moiety (the surface molecule can e.g. be an FC $\gamma$  receptor of macrophages and monocytes, such as FC $\gamma$ RI or, alternatively any other specific surface marker such as CD40 or CTLA-4). It should be noted that all these exemplary targeting molecules can be used as part of an adjuvant also, cf. below.

As an alternative or supplement to targeting the modified IL5 polypeptide to a certain cell type in order to achieve an enhanced immune response, it is possible to increase the level of responsiveness of the immune system by including the above-mentioned second moiety which stimulates the immune system. Typical examples of such second moieties are cytokines, and heat-shock proteins or molecular chaperones, as well as effective parts thereof.

Suitable cytokines to be used according to the invention are those which will normally also function as adjuvants in a vaccine composition, i.e. for instance interferon  $\gamma$  (IFN- $\gamma$ ), Flt3L, interleukin 1 (IL-1), interleukin 2 (IL-2), interleukin 4 (IL-4), interleukin 6 (IL-6), interleukin 12 (IL-12), interleukin 13 (IL-13), interleukin 15 (IL-15), and granulocyte-

macrophage colony stimulating factor (GM-CSF); alternatively, the functional part of the cytokine molecule may suffice as the second moiety. With respect to the use of such cytokines as adjuvant substances, cf. the discussion below. It should be  
5 noted that use of both IL-4 and IL-13 should be exercised very carefully, if at all, as both molecules are known as key effector molecules in the pathophysiology of atopy and asthma.

According to the invention, suitable heat-shock proteins or  
10 molecular chaperones used as the second moiety can be HSP70, HSP90, HSC70, GRP94 (also known as gp96, cf. Wearsch PA et al. 1998, Biochemistry 37: 5709-19), and CRT (calreticulin).

Alternatively, the second moiety can be a toxin, such as listeriolysin (LLO), lipid A and heat-labile enterotoxin. Also,  
15 a number of mycobacterial derivatives such as MDP (muramyl dipeptide) and the trehalose diesters TDM and TDE are interesting possibilities.

Also the possibility of introducing a third moiety which enhances the presentation of the modified IL5 to the immune system is an important embodiment of the invention. The art has shown several examples of this principle. For instance, it is known that the palmitoyl lipidation anchor in the *Borrelia burgdorferi* protein OspA can be utilised so as to provide  
25 self-adjuvating polypeptides (cf. e.g. WO 96/40718). It seems that the lipidated proteins form up micelle-like structures with a core consisting of the lipidation anchor parts of the polypeptides and the remaining parts of the molecule protruding therefrom, resulting in multiple presentations of the antigenic determinants. Hence, the use of this and related approaches using different lipidation anchors (e.g. a myristyl group, a myristyl group, a farnesyl group, a geranyl-geranyl group, a GPI-anchor, and an N-acyl diglyceride group) are preferred  
30 embodiments of the invention, especially since the provision of such a lipidation anchor in a recombinantly produced protein is fairly straightforward and merely requires use of e.g. a naturally occurring signal sequence as a fusion partner  
35

for the modified IL5 polypeptide. Another possibility is use of the C3d fragment of complement factor C3 or C3 itself (cf. Dempsey et al., 1996, Science 271, 348-350 and Lou & Kohler, 1998, Nature Biotechnology 16, 458-462).

5

An alternative embodiment of the invention which also results in the preferred presentation of multiple (e.g. at least 2) copies of the important epitopic regions of IL5 to the immune system is the covalent or non-covalent coupling of IL5, subse-

10 quence or variants thereof to certain carrier molecules. For instance, polymers can be used, e.g. carbohydrates such as dextran, cf. e.g. Lees A et al., 1994, Vaccine 12: 1160-1166; Lees A et al., 1990, J Immunol. 145: 3594-3600, but also man-

15 nose and mannan are useful alternatives. Integral membrane proteins from e.g. *E. coli* and other bacteria are also useful conjugation partners. The traditional carrier molecules such as keyhole limpet hemocyanin (KLH), tetanus toxoid, diphtheria toxoid, and bovine serum albumin (BSA) are also preferred and useful conjugation partners.

20

Certain areas of native IL5 are believed to be superiorly suited for performing modifications. It is predicted that modifications in at least one of loops 1-3 or in the amino acid residues C-terminal to helix D (said loops and said helix

25 D corresponding to those shown in Fig. 3 for human and murine IL5) will be most likely to produce the desired constructs and vaccination results. Considerations underlying these chosen areas are a) preservation of known and predicted B-cell epitopes, b) preservation of tertiary and quaternary structures

30 etc, cf. also the discussion in the preamble to the examples. At any rate, as discussed above, it is fairly easy to screen a set of modified IL5 molecules which have all been subjected to introduction of a T-cell epitope in different locations.

35 Since the most preferred embodiments of the present invention involves down-regulation of human IL5, it is consequently preferred that the IL5 polypeptide discussed above is a human IL5 polypeptide. In this embodiment, it is especially preferred

that the human IL5 polypeptide has been modified by substituting at least one amino acid sequence in SEQ ID NO: 1 with at least one amino acid sequence of equal or different length and containing a foreign T<sub>H</sub> epitope, wherein substituted amino acid residues are selected from the group consisting of residues 87-90, residues 32-43, residues 59-64, residues 86-91, and residues 110-113. The rationale behind such constructs is discussed in detail in the examples.

#### 10 Formulation of IL5 and modified IL5 polypeptides

When effecting presentation of the IL5 polypeptide or the modified IL5 polypeptide to an animal's immune system by means of administration thereof to the animal, the formulation of the polypeptide follows the principles generally acknowledged in the art.

Preparation of vaccines which contain peptide sequences as active ingredients is generally well understood in the art, as exemplified by U.S. Patents 4,608,251; 4,601,903; 4,599,231; 4,599,230; 4,596,792; and 4,578,770, all incorporated herein by reference. Typically, such vaccines are prepared as injectables either as liquid solutions or suspensions; solid forms suitable for solution in, or suspension in, liquid prior to injection may also be prepared. The preparation may also be emulsified. The active immunogenic ingredient is often mixed with excipients which are pharmaceutically acceptable and compatible with the active ingredient. Suitable excipients are, for example, water, saline, dextrose, glycerol, ethanol, or the like, and combinations thereof. In addition, if desired, the vaccine may contain minor amounts of auxiliary substances such as wetting or emulsifying agents, pH buffering agents, or adjuvants which enhance the effectiveness of the vaccines; cf. the detailed discussion of adjuvants below.

The vaccines are conventionally administered parenterally, by injection, for example, either subcutaneously, intracutaneously, intradermally, subdermally or intramuscularly. Addi-

5 tional formulations which are suitable for other modes of administration include suppositories and, in some cases, oral, buccal, sublingual, intraperitoneal, intravaginal, anal, epidural, spinal, and intracranial formulations. For suppositories, traditional binders and carriers may include, for example, polyalkylene glycols or triglycerides; such suppositories may be formed from mixtures containing the active ingredient in the range of 0.5% to 10%, preferably 1-2%. Oral formulations include such normally employed excipients as, for example, pharmaceutical grades of mannitol, lactose, starch, magnesium stearate, sodium saccharine, cellulose, magnesium carbonate, and the like. These compositions take the form of solutions, suspensions, tablets, pills, capsules, sustained release formulations or powders and contain 10-95% of active ingredient, preferably 25-70%. For oral formulations, cholera toxin is an interesting formulation partner (and also a possible conjugation partner).

20 The polypeptides may be formulated into the vaccine as neutral or salt forms. Pharmaceutically acceptable salts include acid addition salts (formed with the free amino groups of the peptide) and which are formed with inorganic acids such as, for example, hydrochloric or phosphoric acids, or such organic acids as acetic, oxalic, tartaric, mandelic, and the like. Salts formed with the free carboxyl groups may also be derived from inorganic bases such as, for example, sodium, potassium, ammonium, calcium, or ferric hydroxides, and such organic bases as isopropylamine, trimethylamine, 2-ethylamino ethanol, histidine, procaine, and the like.

30

The vaccines are administered in a manner compatible with the dosage formulation, and in such amount as will be therapeutically effective and immunogenic. The quantity to be administered depends on the subject to be treated, including, e.g., the capacity of the individual's immune system to mount an immune response, and the degree of protection desired. Suitable dosage ranges are of the order of several hundred micrograms active ingredient per vaccination with a preferred range from



about 0.1  $\mu\text{g}$  to 2,000  $\mu\text{g}$  (even though higher amounts in the 1-10 mg range are contemplated), such as in the range from about 0.5  $\mu\text{g}$  to 1,000  $\mu\text{g}$ , preferably in the range from 1  $\mu\text{g}$  to 500  $\mu\text{g}$  and especially in the range from about 10  $\mu\text{g}$  to 100  $\mu\text{g}$ .

5 Suitable regimens for initial administration and booster shots are also variable but are typified by an initial administration followed by subsequent inoculations or other administrations.

10 The manner of application may be varied widely. Any of the conventional methods for administration of a vaccine are applicable. These include oral application on a solid physiologically acceptable base or in a physiologically acceptable dispersion, parenterally, by injection or the like. The dosage  
15 of the vaccine will depend on the route of administration and will vary according to the age of the person to be vaccinated and the formulation of the antigen.

Some of the polypeptides of the vaccine are sufficiently immunogenic in a vaccine, but for some of the others the immune  
20 response will be enhanced if the vaccine further comprises an adjuvant substance.

Various methods of achieving adjuvant effect for the vaccine  
25 are known. General principles and methods are detailed in "The Theory and Practical Application of Adjuvants", 1995, Duncan E.S. Stewart-Tull (ed.), John Wiley & Sons Ltd, ISBN 0-471-95170-6, and also in "Vaccines: New Generation Immunological Adjuvants", 1995, Gregoriadis G et al. (eds.), Plenum Press,  
30 New York, ISBN 0-306-45283-9, both of which are hereby incorporated by reference herein.

It is especially preferred to use an adjuvant which can be demonstrated to facilitate breaking of the autotolerance to  
35 autoantigens; in fact, this is essential in cases where unmodified IL5 is used as the active ingredient in the autovaccine. Non-limiting examples of suitable adjuvants are selected from the group consisting of an immune targeting adjuvant; an

immune modulating adjuvant such as a toxin, a cytokine, and a mycobacterial derivative; an oil formulation; a polymer; a micelle forming adjuvant; a saponin; an immunostimulating complex matrix (ISCOM matrix); a particle; DDA; aluminium adjuvants; DNA adjuvants;  $\gamma$ -inulin; and an encapsulating adjuvant. In general it should be noted that the disclosures above which relate to compounds and agents useful as first, second and third moieties in the analogues also refer *mutatis mutandis* to their use in the adjuvant of a vaccine of the invention.

The application of adjuvants include use of agents such as aluminium hydroxide or phosphate (alum), commonly used as 0.05 to 0.1 percent solution in buffered saline, admixture with synthetic polymers of sugars (e.g. Carbopol®) used as 0.25 percent solution, aggregation of the protein in the vaccine by heat treatment with temperatures ranging between 70° to 101°C for 30 second to 2 minute periods respectively and also aggregation by means of cross-linking agents are possible. Aggregation by reactivation with pepsin treated antibodies (Fab fragments) to albumin, mixture with bacterial cells such as *C. parvum* or endotoxins or lipopolysaccharide components of gram-negative bacteria, emulsion in physiologically acceptable oil vehicles such as mannide mono-oleate (Aracel A) or emulsion with 20 percent solution of a perfluorocarbon (Fluosol-DA) used as a block substitute may also be employed. Admixture with oils such as squalene and IFA is also preferred.

According to the invention DDA (dimethyldioctadecylammonium bromide) is an interesting candidate for an adjuvant as is DNA and  $\gamma$ -inulin, but also Freund's complete and incomplete adjuvants as well as *quillaja* saponins such as QuilA and QS21 are interesting as is RIBI. Further possibilities are monophosphoryl lipid A (MPL), the above mentioned C3 and C3d, and muramyl dipeptide (MDP).

Liposome formulations are also known to confer adjuvant effects, and therefore liposome adjuvants are preferred according to the invention.

Also immunostimulating complex matrix type (ISCOM® matrix) adjuvants are preferred choices according to the invention, especially since it has been shown that this type of adjuvants

5 are capable of up-regulating MHC Class II expression by APCs.

An ISCOM® matrix consists of (optionally fractionated) saponins (triterpenoids) from *Quillaja saponaria*, cholesterol, and phospholipid. When admixed with the immunogenic protein, the

10 resulting particulate formulation is what is known as an ISCOM particle where the saponin constitutes 60-70% w/w, the cholesterol and phospholipid 10-15% w/w, and the protein 10-15% w/w.

Details relating to composition and use of immunostimulating complexes can e.g. be found in the above-mentioned text-books dealing with adjuvants, but also Morein B et al., 1995, Clin.

15 Immunother. 3: 461-475 as well as Barr IG and Mitchell GF, 1996, Immunol. and Cell Biol. 74: 8-25 (both incorporated by reference herein) provide useful instructions for the preparation of complete immunostimulating complexes.

20 Another highly interesting (and thus, preferred) possibility of achieving adjuvant effect is to employ the technique described in Gosselin et al., 1992 (which is hereby incorporated by reference herein). In brief, the presentation of a relevant antigen such as an antigen of the present invention can be enhanced by conjugating the antigen to antibodies (or antigen binding antibody fragments) against the Fcγ receptors on monocytes/macrophages. Especially conjugates between antigen and anti-FcγRI have been demonstrated to enhance immunogenicity for the purposes of vaccination.

30

Other possibilities involve the use of the targeting and immune modulating substances (i.a. cytokines) mentioned above as candidates for the first and second moieties in the modified versions of IL5. In this connection, also synthetic inducers

35 of cytokines like poly I:C are possibilities.

Suitable mycobacterial derivatives are selected from the group consisting of muramyl dipeptide, complete Freund's adjuvant, RIBI, and a diester of trehalose such as TDM and TDE.

- 5 Suitable immune targeting adjuvants are selected from the group consisting of CD40 ligand and CD40 antibodies or specifically binding fragments thereof (cf. the discussion above), mannose, a Fab fragment, and CTLA-4.
- 10 Suitable polymer adjuvants are selected from the group consisting of a carbohydrate such as dextran, PEG, starch, mannan, and mannose; a plastic polymer such as; and latex such as latex beads.
- 15 Yet another interesting way of modulating an immune response is to include the IL5 immunogen (optionally together with adjuvants and pharmaceutically acceptable carriers and vehicles) in a "virtual lymph node" (VLN) (a proprietary medical device developed by ImmunoTherapy, Inc., 360 Lexington Avenue, New  
20 York, NY 10017-6501). The VLN (a thin tubular device) mimics the structure and function of a lymph node. Insertion of a VLN under the skin creates a site of sterile inflammation with an upsurge of cytokines and chemokines. T- and B-cells as well as APCs rapidly respond to the danger signals, home to the in-  
25 flamed site and accumulate inside the porous matrix of the VLN. It has been shown that the necessary antigen dose required to mount an immune response to an antigen is reduced when using the VLN and that immune protection conferred by vaccination using a VLN surpassed conventional immunization  
30 using Ribi as an adjuvant. The technology is *i.a.* described briefly in Gelber C et al., 1998, "Elicitation of Robust Cellular and Humoral Immune Responses to Small Amounts of Immunogens Using a Novel Medical Device Designated the Virtual Lymph Node", in: "From the Laboratory to the Clinic, Book of Abstracts, October 12<sup>th</sup> - 15<sup>th</sup> 1998, Seascape Resort, Aptos, California".

It is expected that the vaccine should be administered at least once a year, such as at least 1, 2, 3, 4, 5, 6, and 12 times a year. More specifically, 1-12 times per year is expected, such as 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, or 12 times a year to an individual in need thereof. It has previously been shown that the memory immunity induced by the use of the preferred autovaccines according to the invention is not permanent, and therefor the immune system needs to be periodically challenged with the analogues.

Due to genetic variation, different individuals may react with immune responses of varying strength to the same polypeptide. Therefore, the vaccine according to the invention may comprise several different polypeptides in order to increase the immune response, cf. also the discussion above concerning the choice of foreign T-cell epitope introductions. The vaccine may comprise two or more polypeptides, where all of the polypeptides are as defined above.

The vaccine may consequently comprise 3-20 different modified or unmodified polypeptides, such as 3-10 different polypeptides. However, normally the number of polypeptides will be sought kept to a minimum such as 1 or 2 polypeptides.

## 25 Nucleic acid vaccination

As an alternative to classic administration of a peptide-based vaccine, the technology of nucleic acid vaccination (also known as "nucleic acid immunisation", "genetic immunisation", and "gene immunisation") offers a number of attractive features.

First, in contrast to the traditional vaccine approach, nucleic acid vaccination does not require resource consuming large-scale production of the immunogenic agent (e.g. in the form of industrial scale fermentation of microorganisms producing modified IL5). Furthermore, there is no need to devise purification and refolding schemes for the immunogen. And fi-

nally, since nucleic acid vaccination relies on the biochemical apparatus of the vaccinated individual in order to produce the expression product of the nucleic acid introduced, the optimum posttranslational processing of the expression  
5 product is expected to occur; this is especially important in the case of autovaccination, since, as mentioned above, a significant fraction of the original IL5 B-cell epitopes should be preserved in the modified molecule, and since B-cell epitopes in principle can be constituted by parts of any  
10 (bio)molecule (e.g. carbohydrate, lipid, protein etc.). Therefore, native glycosylation and lipidation patterns of the immunogen may very well be of importance for the overall immunogenicity and this is expected to be ensured by having the host producing the immunogen.

Hence, a preferred embodiment of the invention comprises effecting presentation of modified IL5 to the immune system by introducing nucleic acid(s) encoding the modified IL5 into the animal's cells and thereby obtaining *in vivo* expression by the  
20 cells of the nucleic acid(s) introduced.

In this embodiment, the introduced nucleic acid is preferably DNA which can be in the form of naked DNA, DNA formulated with charged or uncharged lipids, DNA formulated in liposomes, DNA  
25 included in a viral vector, DNA formulated with a transfection-facilitating protein or polypeptide, DNA formulated with a targeting protein or polypeptide, DNA formulated with Calcium precipitating agents, DNA coupled to an inert carrier molecule, DNA encapsulated in a polymer, e.g. in PLGA (cf. the  
30 microencapsulation technology described in WO 98/31398) or in chitin or chitosan, and DNA formulated with an adjuvant. In this context it is noted that practically all considerations pertaining to the use of adjuvants in traditional vaccine formulation apply for the formulation of DNA vaccines. Hence, all  
35 disclosures herein which relate to use of adjuvants in the context of polypeptide based vaccines apply *mutatis mutandis* to their use in nucleic acid vaccination technology.

As for routes of administration and administration schemes of polypeptide based vaccines which have been detailed above, these are also applicable for the nucleic acid vaccines of the invention and all discussions above pertaining to routes of administration and administration schemes for polypeptides apply *mutatis mutandis* to nucleic acids. To this should be added that nucleic acid vaccines can suitably be administered intravenously and intraarterially. Furthermore, it is well-known in the art that nucleic acid vaccines can be administered by use of a so-called gene gun, and hence also this and equivalent modes of administration are regarded as part of the present invention. Finally, also the use of a VLN in the administration of nucleic acids has been reported to yield good results, and therefore this particular mode of administration is particularly preferred.

Furthermore, the nucleic acid(s) used as an immunization agent can contain regions encoding the 1<sup>st</sup>, 2<sup>nd</sup> and/or 3<sup>rd</sup> moieties, e.g. in the form of the immunomodulating substances described above such as the cytokines discussed as useful adjuvants. A preferred version of this embodiment encompasses having the coding region for the analogue and the coding region for the immunomodulator in different reading frames or at least under the control of different promoters. Thereby it is avoided that the analogue or epitope is produced as a fusion partner to the immunomodulator. Alternatively, two distinct nucleotide fragments can be used, but this is less preferred because of the advantage of ensured co-expression when having both coding regions included in the same molecule.

Accordingly, the invention also relates to a composition for inducing production of antibodies against IL5, the composition comprising

- a nucleic acid fragment or a vector of the invention (cf. the discussion of vectors below), and
- a pharmaceutically and immunologically acceptable vehicle and/or carrier and/or adjuvant as discussed above.

Under normal circumstances, the IL5 variant-encoding nucleic acid is introduced in the form of a vector wherein expression is under control of a viral promoter. For more detailed discussions of vectors and DNA fragments according to the invention, cf. the discussion below. Also, detailed disclosures relating to the formulation and use of nucleic acid vaccines are available, cf. Donnelly JJ et al, 1997, Annu. Rev. Immunol. 15: 617-648 and Donnelly JJ et al., 1997, Life Sciences 60: 163-172. Both of these references are incorporated by reference herein.

### Live vaccines

A third alternative for effecting presentation of modified IL5 to the immune system is the use of live vaccine technology. In live vaccination, presentation to the immune system is effected by administering, to the animal, a non-pathogenic microorganism which has been transformed with a nucleic acid fragment encoding a modified IL5 or with a vector incorporating such a nucleic acid fragment. The non-pathogenic microorganism can be any suitable attenuated bacterial strain (attenuated by means of passaging or by means of removal of pathogenic expression products by recombinant DNA technology), e.g. *Mycobacterium bovis* BCG., non-pathogenic *Streptococcus* spp., *E. coli*, *Salmonella* spp., *Vibrio cholerae*, *Shigella*, etc. Reviews dealing with preparation of state-of-the-art live vaccines can e.g. be found in Saliou P, 1995, Rev. Prat. 45: 1492-1496 and Walker PD, 1992, Vaccine 10: 977-990, both incorporated by reference herein. For details about the nucleic acid fragments and vectors used in such live vaccines, cf. the discussion below.

As an alternative to bacterial live vaccines, the nucleic acid fragment of the invention discussed below can be incorporated in a non-virulent viral vaccine vector such as a vaccinia strain or any other suitable pox virus.



Normally, the non-pathogenic microorganism or virus is administered only once to the animal, but in certain cases it may be necessary to administer the microorganism more than once in a lifetime in order to maintain protective immunity. It is even contemplated that immunization schemes as those detailed above for polypeptide vaccination will be useful when using live or virus vaccines.

Alternatively, live or virus vaccination is combined with previous or subsequent polypeptide and/or nucleic acid vaccination. For instance, it is possible to effect primary immunization with a live or virus vaccine followed by subsequent booster immunizations using the polypeptide or nucleic acid approach.

The microorganism or virus can be transformed with nucleic acid(s) containing regions encoding the 1<sup>st</sup>, 2<sup>nd</sup> and/or 3<sup>rd</sup> moieties, e.g. in the form of the immunomodulating substances described above such as the cytokines discussed as useful adjuvants. A preferred version of this embodiment encompasses having the coding region for the analogue and the coding region for the immunomodulator in different reading frames or at least under the control of different promoters. Thereby it is avoided that the analogue or epitopes are produced as fusion partners to the immunomodulator. Alternatively, two distinct nucleotide fragments can be used as transforming agents. Of course, having the 1<sup>st</sup> and/or 2<sup>nd</sup> and/or 3<sup>rd</sup> moieties in the same reading frame can provide as an expression product, an analogue of the invention, and such an embodiment is especially preferred according to the present invention.

#### Use of the method of the invention in disease treatment

As will be appreciated from the discussions above, the provision of the method of the invention allows for control of diseases characterized by eosinophilia. In this context, asthma is the key target for the inventive method but also other chronic allergic conditions such as multiple allergy and al-

lergic rhinitis are feasible targets for treatment/amelioration. Hence, an important embodiment of the method of the invention for down-regulating IL5 activity comprises treating and/or preventing and/or ameliorating asthma or other chronic allergic conditions characterized by eosinophilia, the method comprising down-regulating IL5 activity according to the method of the invention to such an extent that the number of eosinophil cells is significantly reduced.

- 10 In the present context such a significant reduction in eosinophil cell numbers is at least 20% compared to the eosinophil number prior to treatment, but higher percentages are contemplated, such as at least 30%, at least 40%, at least 50%, at least 60%, at least 70%, at least 80% and even at least 90%.  
15 The reduction may be systemic or, more often, locally in e.g. the lungs.

Eosinophil cell numbers are determined by methods known in the art, typically using microscopy of a suitable sample (such as a BAL fluid) and counting the number of eosinophil cells manually under microscope. Alternatively, eosinophil numbers can be counted using flow cytometric methods or any other convenient method of cytometry capable of distinguishing eosinophils.

25

#### Peptides, polypeptides, and compositions of the invention

- As will be apparent from the above, the present invention is based on the concept of immunising individuals against the IL5 antigen in order to indirectly obtain a reduction in eosinophil cell numbers. The preferred way of obtaining such an immunization is to use modified versions of IL5, thereby providing molecules which have not previously been disclosed in the art.

35

It is believed that the modified IL5 molecules discussed herein are inventive in their own right, and therefore an important part of the invention pertains to an IL5 analogue

which is derived from an animal IL5 wherein is introduced a modification which has as a result that immunization of the animal with the analogue induces production of antibodies cross-reacting with the unmodified IL5 polypeptide. Preferably, the nature of the modification conforms with the types of modifications described above when discussing various embodiments of the method of the invention when using modified IL5. Hence, any disclosure presented herein pertaining to modified IL5 molecules are relevant for the purpose of describing the IL5 analogues of the invention, and any such disclosures apply *mutatis mutandis* to the description of these analogues.

It should be noted that preferred modified IL5 molecules comprise modifications which results in a polypeptide having a sequence identity of at least 70% with IL5 or with a subsequence thereof of at least 10 amino acids in length. Higher sequence identities are preferred, e.g. at least 75% or even at least 80% or 85%. The sequence identity for proteins and nucleic acids can be calculated as  $(N_{ref} - N_{dif}) \cdot 100 / N_{ref}$ , wherein  $N_{dif}$  is the total number of non-identical residues in the two sequences when aligned and wherein  $N_{ref}$  is the number of residues in one of the sequences. Hence, the DNA sequence AGTCAGTC will have a sequence identity of 75% with the sequence AATCAATC ( $N_{dif}=2$  and  $N_{ref}=8$ ).

The invention also pertains to compositions useful in exercising the method of the invention. Hence, the invention also relates to an immunogenic composition comprising an immunogenically effective amount of an IL5 polypeptide which is a self-protein in an animal, said IL5 polypeptide being formulated together with an immunologically acceptable adjuvant so as to break the animal's autotolerance towards the IL5 polypeptide, the composition further comprising a pharmaceutically and immunologically acceptable vehicle and/or carrier. In other words, this part of the invention pertains to the formulations of naturally occurring IL5 polypeptides which have

been described in connection with embodiments of the method of the invention.

The invention also relates to an immunogenic composition comprising an immunologically effective amount of an IL5 analogue defined above, said composition further comprising a pharmaceutically and immunologically acceptable diluent and/or vehicle and/or carrier and/or excipient and optionally an adjuvant. In other words, this part of the invention concerns formulations of modified IL5, essentially as described hereinabove. The choice of adjuvants, carriers, and vehicles is accordingly in line with what has been discussed above when referring to formulation of modified and unmodified IL5 for use in the inventive method for the down-regulation of IL5.

The polypeptides are prepared according to methods well-known in the art. Longer polypeptides are normally prepared by means of recombinant gene technology including introduction of a nucleic acid sequence encoding the IL5 analogue into a suitable vector, transformation of a suitable host cell with the vector, expression of the nucleic acid sequence, recovery of the expression product from the host cells or their culture supernatant, and subsequent purification and optional further modification, e.g. refolding or derivatization.

Shorter peptides are preferably prepared by means of the well-known techniques of solid- or liquid-phase peptide synthesis. However, recent advances in this technology has rendered possible the production of full-length polypeptides and proteins by these means, and therefore it is also within the scope of the present invention to prepare the long constructs by synthetic means.

#### Nucleic acid fragments and vectors of the invention

It will be appreciated from the above disclosure that modified IL5 polypeptides can be prepared by means of recombinant gene technology but also by means of chemical synthesis or semisyn-

thesis; the latter two options are especially relevant when the modification consists in coupling to protein carriers (such as KLH, diphtheria toxoid, tetanus toxoid, and BSA) and non-proteinaceous molecules such as carbohydrate polymers and  
5 of course also when the modification comprises addition of side chains or side groups to an IL5 polypeptide-derived peptide chain.

For the purpose of recombinant gene technology, and of course  
10 also for the purpose of nucleic acid immunization, nucleic acid fragments encoding modified IL5 are important chemical products. Hence, an important part of the invention pertains to a nucleic acid fragment which encodes an IL5 analogue, i.e. an IL5 derived polypeptide which either comprises the natural  
15 IL5 sequence to which has been added or inserted a fusion partner or, preferably an IL5 derived polypeptide wherein has been introduced a foreign T-cell epitope by means of insertion and/or addition, preferably by means of substitution and/or deletion. The nucleic acid fragments of the invention are ei-  
20 ther DNA or RNA fragments.

The nucleic acid fragments of the invention will normally be inserted in suitable vectors to form cloning or expression vectors carrying the nucleic acid fragments of the invention;  
25 such novel vectors are also part of the invention. Details concerning the construction of these vectors of the invention will be discussed in context of transformed cells and microorganisms below. The vectors can, depending on purpose and type of application, be in the form of plasmids, phages, cosmids,  
30 mini-chromosomes, or virus, but also naked DNA which is only expressed transiently in certain cells is an important vector. Preferred cloning and expression vectors of the invention are capable of autonomous replication, thereby enabling high copy-numbers for the purposes of high-level expression or high-  
35 level replication for subsequent cloning.

The general outline of a vector of the invention comprises the following features in the 5'→3' direction and in operable

linkage: a promoter for driving expression of the nucleic acid fragment of the invention, optionally a nucleic acid sequence encoding a leader peptide enabling secretion (to the extracellular phase or, where applicable, into the periplasma) of or  
5 integration into the membrane of the polypeptide fragment, the nucleic acid fragment of the invention, and optionally a nucleic acid sequence encoding a terminator. When operating with expression vectors in producer strains or cell-lines it is for the purposes of genetic stability of the transformed cell preferred that the vector when introduced into a host cell is integrated in the host cell genome. In contrast, when working with vectors to be used for effecting *in vivo* expression in an animal (*i.e.* when using the vector in DNA vaccination) it is for security reasons preferred that the vector is not incapable of being integrated in the host cell genome; typically,  
15 naked DNA or non-integrating viral vectors are used, the choices of which are well-known to the person skilled in the art.

20 The vectors of the invention are used to transform host cells to produce the modified IL5 polypeptide of the invention. Such transformed cells, which are also part of the invention, can be cultured cells or cell lines used for propagation of the nucleic acid fragments and vectors of the invention, or used  
25 for recombinant production of the modified IL5 polypeptides of the invention. Alternatively, the transformed cells can be suitable live vaccine strains wherein the nucleic acid fragment (one single or multiple copies) have been inserted so as to effect secretion or integration into the bacterial membrane  
30 or cell-wall of the modified IL5.

Preferred transformed cells of the invention are microorganisms such as bacteria (such as the species *Escherichia* [*e.g.* *E. coli*], *Bacillus* [*e.g.* *Bacillus subtilis*], *Salmonella*, or  
35 *Mycobacterium* [preferably non-pathogenic, *e.g.* *M. bovis* BCG]), yeasts (such as *Saccharomyces cerevisiae*), and protozoans. Alternatively, the transformed cells are derived from a multicellular organism such as a fungus, an insect cell, a plant

cell, or a mammalian cell. Most preferred are cells derived from a human being, cf. the discussion of cell lines and vectors below. Recent results have shown great promise in the use of a commercially available *Drosophila melanogaster* cell line (the Schneider 2 (S<sub>2</sub>) cell line and vector system available from Invitrogen) for the recombinant production of IL5 analogues of the invention, and therefore this expression system is particularly preferred.

10 For the purposes of cloning and/or optimized expression it is preferred that the transformed cell is capable of replicating the nucleic acid fragment of the invention. Cells expressing the nucleic acid fragment are preferred useful embodiments of the invention; they can be used for small-scale or large-scale  
15 preparation of the modified IL5 or, in the case of non-pathogenic bacteria, as vaccine constituents in a live vaccine.

When producing the modified IL5 of the invention by means of transformed cells, it is convenient, although far from essential, that the expression product is either exported out into  
20 the culture medium or carried on the surface of the transformed cell.

When an effective producer cell has been identified it is preferred, on the basis thereof, to establish a stable cell line which carries the vector of the invention and which expresses the nucleic acid fragment encoding the modified IL5. Preferably, this stable cell line secretes or carries the IL5 analogue of the invention, thereby facilitating purification  
25 thereof.  
30 thereof.

In general, plasmid vectors containing replicon and control sequences which are derived from species compatible with the host cell are used in connection with the hosts. The vector  
35 ordinarily carries a replication site, as well as marking sequences which are capable of providing phenotypic selection in transformed cells. For example, *E. coli* is typically transformed using pBR322, a plasmid derived from an *E. coli* species

(see, e.g., Bolivar et al., 1977). The pBR322 plasmid contains genes for ampicillin and tetracycline resistance and thus provides easy means for identifying transformed cells. The pBR plasmid, or other microbial plasmid or phage must also contain, or be modified to contain, promoters which can be used by the prokaryotic microorganism for expression.

Those promoters most commonly used in prokaryotic recombinant DNA construction include the B-lactamase (penicillinase) and lactose promoter systems (Chang et al., 1978; Itakura et al., 1977; Goeddel et al., 1979) and a tryptophan (trp) promoter system (Goeddel et al., 1979; EP-A-0 036 776). While these are the most commonly used, other microbial promoters have been discovered and utilized, and details concerning their nucleotide sequences have been published, enabling a skilled worker to ligate them functionally with plasmid vectors (Siebwenlist et al., 1980). Certain genes from prokaryotes may be expressed efficiently in *E. coli* from their own promoter sequences, precluding the need for addition of another promoter by artificial means.

In addition to prokaryotes, eukaryotic microbes, such as yeast cultures may also be used, and here the promoter should be capable of driving expression. *Saccharomyces cerevisiae*, or common baker's yeast is the most commonly used among eukaryotic microorganisms, although a number of other strains are commonly available. For expression in *Saccharomyces*, the plasmid YRp7, for example, is commonly used (Stinchcomb et al., 1979; Kingsman et al., 1979; Tschemper et al., 1980). This plasmid already contains the trp1 gene which provides a selection marker for a mutant strain of yeast lacking the ability to grow in tryptophan for example ATCC No. 44076 or PEP4-1 (Jones, 1977). The presence of the trp1 lesion as a characteristic of the yeast host cell genome then provides an effective environment for detecting transformation by growth in the absence of tryptophan.



Suitable promoting sequences in yeast vectors include the promoters for 3-phosphoglycerate kinase (Hitzman et al., 1980) or other glycolytic enzymes (Hess et al., 1968; Holland et al., 1978), such as enolase, glyceraldehyde-3-phosphate dehydrogenase, hexokinase, pyruvate decarboxylase, phosphofructokinase, glucose-6-phosphate isomerase, 3-phosphoglycerate mutase, pyruvate kinase, triosephosphate isomerase, phosphoglucose isomerase, and glucokinase. In constructing suitable expression plasmids, the termination sequences associated with these genes are also ligated into the expression vector 3' of the sequence desired to be expressed to provide polyadenylation of the mRNA and termination.

Other promoters, which have the additional advantage of transcription controlled by growth conditions are the promoter region for alcohol dehydrogenase 2, isocytochrome C, acid phosphatase, degradative enzymes associated with nitrogen metabolism, and the aforementioned glyceraldehyde-3-phosphate dehydrogenase, and enzymes responsible for maltose and galactose utilization. Any plasmid vector containing a yeast-compatible promoter, origin of replication and termination sequences is suitable.

In addition to microorganisms, cultures of cells derived from multicellular organisms may also be used as hosts. In principle, any such cell culture is workable, whether from vertebrate or invertebrate culture. However, interest has been greatest in vertebrate cells, and propagation of vertebrate in culture (tissue culture) has become a routine procedure in recent years (Tissue Culture, 1973). Examples of such useful host cell lines are VERO and HeLa cells, Chinese hamster ovary (CHO) cell lines, and W138, BHK, COS-7 293, *Spodoptera frugiperda* (SF) cells (commercially available as complete expression systems from i.a. Protein Sciences, 1000 Research Parkway, Meriden, CT 06450, U.S.A. and from Invitrogen), and MDCK cell lines. In the present invention, an especially preferred cell line is S<sub>2</sub> available from Invitrogen, PO Box 2312, 9704 CH Groningen, The Netherlands.

Expression vectors for such cells ordinarily include (if necessary) an origin of replication, a promoter located in front of the gene to be expressed, along with any necessary ribosome binding sites, RNA splice sites, polyadenylation site, and transcriptional terminator sequences.

For use in mammalian cells, the control functions on the expression vectors are often provided by viral material. For example, commonly used promoters are derived from polyoma, Adenovirus 2, and most frequently Simian Virus 40 (SV40). The early and late promoters of SV40 virus are particularly useful because both are obtained easily from the virus as a fragment which also contains the SV40 viral origin of replication (Fiers et al., 1978). Smaller or larger SV40 fragments may also be used, provided there is included the approximately 250 bp sequence extending from the *HindIII* site toward the *BglI* site located in the viral origin of replication. Further, it is also possible, and often desirable, to utilize promoter or control sequences normally associated with the desired gene sequence, provided such control sequences are compatible with the host cell systems.

An origin of replication may be provided either by construction of the vector to include an exogenous origin, such as may be derived from SV40 or other viral (e.g., Polyoma, Adeno, VSV, BPV) or may be provided by the host cell chromosomal replication mechanism. If the vector is integrated into the host cell chromosome, the latter is often sufficient.

#### Identification of useful IL5 analogues

It will be clear to the skilled person that not all variants or modifications of native IL5 will have the ability to elicit antibodies in an animal which are cross-reactive with the native form. It is, however, not difficult to set up an effective standard screen for modified IL5 molecules which fulfill the minimum requirements for immunological reactivity dis-

cussed herein. Hence, another part of the invention concerns a method for the identification of a modified IL5 polypeptide which is capable of inducing antibodies against unmodified IL5 in an animal species where the unmodified IL5 polypeptide is a self-protein, the method comprising

- preparing, by means of peptide synthesis or by molecular biological means, a set of mutually distinct modified IL5 polypeptides wherein amino acids have been added to, inserted in, deleted from, or substituted into the amino acid sequence of an IL5 polypeptide of the animal species thereby giving rise to amino acid sequences in the set which comprise T-cell epitopes which are and foreign to the animal species, or preparing a set of nucleic acid fragments encoding the set of mutually distinct modified IL5 polypeptides,
- testing members of the set for their ability to induce production of antibodies by the animal species against the unmodified IL5, and
- identifying and optionally isolating the member(s) of the set which significantly induces antibody production against unmodified IL5 in the animal species, or identifying and optionally isolating the polypeptide expression products encoded by members of the set of nucleic acid fragments which significantly induces antibody production against unmodified IL5 polypeptide in the animal species.

In this context, the "set of mutually distinct modified IL5 polypeptides" is a collection of non-identical modified IL5 polypeptides which have e.g. been selected on the basis of the criteria discussed above (e.g. in combination with studies of circular dichroism, NMR spectra, and/or X-ray diffraction patterns). The set may consist of only a few members but it is contemplated that the set may contain several hundred members. Likewise, the set of nucleic acid fragments is a collection of non-identical nucleic acid fragments, each encoding a modified IL5 polypeptide selected in the same manner.

The test of members of the set can ultimately be performed *in vivo*, but a number of *in vitro* tests can be applied which narrow down the number of modified molecules which will serve the purpose of the invention.

5

Since the goal of introducing the foreign T-cell epitopes is to support the B-cell response by T-cell help, a prerequisite is that T-cell proliferation is induced by the modified IL5. T-cell proliferation can be tested by standardized proliferation assays *in vitro*. In short, a sample enriched for T-cells is obtained from a subject and subsequently kept in culture. The cultured T-cells are contacted with APCs of the subject which have previously taken up the modified molecule and processed it to present its T-cell epitopes. The proliferation of T-cells is monitored and compared to a suitable control (e.g. T-cells in culture contacted with APCs which have processed intact, native IL5). Alternatively, proliferation can be measured by determining the concentration of relevant cytokines released by the T-cells in response to their recognition of foreign T-cells.

Having rendered highly probable that at least one modified IL5 of the set is capable of inducing antibody production against IL5, it is possible to prepare an immunogenic composition comprising at least one modified IL5 polypeptide which is capable of inducing antibodies against unmodified IL5 in an animal species where the unmodified IL5 polypeptide is a self-protein, the method comprising admixing the member(s) of the set which significantly induces production of antibodies in the animal species which are reactive with IL5 with a pharmaceutically and immunologically acceptable carrier and/or vehicle and/or diluent and/or excipient, optionally in combination with at least one pharmaceutically and immunologically acceptable adjuvant.

35

Likewise, it is also possible to prepare an immunogenic composition which as an immunogen contains a nucleic acid fragment

encoding a immunogenic IL5 analogue, cf. the discussion of nucleic acid vaccination above.

The above aspects of the invention are conveniently carried out by initially preparing a number of mutually distinct nucleic acid sequences or vectors of the invention, inserting these into appropriate expression vectors, transforming suitable host cells with the vectors, and expressing the nucleic acid sequences of the invention. These steps can be followed by isolation of the expression products. It is preferred that the nucleic acid sequences and/or vectors are prepared by methods comprising exercise of a molecular amplification technique such as PCR or by means of nucleic acid synthesis.

#### PREAMBLE TO EXAMPLES

##### Vaccine design

The exemplary candidates for an IL5 autovaccine are constructed according to the AutoVac™ concept (described in detail in WO 95/05849) by substitution with known promiscuous T cell epitopes into the human IL5 wild type protein. The substitutions are peptide substitutions, where the inserted peptide may be of the same or different length than the deleted peptide in the wild-type sequence.

For initial proof of concept by *in vivo* testing and screening, it was decided to prepare the constructs in the murine IL5 sequence. By way of example, the tetanus toxoid epitopes P2 (SEQ ID NO: 23) and P30 (SEQ ID NO: 24) are used as substituting peptides, but any other suitable peptide containing or constituting a promiscuous T<sub>H</sub> epitope could, according to the present invention, be used.

It should be emphasized that the size of the molecule (115 res.) compared to the size of the substitutions (15 or 21 residues for P2 and P30, respectively) strongly limits the

possible sites of structural non-destructive inserts. As the disulfide bridges are important, but not imperative, for the dimerization, some variants are made in pairs +/- elimination of the cysteines.

5

In the construction of the candidate molecules, two basic parameters have been considered. First, it is attempted to conserve a maximum fraction of the three-dimensional structure of the wild type hIL5, thereby conserving the native B-cell epitope repertoire. This is supported by Dickason et al., (1994) who demonstrated that IL5 B-cell epitopes known to be neutralising are conformational. Conservation of the tertiary structure is sought achieved by introducing the modifications at structurally "neutral" sites, such as loops or separate segments. The fact that the N-terminal helix "A" together with the helices "B" and "C" are able to fold into a quaternary structure with a second molecule, indicates that these 3 helices constitute a stable folding-scaffold.

20 Second, the biological activity in relation to the vaccine concept has been considered. In general, an inactive construct is preferable with a view to reducing putative toxic effects of the molecules and in general for evaluating the immune response. On the other hand, the optimum neutralising antibodies should theoretically exhibit specificity for the part of IL5 that interacts with the IL5R. This is most likely achieved by immunising with an active variant. Finally, it is not impossible that the biological effect of IL5 on the immune system might act as an enhancer on the immune response, thus improving the overall effect. Based on Applicant's previous experiences with other molecules, however, the majority of "theoretically possible active" constructs is expected to have low or no activity.

35 Therefore, all variants suggested are potentially active but can, if desirable, with relative ease be rendered inactive by hindering the formation of the active dimer or by alterations

in the areas of the "A"- and "D"-helices that are involved in the receptor binding/activation.

In summary, the above considerations of structure conservation and biological activity defines the target areas as any one of loops 1-3 as well as the C-terminal flexible area.

Loop 3 is selected as the primary target area since it is structurally separated from the assumed tri-helical folding scaffold. As it is furthermore possible to produce a biologically active monomer, by elongation of loop 3 (Dickason, 1996), this area holds the possibilities for generating all types of variants: monomer/dimer and active/inactivated.

"Loop 1" is a second area containing a non-helical stretch of a suitable length for substitutions. Variants from this region would theoretically be active only if capable of dimerising, but since the length of the wild-type loop makes it rather flexible it is reasonable to expect a correct folding of the protein after substitution.

Variants containing substitutions in the "loop 2" area will also only be active as dimers. The area that can be substituted is short compared to the inserts and has a central position in the assumed folding scaffold, two characteristics of loop 2 which might be of hindrance to the correct folding of the protein after substitution. On the other hand, loop 2 is situated opposite to the area interacting with the IL5R, resulting in an expected optimum presentation of the wild-type neutralising epitopes if the modified protein is correctly folded.

Finally, inserts in the C-terminal flexible region following "helix D" are proposed. From a protein structure point of view this concept appears fairly safe, but it is likely that modifications in this region will affect both dimerization and biological activity (if the modified protein is dimerized)

since the C-terminal is located in the area of both receptor binding and in the dimer interface.

The amino acid sequence of 10 variants initially constructed according to the above considerations are set forth as SEQ ID NOs: 2-11 and 13-22. Further variants constructed at a later stage are set forth in SEQ ID NOs: 27-59 (including both DNA nucleic acid sequences and amino acid sequences).

10 It should be noted, that all inserts except from the ones according to Example 2 are prepared so as to include flanking amino acid residues that are conserved from hIL5 to mIL5 in order to promote the process of successful transfer of positive constructs from mice to man.

15 In the following examples, positions for substitution are indexed according to the murine amino acid residue sequence numbers; the corresponding human positions are given in parentheses.

#### 20 EXAMPLE 1

*Variants with P2 substituting positions in loop 3 while preserving Cys84(86)*

The P2 epitope (SEQ ID NO: 23) is substituted into loop 3 while avoiding elimination of Cys84(86). These variants (SEQ ID NOs: 2 and 28 (human), where amino acids 87-90 or 88-91 are substituted and 13 and 46 (murine) where amino acids 85-88 or 86-89 are substituted) are potentially active as both monomers (due to the elongation of loop 3) and as dimers. SEQ ID Nos: 28 and 46 are also denoted hIL5.1 and mIL5.1, respectively.



## EXAMPLE 2

*Variants with P2 substituting positions in loop 1 while preserving Cys42(44)*

5

The P2 epitope (SEQ ID NO: 23) is substituted into loop 1 while avoiding elimination of Cys42(44). These variants (SEQ ID NOS: 3 and 36 (human) where amino acids 32-43 or 33-43 are substituted and 14 and 56 (murine) where amino acids 30-41 or 10 31-41 are substituted) are potentially active as dimers only. SEQ ID Nos: 36 and 56 are also denoted hIL5.5 and mIL5.5, respectively.

## 15 EXAMPLE 3

*Variants with P2 substituting positions in loop 2*

The P2 epitope (SEQ ID NO: 23) is substituted into loop 2.

20 These variants (SEQ ID NOS: 4 and 34 (human) where amino acids 59-64 are substituted and 15 and 50 (murine) where amino acids 57-62 are substituted) are potentially active as dimers only. SEQ ID Nos: 34 and 50 are also denoted hIL5.4 and mIL5.4, respectively.

25

## EXAMPLE 4

*Variants with P2 substituting positions in loop 3 while eliminating Cys84(86)*

35 The P2 epitope (SEQ ID NO: 23) is substituted into loop 3 while eliminating Cys84(86). These variants (SEQ ID NOS: 5 and 38 (human) where amino acids 86-91 are substituted and 16 and 54 (murine) where amino acids 84-89 are substituted) are in principle similar to the variants of type #1 (SEQ ID NOS: 2 and 28 and 13 and 46), but the generation of monomer products has been facilitated by inhibiting the formation of disulfide

bridging and adjusting the length of loop 3. SEQ ID Nos: 38 and 54 are also denoted hIL5.6 and mIL5.6, respectively.

#### 5 EXAMPLE 5

*Variants with P2 substituting positions 108-111 (110-113) in the C-terminus*

- 10 The P2 epitope (SEQ ID NO: 23) is substituted into the C-terminal area succeeding helix D. These variants (SEQ ID NOs: 6 and 17) are potentially active as a dimer only.

#### 15 EXAMPLE 6

*Variants with P30 substituting positions in loop 3 while preserving Cys84(86)*

- 20 The P30 epitope (SEQ ID NO: 24) is substituted into loop 3 avoiding elimination of Cys84(86). These variants (SEQ ID NOs: 7 and 40 (human) where amino acids 88-91 or 87-90 are substituted and 18 and 58 (murine) where amino acids 85-88 or 86-89 are substituted) are potentially active both as monomers (due  
25 to the elongation of loop 3) and as dimers. SEQ ID Nos: 40 and 58 are also denoted hIL5.7 and mIL5.7, respectively.

#### EXAMPLE 7

30

*Variants with P30 substituting positions in loop 1 while preserving Cys42(44)*

- 35 The P30 epitope (SEQ ID NO: 24) is substituted into loop 1, avoiding elimination of Cys42(44). These variants (SEQ ID NOs: 8 and 30 (human) where amino acids 32-43 are substituted and 19 and 48 (murine) where amino acids 30-41 are substituted)

are potentially active as dimers only. SEQ ID Nos: 30 and 48 are also denoted hIL5.2 and mIL5.2, respectively.

#### 5 EXAMPLE 8

*Variants with P30 substituting positions in loop 2*

The P30 epitope (SEQ ID NO: 24) is substituted into loop 2.

- 10 These variants (SEQ ID NOs: 9 and 20 where amino acids 59-64 and 57-62 are substituted, respectively) are potentially active as dimers only.

#### 15 EXAMPLE 9

*Variants with P30 substituting positions in the C terminus*

The P30 epitope (SEQ ID NO: 24) is substituted into the

- 20 C-terminal area succeeding helix D. These variants (SEQ ID NOs: 10 and 21 where amino acids 110-113 and 108-111 are substituted, respectively) are potentially active as dimers only.

#### 25 EXAMPLE 10

*Variants with P2 substituting positions 84-89 (86-91) and P30 substituting positions 110-113*

- 30 The P2 epitope (SEQ ID NO: 23) is substituted into loop 3 eliminating Cys84(86) and the P30 epitope (SEQ ID NO: 24) is substituted into the C-terminal area succeeding helix-D. These variants (SEQ ID NOs: 11 and 22) are together with variants of type #12 the only ones containing both epitopes and are poten-
- 35 tially active monomers.

## EXAMPLE 11

*Variants with P30 substituting positions in loop 3 while eliminating Cys84(86)*

5

The P2 epitope (SEQ ID NO: 24) is substituted into loop 3 while eliminating Cys84(86). These variants (SEQ ID NOs: 42 (human) where amino acids 86-91 are substituted and 58 (murine) where amino acids 84-89 are substituted) are in principle similar to the variants of type #6, but the generation of monomer products has been facilitated by inhibiting the formation of disulfide bridging and adjusting the length of loop 3. SEQ ID Nos: 42 and 58 are also denoted hIL5.12 and mIL5.12, respectively.

15

## EXAMPLE 12

*Variants with P2 and P30 substituting positions in loop 3*

20

The P2 (SEQ ID NO: 23) and P30 (SEQ ID NO: 24) epitopes are substituted into loop 3 while preserving Cys84(86). These variants (SEQ ID NOs: 44 and 60 where amino acids 88-91 and 86-89 are substituted, respectively) contain both epitopes and are potentially active monomers. SEQ ID NOs: 44 and 60 are also denoted hIL5.13 and mIL5.13, respectively.

25

## EXAMPLE 13

30

*Choice of expression system*

35

Recombinant IL5 has been expressed in a number of different expression systems including yeast, insect cells and CHO cells (Tavernier et al., 1989).

According to the present invention, one suitable expression system is *E. coli*, based on previous studies reporting the use

of this host for production hIL5 (Proudfoot et al., 1990, Graber et al., 1993). The recombinant protein is expressed as inclusion bodies that are converted into the biologically active dimer upon purification and re-folding (e.g. using the generally applicable refolding methods disclosed in US 5,739,281). The speed and simplicity of *E. coli* expression allows immediate initiation of the production of protein when the genetic constructs are ready, thus facilitating rapid generation of material to establish an *in vivo* proof of the IL5 autovaccine concept.

If for some reason the feasibility is found to be too low (e.g. low yield following re-folding, instability of the products or improved pharmacokinetical parameters related to glycosylation etc), production in yeast could be considered in a further development of the autovaccine.

Recently, promising results have been obtained using the *Drosophila melanogaster* expression system using S<sub>2</sub> cells (available from Invitrogen) and at present this system is the preferred embodiment for expression of the IL5 analogues of the invention.

IL5 variant protein was produced from S2 drosophila cells stably expressing the IL5 constructs. Several different transfection methods were tested, and both Ca<sub>2</sub>PO<sub>4</sub> and Lipofectin were chosen. Two different subclones of S2 cells were used and transfected with Ca<sub>2</sub>PO<sub>4</sub> and Lipofectin, respectively. The two clones were obtained from ATCC and Lars Søndergaard of the University of Copenhagen, respectively. Using both methods suitable stable lines were selected expressing mIL5 and mIL5.1 proteins in the 2-10 mg/L range.

#### Materials & Methods:

35

S2 cells were grown and maintained in Schneider's medium (Sigma) containing 5-10% fetal calf serum (FCS), 0.1% pluronic

F68 (Sigma), penicillin/streptomycin (Life Technologies) grown in shake flasks at 25°C and 120 rpm.

Lipofectin transfections were performed in 250 ml or 1 l shake  
5 flasks. S2 cells were split to  $2.5-3 \times 10^6$ /ml into 50 ml Ex-  
cell 420 (JRH Biosciences) without antibiotics, and grown  
overnight in a 250 ml shake flask. The next morning the Li-  
pofectin reagents were prepared: tube 1) 300-1200 µg plasmid  
DNA containing the gene of interest, plus 15-60 µg pCoHYGRO  
10 hygromycin selection plasmid (20:1 ratio of plasmids) in 15-45  
ml serum and supplement-free medium; tube 2) 1ml Lipofectin in  
5 ml serum and supplement-free medium. After 1 hour at room  
temperature, tubes 1 and 2 were mixed and rested for 15 mi-  
nutes at room temperature before gently adding to S2 cells.  
15 After growing cells overnight new media was added containing  
full supplements plus 150-300 µg/ml Hygromycin.

Transient and stable lines were induced with either 500 µM  
copper sulfate or 10 µM cadmium chloride for 48-72 hours in  
20 serum-free Ex-cell 420 medium (JRH Biosciences).

### Results:

33 stable lines were generated by  $\text{Ca}_2\text{PO}_4$  and 23 by Lipofectin.  
25 The expression yields varied from non-detectable up to 11  
mg/L. The following table summarizes a few of the lines used  
for protein production.

Expression result summary from best mIL5 S2 cell transfections.

30

Plasmid	Construct	S2 cells	Transfection Method	Yield
p612	IL5/His15/mIL5wt	ATCC	$\text{Ca}_2\text{PO}_4$	3.5 mg/L
p767	Bip/His15/mIL5wt	LS	Lipofectin	11 mg/L
p613	IL5/His15/mIL5.1	ATCC	$\text{Ca}_2\text{PO}_4$	2.6 mg/L
p768	Bip/His15/mIL5.1	ATCC	$\text{Ca}_2\text{PO}_4$	0*
p614	IL5/His15/mIL5.5	LS	Lipofectin	0*

\* Expression plasmid contained sequence mutations.

Hence, S2 cells can be transfected by either calcium phosphate precipitation or Lipofectin. Due to the difference in expression level between plasmids p612 and p767, it seems that the Bip signal peptide is a more efficient leader sequence than the endogenous mIL5 leader in S2 cells.

#### EXAMPLE 14

##### 10 *Screening and selection of the modified molecules*

Following expression, the recombinant protein is purified and characterised. The characterisation of the autovaccine candidates will include analytical chromatography, iso-electric focussing (IEF), SDS-PAGE, amino acid composition analysis, N-terminal sequence analysis, mass spectrometry, low angle laser light scattering, standard spectroscopy, and Circular Dichroism to an extent that precisely document the relevant parameters defining the intended protein product.

The His tagged proteins have been purified using a two-step procedure until recently. However, the yield and purity were not as high as expected after the final chelate-step. A new one-step purification procedure has been implied with 3 major advantages achieved: higher yield, higher through-put and higher purity of the final product. Cleavage conditions for removal of the histag have also been established.

##### The two-step IL5 purification procedure:

Expression of the protein is induced by addition of metal ions to the media. These metal-ions have to be removed before application of the protein to the chelate column. Thus, a total of 20 mM EDTA is added to complex the metal-ions and the supernatant is then passed over a SP-sepharose column to capture the protein. After washing to remove unbound protein, bound

protein is eluted by a step-gradient of NaCl. This step serves two purposes: a concentrating step reducing the volume by a factor of 30, and buffer-exchange.

- 5 Relevant fractions (as determined by SDS-PAGE) are pooled and further purified on the metal chelate column.

The protein is applied to a  $\text{Ni}^{2+}$ -charged chelate column and unbound protein washed off. Bound protein is then eluted using  
10 an Imidazole gradient. All fractions, flow-through and EDTA-washes of the column, are then checked by both SDS-PAGE and dot-blot.

Relevant fractions (as determined by SDS-PAGE and dot-blot)  
15 are pooled and dialyzed twice against 10 X volume of PBS, pH adjusted to 6.9.

After filtration, the dialyzed material is concentrated until a suitable concentration is achieved (preferably 1 mg/ml). Fi-  
20 nally, the protein is aliquoted and stored at  $-20^{\circ}\text{C}$ .

The following specific protocol has been applied:

- 1) The received supernatant is centrifuged at  $2500 \times g$  for 15 min  
25 (if infection has occurred, it needs centrifugation at  $22000 \times g$  for 30 min. The supernatant is then filtered using a  $0.45 \mu\text{m}$  filter followed by a  $0.22 \mu\text{m}$  filter (sometimes it is necessary to filter through a  $5 \mu\text{m}$  filter first).  
The supernatant is then mixed 1:1 with buffer A (see step 2)  
30 containing 40 mM EDTA, resulting in a final buffer composition of 0.2 M  $\text{NaH}_2\text{PO}_4$ , 10% glycerol, 20 mM EDTA, pH 6.0
- 2) The filtered supernatant is subsequently applied to a SP-Sepha-  
rose column equilibrated in buffer A. A total of 1-2 L (depen-  
35 ding on protein concentration, the above holds for 1-10 mg IL5 /L) can be applied to an 80 ml column. Flow during application:



1-2 ml/min (usually over night), the flow-through is collected and saved for later analysis. Following application, the column is washed with 2-3 column volumes (CV) of A-buffer until a stable baseline is achieved. Bound protein is eluted using a step gradient: 0-100-500-1000 mM NaCl, fractions of 10 ml are collected, flow is 10 ml/min. Purification is performed at 5°C.

The column is cleaned with 2 CV 1 M NaOH, flow 5 ml/min after each run and re-equilibrated in buffer A.

Buffer A: 0.2 M  $\text{NaH}_2\text{PO}_4$ , 10% glycerol, pH 6.0

Buffer B: 0.2 M  $\text{NaH}_2\text{PO}_4$ , 1 M NaCl, 40 mM Imidazole, 10% glycerol, pH 6.0

The same procedure is used for both wt and variants.

All fractions, starting material and flow-through are tested in dot-blot and SDS-PAGE. The fractions containing IL5 are pooled and further purified using a chelate-column.

#### The one-step IL5 purification procedure:

The supernatant is applied directly to a 70-ml chelate-column charged with  $\text{ZnCl}_2$ . After removal of the unbound material by washing, bound protein (IL5 and contaminants) is eluted by applying a gradient of Imidazole. This method takes full advantage of the His tag giving a one-step purification procedure with a high degree of purity of the final product (>95%).

Relevant fractions (as determined by SDS-PAGE and dot-blot) are pooled and dialyzed twice against 10 X volume of PBS, pH adjusted to 6.9 and concentration of NaCl adjusted to 400 mM.

After filtration, the dialyzed material is concentrated until a suitable concentration is achieved (preferably 1 mg/ml). Finally, the protein is aliquoted and stored at -20°C.

A specific protocol follows the following steps

- 1) The supernatant is filtered through a 0.45  $\mu$ m filter to remove impurities and diluted 1:1 with buffer A.

5 A 70-ml Fast Flow chelate column is rinsed with 5 CV water and then charged with 10 CV 10 mM  $\text{ZnCl}_2$ , pH 7. After equilibration with 5 CV A-buffer, the sample is applied using the pump (flow 10 ml/min). The flow-through is collected and saved for later analysis. Bound protein is eluted using an Imidazole-gradient going from 0 to 250 mM Imidazole over 30 CV. Finally, the column is stripped by 5 CV of buffer C. Fractions of 10 ml are collected.

Buffers:

A: 20 mM  $\text{NaH}_2\text{PO}_4$ , 0.5 M NaCl, 10% glycerol, pH 7.

B: 20 mM  $\text{NaH}_2\text{PO}_4$ , 0.5 M NaCl, 10% glycerol, pH 7, 0.25 M imidazole

C: 20 mM  $\text{NaH}_2\text{PO}_4$ , 0.5 M NaCl, 0.1 M EDTA pH 7.0.

All fractions, flow-through and starting material is tested in SDS-PAGE.

- 2) The purest fractions (as determined by SDS-PAGE) containing IL5 are pooled (50  $\mu$ l are saved for later analysis) and dialyzed twice against 10 X volume of PBS, pH adjusted to 6.9. at 6°C, MWCO 12-14 kDa. The dialysate is filtered through a 0.22  $\mu$ m filter (50  $\mu$ l is saved for later analysis) and  $A_{280}$  is measured using dialysis-buffer (filtered through 0.45  $\mu$ m) as reference. The volume before and after dialysis is measured and samples showing the dialysis/concentrating step are saved for later analysis by SDS-PAGE (after step 3).

- 3) NaCl is added to the dialyzed protein until a total concentration of 400 mM and it is then concentrated using either an Amicon apparatus (for volumes larger than 50 ml) or Vivaspin concentrating device (for 10-50 ml). In both cases, the membrane is saturated with 10 ml PBS-buffer before the sample is applied. The sample should be concentrated until a concentration of preferably 1 mg/ml is achieved (as measured by  $A_{280}$ ). The dialyzed, concentrated sample is filtered through a 0.22  $\mu$ m filter

and marked with an E-nr. The  $A_{280}$  is measured using the flow-through as reference.

All samples from the dialysis and concentrating step are analyzed by SDS-PAGE and Coomassie-stained. The purified protein is stored frozen in aliquots and a sheet describing the sample is filed in the "IL5-protein"-folder.

The above-described procedure gives a protein with a purity of approximately 90-95%, still containing the His Tag.

When sequenced, both IL5wt and variant IL5.1 gave the expected N-terminal sequences including the His Tag.

The purification procedure referred to above has been implemented in the following specific setup:

- 1) The pooled fractions from the SP-sepharose column are filtered through a 0.45  $\mu$ m filter to remove impurities.

A 5-ml HiTrap chelate column (use only dedicated columns) is rinsed with 15 ml water (using a syringe) and then charged with 15 ml 0.1 M  $\text{NiSO}_4$  and washed with 15 ml water. The column is connected to the Äkta-system and equilibrated with 2-3 CV A-buffer. The sample is applied using either the loop or pump - depending on the volume (flow 4 ml/min), the flow-through is collected and saved for later analysis. Bound protein is eluted using an Imidazole-gradient going from 0 to 500 mM Imidazole over 20 CV. Fractions of 5 ml are collected. Finally, the column is stripped using 5 CV of buffer B2.

Buffer A: 0.2 M  $\text{NaH}_2\text{PO}_4$ , 0.5 M NaCl, 10 % glycerol, pH 5.0

Buffer B1: 0.2 M  $\text{NaH}_2\text{PO}_4$ , 0.5 M NaCl, 0.5 M Imidazole, 10 % glycerol, pH 5.0

Buffer B2: 50 mM Na-acetate, 0.5 M NaCl, 0.1 M EDTA, 10 % glycerol, pH 4.5

All fractions, flow-through and starting material are tested in dot-blot, all relevant fractions are tested in SDS-PAGE.

2) The purest fractions (as determined by SDS-PAGE) containing IL5 are pooled (save 50  $\mu$ l for later analysis) and dialyzed twice against 10 X volume of PBS, pH adjusted to 6.9. at 6 °C, MWCO 12-14 kDa. The dialysate is filtered through a 0.22  $\mu$ m filter (save 50  $\mu$ l for later analysis) and  $A_{280}$  is measured using filtered dialysis-buffer as reference. The volume before and after dialysis is measured and samples showing the dialysis/concentrating step are saved for later analysis by SDS-PAGE (after step 3).

3) After addition of extra NaCl up to a final concentration of 400 mM, the dialyzed protein is concentrated using either an Amicon apparatus (for volumes larger than 50 ml) or Vivaspin concentrating device (for 10-50 ml). In both cases, the membrane is saturated with 10 ml PBS buffer before the sample is applied. The sample should be concentrated until a concentration of preferably 1 mg/ml is achieved (as measured by  $A_{280}$ ). The  $A_{280}$  is measured using the flow-through as reference. The dialyzed, concentrated sample is filtered through a 0.22  $\mu$ m filter and marked with an E-nr.

All samples from the dialysis and concentrating step are analyzed by SDS-PAGE and Coomassie-stained. The purified protein is stored frozen in aliquots.

Other purification procedures that have been evaluated are:

*Zn<sup>2+</sup>-chelate purification:* Elution of the protein using an increasing Imidazole gradient has proved very efficient as the wt-protein binds strongly to the column. The *Drosophila* supernatant can be directly applied and after washing, the IL5wt can be eluted by Imidazole. The column is charged with 10 CV 10 mM  $ZnCl_2$ , and washed with water. The pH of the binding and elution buffers has to be above 6.5 as otherwise the  $ZnCl_2$  will precipitate.

Con A affinity chromatography is under investigation. The possibility of using the glycosylation present on IL5 as an affinity-tag and elute by application of a monosaccharide-analog would be interesting since it could be applied to the non-His 5 tagged constructs as well.

#### Removal of Histag:

Removal of the 15 aa His tag (SEQ ID NO: 25) has been performed according to suppliers (Unizyme) instructions:

The purified and dialyzed/concentrated His tagged IL5 is de-His tagged by the sequential addition of two enzymes, DAP1 and Glutamine cyclotransferase. DAP1 removes two amino acids from the free N-terminus while the QCT

The enzyme needs to be activated first:

9 µl HT-DAP1 (10 U/ml) is mixed with 9 µl 20 mM cysteamine-HCl. After 5 min incubation at room temperature, a total 108 µl HP-GCT (100 U/ml) and 54 µl TAGZyme buffer is added. This must be used within 15 min.

This portion will digest 1 mg of His tagged protein.

The His tagged protein is mixed with 150 µl activated enzyme and incubated at 37°C for 120 min. Samples are withdrawn for SDS-PAGE analysis (10 µl) after 0, 10, 30, 60 and 120 min. The samples are put on ice to stop the digestion.

#### Buffers:

1. TAGZyme buffer: 20 mM NaPO<sub>4</sub> buffer, pH 7.5; 150 mM NaCl
2. 20 mM Cysteamine-HCl

The digested protein (as determined from SDS-PAGE analysis or N-terminal sequencing) is applied to a 1-ml Ni-chelate column equilibrated in PBS. Everything is collected.

The flow-through from the application is saved for later analysis. The column is eluted by addition of 3 CV PBS, fractions of 0.5 ml are collected. The column is cleaned by washing with 2 CV 0.5 M Imidazole, and fractions are saved for analysis.

All fractions are tested in SDS-PAGE, and fractions containing IL5 are pooled and  $A_{280}$  is measured using PBS as reference. Finally, the protein is concentrated using a Vivaspin concentrating device until a concentration of 1 mg/ml is achieved.

Removal of His tag has been performed in small-scale experiments (0.1-1 mg) and has not been up-scaled. It should be noted that removal of the tag requires an unblocked and non-modified N-terminus.

The His tagged protein is incubated with two enzymes, a dipeptidyl amino peptidase which removes two amino acids at a time and a glutamic acid cyclotransferase which catalyze the conversion of a glutamic acid into a pyro-glutamic acid. This conversion blocks further degradation by the dipeptidyl amino peptidase. The digestion mixture is then passed through a chelate column which should retain the enzymes (which are His tagged), contaminating proteins binding to the column and non-degraded or partially degraded protein. The de-tagged protein passes the column and is collected in the flow-through. After a second digestion with an enzyme that removes the pyro-glutamic acid, the protein is again passed over a chelate-column to remove the second enzyme. It is expected that the protein needs to be concentrated again at this final stage.

#### General observations:

The pI of UniHis-IL5wt is 9.5 and the optimum pH-value for the protein seems to be 6.5-7.0 (has not been investigated tho-

roughly). A NaCl-concentration of 400 mM seems to stabilize the protein during concentration.

#### EXAMPLE 15

5

##### *In vitro screening*

The primary *in vitro* screening will be in the form of an enzyme-linked immunosorbent assay (ELISA): A competitive ELISA  
10 towards wild-type IL5 provides an estimate of the presence of relevant B-cell epitopes in the modified IL5 constructs before introduction thereof into animals.

A conventional ELISA assay can be used to measure titres of  
15 auto-antibodies in the serum of vaccinated animals. Antibodies (both mono-specific and monoclonal) towards the human as well as towards the murine IL5 are commercially available from R&D Systems, 614 McKinley Place NE, Minneapolis, MN 55413, USA.

20 The biological activity of the product and/or the neutralising capacity of induced auto-antibodies can be tested in an IL5 bioassay. Previously reported examples of such bioassays are: Assessment of IL5 induced proliferation of TF1 cells (for human IL5) and assessment of IL5 induced proliferation of BCL1  
25 cells or B13 B cells (for murine IL5) (Callard & Gearing 1994, Dickason et al., 1994).

The effect on airway responsiveness of the autovaccine can also be tested in an *in vitro* assay wherein the trachea from  
30 vaccinated mice are removed and placed on a hook in an organ bath. The tension of the trachea after histamine challenge is measured (van Oosterhout et al., 1995).

Consequently, in order to be able to determine the biological  
35 activity of recombinant mIL5 (and mIL5 AutoVac) protein samples, a cellular bioactivity assay for murine IL5 is being established. The assay is based on the ability of the B cell lymphoma line BCL1 to proliferate in response to mIL5 added to

the culture medium. Two different BCL1 clones were obtained from ATCC, BCL1 clone 5B1b (ATCC CRL-1669) and BCL1 clone CW13.20.3B3 (ATCC TIB-197).

5 In a typical BCL1 proliferation experiment, the cells are plated in complete RPMI medium supplemented with fetal calf serum (FCS) in microtiter plates and incubated with dilution series of murine IL5. Proliferation of the BCL1 cells is measured by incorporation of tritiated thymidine. Several optimization experiments have been performed using dilution series of purchased recombinant mIL5 (R&D Systems) for stimulation. The variable parameters include: incubation intervals, <sup>3</sup>H-thymidine pulsing intervals, numbers of cells plated per well, fetal calf serum (FCS) concentrations and concentrations of added mIL5. Dose-dependent proliferation of the BCL1 cells with a maximal proliferation of about 3 times the background (BCL1 cells with no mIL5 added) has been obtained.

The BCL1 assay has been used to determine the biological activity of the following samples expressed from Drosophila S2 cells and purified as described above: HIS-mIL5wt material (E1320), HIS-mIL5wt material (E1422), HIS-mIL5.1 material (E1396) and an "S2-background-preparation" (E0016). The proliferation in response to one HIS-mIL5wt (E1320) preparation was significantly higher than the proliferation in response to the "S2-background-preparation", whereas the mIL5.1 variant and one wild type preparation (E1422) were determined as biologically inactive.

30 Ongoing work includes inhibition of the BCL1 proliferation with anti-mIL5, and the anti-mIL5 monoclonal antibody TRFK5 is used for optimization studies. This is done in order to use this assay to determine the ability of anti-mIL5 antisera from immunised mice to inhibit the biological activity of mIL5.



## EXAMPLE 16

5 *In vivo models*

For measuring the *in vivo* effect of the autovaccine, well-known animal models for asthma exists. Normally, the animal is sensitised with a compound (allergen/antigen) and after  
10 challenge with the aerosolised compound, broncho-constriction (airway conduction) is measured using a body plethysmograph. The eosinophil cell counts in the BAL fluid are also determined.

15 Several of the studies investigating the effect of anti-IL5 mAb's have been successfully performed in mice. Against use of the murine model speaks the fact the IL5 acts as a B-cell growth factor, rendering possible interference with the murine antibody response. However, as shown in a study using IL5  
20 knock-out mice, the T-cell dependent antibody response against ovalbumin as well as cytotoxic T-cell development appeared normal (Kopf et al., 1996). As the mouse is also the most practical and economical model in comparison to guinea pigs or monkeys, the ovalbumin sensitised Bal/c mice model of  
25 asthma/airway hypersensitivity as used by Hamelman et al. (1997) will be used.

If, however, the effect of IL5 on B-cells in the murine model turns out to be a problem, the use of other suitable animal  
30 models known in the art will be applied.

## EXAMPLE 17

*Preparation of DNA constructs encoding murine IL5 and variants thereof*

5

Construction of variants in pcDNA3.1+:

Insertion of P2 and P30 epitopes into wildtype mIL5 was done by SOE-PCR with overlapping primers containing the epitope sequences. Wildtype mIL5 gene including leader sequence (SEQ ID NO: 63), cloned into pcDNA3.1+ with consensus Kozak sequence (obtaining plasmid p815), was used as template for the PCR reactions. The resulting fragments were digested with *NheI* and *NotI*, purified and cloned into p815 was used as template for the PCR.

15

Cloning of variants into pMT Drosophila vector with BiP leader and UNI-His tag:

Wildtype mIL5 was cloned into the pMT *Drosophila* expression vector series (Invitrogen) by generating a PCR fragment with mIL5 specific primers containing appropriate restriction sites and, in addition, containing sequences encoding a *Drosophila* Kozak like sequence followed by the *Drosophila* BiP leader sequence followed by a sequence encoding a UNI-HIS tag (SEQ ID NO: 25) fused to the 5' end of the sequence encoding mature mIL5. Wildtype mIL5 cDNA sequence was used as template. The resulting fragment was digested with *EcoRI* and *NotI* and was subsequently cloned into the pMT/V5-HisA vector (Invitrogen). The resulting plasmid (p818) was used for cloning of epitope containing variants into pMT. These were cloned by digesting the variants made in pcDNA3.1+ with *SacI* and *NotI* and cloning the resulting fragments into p818.

Cloning of variants into pAC5:

Wildtype and variants of mIL5 were cloned into the pAC5 constitutive Drosophila expression vector by digestion of variants in pMT with *EcoRI* and *NotI* and cloning the resulting 5 fragments into the pAC5.1/V5-HisA vector (Invitrogen).

## EXAMPLE 18

10 *Preparation of DNA constructs encoding human IL5 and variants thereof*

Five lines of plasmids are contemplated containing unmodified IL5 and all or some of the nine IL5 variants. The lines include: 1) human IL5 for DNA vaccination in the pCI vector suited for expression in human cells, 2) human IL5 with the BiP leader sequence and a 15 aa His tag (SEQ ID NO: 25, obtained from UNIZYME in Hørsholm, Denmark. The tag is termed "UNI" or "UNI-His tag" herein) in the pMT/V5/HIS vector for inducible expression in Drosophila, 3) as in 2 but without the His tag, 4) as in 3 but with murine IL5 and 5) human IL5 with the DAPI leader sequence and the 15 aa HIS tag in the vector pVL1393 for expression in the baculo-virus system.

Plasmids for DNA-vaccination in the pCI vector:

Name	ref #	Strain #	Epitope
hIL5 (pCI)	p888	MR#1237	none
hIL5.1 (pCI)	p889	MR#1238	P2, Loop 3
hIL5.2 (pCI)	p890	MR#1239	P30, Loop 1
hIL5.3 (pCI)	p891	MR#1240	P30, Loop 2
hIL5.4 (pCI)	p892	MR#1241	P2, Loop 2
hIL5.5 (pCI)	p893	MR#1242	P2, Loop 1
hIL5.6 (pCI)	p894	MR#1243	P2, Loop 3
hIL5.7 (pCI)	p895	MR#1244	P30, Loop 3
hIL5.12 (pCI)	p896	MR#1245	P30, Loop 3
hIL5.13 (pCI)	p897	MR#1246	P2 and P30, Loop 3

5 Plasmids for human IL5 expression in Drosophila with the UNI-HIS tag and BiP leader sequence in pMT/V5/HIS :

Name	Ref #	Strain #	Epitope
hIL5m-UNI-BiP (pMT/V5-HisA)	p899	MR#1247	none
hIL5.1m-UNI-BiP (pMT/V5-HisA)	p900	MR#1248	P2, Loop 3
hIL5.2m-UNI-BiP (pMT/V5-HisA)	p901	MR#1249	P30, Loop 1
hIL5.3m-UNI-BiP (pMT/V5-HisA)	p929	MR#1277	P30, Loop 2
hIL5.4m-UNI-BiP (pMT/V5-HisA)	p902	MR#1250	P2, Loop 2
hIL5.5m-UNI-BiP (pMT/V5-HisA)	p903	MR#1251	P2, Loop 1
hIL5.6m-UNI-BiP (pMT/V5-HisA)	p904	MR#1252	P2, Loop 3
hIL5.7m-UNI-BiP (pMT/V5-HisA)	p905	MR#1253	P30, Loop 3
hIL5.12m-UNI-BiP (pMT/V5-HisA)	p906	MR#1254	P30, Loop 3
hIL5.13m-UNI-BiP (pMT/V5-HisA)	p907	MR#1255	P2 and P30, Loop 3

10 Plasmids for human IL5 expression in Drosophila with the BiP leader sequence, but without the UNI-HIS tag in pMT/V5/HIS:

80

Name	Ref #	Strain #	Epitope
hIL5m-BiP (pMT/V5-HisA)	p908	MR#1256	none
hIL5.1m-BiP (pMT/V5-HisA)	p909	MR#1257	P2, Loop 3
hIL5.2m-BiP (pMT/V5-HisA)	p921	MR#1269	P30, Loop 1
hIL5.3m-BiP (pMT/V5-HisA)	p922	MR#1270	P30, Loop 2
hIL5.4m-BiP (pMT/V5-HisA)	p923	MR#1271	P2, Loop 2
hIL5.5m-BiP (pMT/V5-HisA)	p924	MR#1272	P2, Loop 1
hIL5.6m-BiP (pMT/V5-HisA)	p925	MR#1273	P2, Loop 3
hIL5.7m-BiP (pMT/V5-HisA)	p926	MR#1274	P30, Loop 3
hIL5.12m-BiP (pMT/V5-HisA)	p927	MR#1275	P30, Loop 3
hIL5.13m-BiP (pMT/V5-HisA)	p928	MR#1276	P2 and P30, Loop 3

Plasmids for murine IL5 expression in Drosophila with the BiP leader sequence, but without the 15 aa His tag in pMT/V5/HIS:

5

Name	ref #	Strain #	Epitope
mIL5m-BiP (pMT/V5-HisA)	p918	MR#1266	none
mIL5.1m-BiP (pMT/V5-HisA)	p919	MR#1267	P2, Loop 3
mIL5.2m-BiP (pMT/V5-HisA)	p920	MR#1268	P30, Loop 1

Plasmids for human IL-5 expression in the baculo-virus system with the UNI-HIS tag and DAP1 leader sequence pVL1393 in

10 pVL1393:

Name	Ref #	Strain #	Epitope
hIL5m-UNI-DAP1 (pVL1393)	p916	MR#1264	none
hIL5.1m-UNI-DAP1 (pVL1393)	p917	MR#1265	P2, Loop 3

## EXAMPLE 19

*DNA immunization studies*

5 Generation of vectors encoding mIL5wt, mIL5.1 and mIL5.5 with Kozak sequences for DNA vaccination experiments:

DNA fragments encoding mIL5wt, mIL5.1 and mIL5.5 including the natural leader sequence (SEQ ID NO: 63) were inserted into pcDNA3.1 thus yielding new plasmids p521, 522, and p523. In order to enhance expression of cDNA in mammalian cells, Kozak consensus sequences were inserted upstream of the coding sequences using PCR technology. PCR reactions were performed using p521, p522 and p523 as templates and a forward primer encoding the Kozak sequence immediately upstream of the mIL5 leader start codon. Purified PCR products were cloned into pcDNA3.1+ vector using restriction endonucleases *Bam*HI and *Not*I. The resulting plasmids p815, p816 and p817, respectively, were verified by DNA sequencing. All other plasmids used for DNA vaccination experiments were constructed using the p521 plasmid as starting material.

In vitro translation of DNA vaccination plasmids using Promega Kit:

A commercial kit using rabbit reticulocyte extract to generate in vitro translated protein product plasmid DNA, has previously been successfully used in our lab to monitor protein expression from pcDNA plasmid encoding e.g ovalbumin cDNA. Murine IL5 DNA vaccination plasmids were added to the kit reagents according to the standard procedure. However, several attempts to detect expressed mIL5 material on autoradiograms failed whereas positive controls worked. Results from COS cell transfections and DNA vaccination shows that the gene products are expressed, so we did not investigate these technical problems further.

Transient transfection of COS cells with DNA vaccination plasmids to determine expression levels:

In order to monitor the transfection/expression efficiency of the plasmids used for DNA vaccination experiments, a transient transfection assay using COS cells was established. COS cells were trypsinized and plated in DMEM medium supplemented with 10 % FCS in T25 culture flasks. The cells were transfected at day 2 using the Dotap kit (Roche Diagnostics) and harvested at day 5. Culture supernatant, whole cell lysate and membrane enriched preparations were tested in Western blotting to detect anti-mIL5 reactive expression product. The anti-mIL5 reactive product in the cell preparations consistently migrated as 2-3 separate bands of 21-28 kD in SDS-PAGE, whereas the MW of the mIL5 monomer used as standard (expressed in baculovirus, R&D Systems) is only 15-18 kD. Using non-denaturing circumstances, the 21-28 kD substances form dimers so we believe the material is mIL5, possibly in several differently glycosylated forms. DNA vaccination results (see below) clearly support this conclusion.

DNA vaccination of mice using murine IL5 AutoVac constructs:

A DNA vaccination study was performed in order to investigate whether antibody responses specific for murine IL5 can be induced by immunising mice with naked plasmid DNA encoding 8 different murine IL5 mutants. Since IL5 previously has been reported to play a role in B cell differentiation, it is essential to demonstrate that anti-mIL5 autoantibodies can be generated in mice and B cell tolerance to mIL5 can be broken.

The general setup of the DNA vaccination experiments use either C3H/He mice (H-2<sup>k</sup>) or Balb/cA mice (H-2<sup>d</sup>), 6-8 weeks old divided into groups of 5 mice each. At days 0, 14, 28, 42, 62 and 76 the mice were anaesthetized using hypnorm/dormicum s.c.

and injected with expression plasmids encoding ovalbumin (control), mIL5wt (wild type), or the mIL5 variants to be tested. The DNA material was prepared using endofree GigaPrep kits (Qiagen) and dissolved at 1 µg/ml in 0.15 M NaCl or 0.15 M NaCl containing 0.1% bupivacaine. 100 µl material was injected i.d. in each mouse at the lower back distributed at two injection sites. Prebleeds were obtained at day minus 2, and the test bleedings were obtained at weeks 3, 5, 8 and 11. Sera were isolated by centrifugation and stored at -20°C until testing in ELISA for reactivity against purified ovalbumin and mIL5 proteins.

A Typical result of a DNA vaccination experiment is shown in Fig. 4. According to the general setup described above, 40 Balb/cA mice were immunized with ovalbumin control plasmid, mIL5wt encoding plasmid or plasmids encoding the mIL5 AutoVac variants mIL5.1 or mIL5.5. In this experiment, 9 out of 9 mice immunized with ovalbumin encoding plasmid developed anti-ovalbumin antibodies, whereas no anti-ovalbumin response was induced in mice receiving the mIL5 wild type or mIL5 variant encoding DNA. Injection of mIL5wt encoding plasmid did not give raise to an anti-mIL5 response, whereas the B cell tolerance to mIL5 was broken in 4 out of 10 mice immunized with mIL5.1 plasmid and 7 out of 9 mice immunized with mIL5.5 encoding plasmid DNA.

The main result of the whole series of DNA vaccination experiments is summarized in the table below. The number of responders within an immunisation group differs between the different mIL5 AutoVac constructs and is dependent on the mouse strain. Clearly, the mIL5.2 AutoVac construct is superior to the other variants, being able to induce anti-mIL5 antibody responses in both mouse strains with a penetrance of 100 %.



This plasmid (p820) also gave the highest expression levels in the COS transfection assay.

Another example to emphasize is the apparent MHC restriction

5 seen when using mIL5.4 encoding plasmid DNA as immunogen.

Whereas only 1/10 C3H/Hen mice responds to the DNA vaccine, 9 out of 10 Balb/cA mice are responders. The opposite phenomenon (although not quite as pronounced) is seen with the mIL5.6 construct. The mIL5.2 DNA vaccine, however, seem to be

10 promiscuously immunogenic.

	OVAwt-pVax	mIL5wt-pcDNA	mIL5.1-pcDNA	mIL5.2-pcDNA	mIL5.4-pcDNA
Balb/cA	28/28	0/28	4/10	9/10	9/10
C3H/Hen	29/29	0/30	3/10	10/10	1/10

	mIL5.5-pcDNA	mIL5.6-pcDNA	mIL5.7-pcDNA	mIL5.12-pcDNA	mIL5.13-pcDNA
Balb/cA	7/9	0/10	2/10	0/10	0/10 *
C3H/Hen	5/10	6/10	2/10	2/10	2/10 *

15 Summary of the result of DNA vaccination of 280 mice. 6 mice died during the experiment for reasons not connected to the effects of the DNA vaccination. The number of responders (with high or intermediate anti-mIL5 titers) is shown in relation to the total number of mice within each immunization

20 group. \*) bleedings obtained at day 55. All the other bleedings were obtained at day 77.

Another feature to mention is the tendency of mIL5 variants with the foreign T helper epitope inserted in mIL5 loop1 to be

25 stronger DNA vaccination immunogens than variants with the T helper epitope inserted in loop 3. This could be due to the relatively high expression levels. The only loop 2 variant tested, mIL5.4-pcDNA is only a strong immunogen in the Balb/cA strain, as mentioned above.

Further characterization of the antibody responses induced by DNA vaccination:

ELISA experiments were set up in order to determine whether antibodies specific for the inserted T helper epitope could be detected in anti-mIL5 positive mice. For each immunisation group, sera from anti-mIL5 positive mice were pooled and tested for reactivity against P2 or P30 peptides which had been immobilised in AquaBind microtiter plates. Antisera induced by DNA vaccination against mIL5.2 in both mouse strains clearly contained reactivity against the inserted P30, whereas none of the other antisera were reactive with P2 or P30. This is probably connected to the higher antibody titers and penetration that is generally observed with the mIL5.2 DNA vaccination construct. It should be mentioned that using this ELISA setup we were able to detect anti-P2 reactivity in antisera induced against mIL5.1.

The positive anti-mIL5 antiserum pools from the DNA vaccinated mice were also tested in a competitive ELISA for their ability to inhibit the interaction between soluble native murine IL5 and monoclonal antibodies TRFK4 or TRFK5, which are both neutralizing antibodies. Dilution series of anti-mIL5 antiserum pools were preincubated with soluble native mIL5 and the sample was added to ELISA plates coated with catching antibody TRFK5. Bound murine IL5 (which was not absorbed by the antisera) was next visualised using layers of biotinylated TRFK4 and subsequently horse radish peroxidase labeled streptavidin. Not all the anti-mIL5 positive antisera induced by DNA vaccination could inhibit the interaction between soluble mIL5 and TRFK4 or TRFK5. The antiserum with the highest TRFK4/5 inhibiting capability was from C3H/Hen mice immunized with mIL5.2 encoding DNA. It has not been tested whether the observed differences in inhibition is a direct measure of titer differences or it is connected to the fine specificity of the dif-

ferent antisera. Most likely, it is a combination of these two factors.

Animal model of eosinophilia in mIL5 AutoVac DNA immunized

5 mice:

40 DNA vaccinated mice were chosen for testing in an animal model of eosinophilia: 10 Balb/cA mice immunized with mIL5wt DNA, 10 Balb/cA mice immunized with mIL5.2 DNA, 10 C3H/Hen mice immunized with mIL5wt DNA and 10 C3H/Hen mice immunized with mIL5.2 DNA. A sensitization/challenging regimen with ovalbumin to induce eosinophilia was given to in each of these mice. The mice were sensitized with subcutaneous injections of 50 µg ovalbumin (OVA) in 0.9 % saline mixed 1:1 with Adjuphos once per week for three weeks. Four days after the last OVA sensitization the mice were challenged intranasally with 12.5 µg OVA in 0.9 % saline every other day for a total of 3 challenges. Bronchoalveolar lavage fluid (BALF) was collected one day after the last sensitization by cannulating the tracheae and washing the airway lumina with 1 ml PBS.

20

Approximately 30,000-60,000 BALF cells were spun unto slides at 1,500 rpm for 20 minutes. The slides were dried overnight and stained for 2.5 minutes with May-Grunwald stain (Sigma), washed for 4 minutes in tris buffered saline, stained for 20-30 minutes with Geimsa stain (1:20 with ddH2O; Sigma) and briefly rinsed with ddH2O. Stained slides were dried overnight and cell types were identified using light microscopy. Approximately 100-200 cells were counted per slide and 3 slides were counted per mouse. The eosinophil counts were expressed as the number of eosinophils per 100 cells counted. In mIL5.2 DNA vaccinated C3H/Hen mice, the induction of lung eosinophilia was significantly down-regulated compared to the wild type mIL5wt DNA vaccinated group (mIL5.2 DNA:  $14.6 \pm 8.9$  eosinophils per 100 cells; mIL5wt DNA:  $51.1 \pm 9.9$  eosinophils per

100 cells). However, in the Balb/cA strain, there was no significant difference in eosinophil counts between the immunization groups (mIL5.2 DNA:  $23.3 \pm 6.8$  eosinophils per 100 cells; mIL5wt DNA:  $27.7 \pm 9.3$  eosinophils per 100 cells). A possible explanation is that Balb/cA mice are only weakly susceptible to the model. This is supported by anti-ovalbumin ELISA data showing that one week before the BALF collection the anti-ovalbumin titers in serum from the Balb/cA mice were lower than in serum from C3H/Hen. The Balb/cJ substrain is reported to be susceptible to the OVA sensitization/challenge model.

#### EXAMPLE 20

##### *Protein vaccination study*

Balb/c J mice were immunized with murine IL5 (mIL5) protein and subjected to an ovalbumin intranasal model that induces eosinophils in the lungs of treated mice. Both the UniHis-mIL5 and the UniHis-mIL5.1 proteins induced antibodies that cross-react with mIL-5 made in sf9 cells from R&D Systems. The eosinophilia model induced high numbers of eosinophils in the OVA control group and the UniHis-mIL5.1 groups, while the numbers of eosinophils were reduced in both the PBS group and the UniHis-mIL5 group. This result led us to believe that the groups may have been mixed.

##### Materials & Methods:

UniHis-mIL-5	E1320 & E01397
UniHis-mIL-5.1	E01337 & E01396

### Immunizations:

6-8 week old female Balb/c J (M&B) mice were immunized with either 1) nothing, 2) PBS, 3) UniHis-mIL5, or 4) UniHis-mIL-5.1 in Complete Freund's Adjuvant (CFA; Sigma) and boosted 3 times at three week intervals with antigen in Incomplete Freund's Adjuvant (IFA; Sigma). Sera was collected and tested in an ELISA 10 days after each boost.

### 10 ELISAs:

#### Anti-UniHis-mIL5 ELISA:

Sera were obtained at days 32 (bleed 1) and 54 (bleed 2) after 2 and 3 immunizations, respectively. Polystyrene microtiter plates (Maxisorp, Nunc) were coated with purified HIS-mIL5wt (0.1 µg/well, E1320). The reactivities of diluted sera added to the wells were visualised using a goat anti-mouse secondary antibody. OD490 readings of control sera from mice immunized with PBS in Freund's adjuvants were subtracted from the OD490 readings of the test samples.

#### Anti-mIL5 ELISA:

Sera were obtained at day 75 (bleed 3). Polystyrene microtiter plates (Maxisorp, Nunc) were coated with purchased mIL5 (0.1 µg/well, R&D cat. no. 405-ML). The reactivities of 1:1000 diluted sera added to the wells were visualised using a goat anti-mouse secondary antibody. The reactivity of TRFK5 (2 µg/ml) was visualised using a rabbit anti-rat secondary antibody.

### 30

#### Competitive ELISA:

Dilutions of antisera were preincubated with soluble IL5 for 1 hour and added to polystyrene microtiter plates (Maxisorp, Nunc) which were coated with catching antibody TRFK5. Bound

mIL5 was visualised using biotinylated TRFK4 and a HRP labelled goat anti-mouse secondary antibody.

Anti-P2 ELISA:

5 Pools of antisera from HIS-mIL5wt, HIS-mIL5.1 or PBS immunised mice were tested for reactivity against P2 peptide in ELISA. Specialized microtiter plates (Aquabind, M&E Biotech) were coated with 0.5 µg/well synthetic P2 peptide. The reactivities of diluted sera added to the wells were visualised using a HRP  
10 labelled goat anti-mouse secondary antibody (1:2000, Dako).

Anti-UniHis ELISA:

Pools of antisera from HIS-mIL5wt, HIS-mIL5.1 or PBS immunised mice were tested for reactivity against HIS-tag peptide  
15 (UNIZYME) in ELISA. Specialized microtiter plates (AquaBind, M&E Biotech) were coated with 0.5 µg/well synthetic HIS-tag peptide. The reactivities of diluted sera added to the wells were visualised using a HRP labelled goat anti-mouse secondary antibody (1:2000, Dako).

Anti-S2 background protein ELISA:

Pools of antisera from HIS-mIL5wt, HIS-mIL5.1 or PBS immunised mice were tested for reactivity against S2 background preparation in ELISA. Polystyrene microtiter plates (Maxisorp, Nunc)  
25 were coated with 0.1 µg/well S2 background preparation. The reactivities of diluted sera added to the wells were visualised using a HRP labelled goat anti-mouse secondary antibody (1:2000, Dako).

30 Anti-BSA ELISA:

Pools of antisera from HIS-mIL5wt, HIS-mIL5.1 or PBS immunised mice were tested for reactivity against BSA in ELISA. Polystyrene microtiter plates (Maxisorp, Nunc) were coated with 10 µg/well BSA (Intergen). The reactivities of diluted sera added

to the wells were visualised using a HRP labelled goat anti-mouse secondary antibody (1:2000, Dako).

#### Eosinophilia Model:

5 Balb/c J mice were sensitized with subcutaneous injections of 50 µg ovalbumin (OVA) in 0.9% saline mixed 1:1 with Adjuphos as alum adjuvant. OVA immunizations were repeated once per week for four weeks. One week after the last OVA sensitization, the mice were challenged with 12.5 µg OVA in 0.9% saline  
10 intranasal every other day for a total of 3 challenges. Bronchoalveolar lavage fluid (BALF) was collected one day after the last sensitization by cannulating the tracheae and washing the airway lumina with 1 ml 0.9% saline, or PBS.

#### BAL staining:

15 Approximately 30,000-60,000 BALF cells were spun unto slides at 1,500 rpm for 20 minutes. The slides were dried overnight and stained for 2.5 minutes with May-Grunwald stain (Sigma), washed for 4 minutes in TBS, stained for 20-30 minutes with  
20 Giemsa stain (1:20 with ddH<sub>2</sub>O; Sigma) and briefly rinsed with ddH<sub>2</sub>O. Stained slides were dried overnight and cell types were identified using light microscopy. Approximately 100-200 cells were counted per slide and 3 slides were counted per mouse.

#### 25 Results:

##### Detection of anti-mIL5 antibodies:

A series of ELISA experiments were performed in order to investigate whether antibody responses specific for murine IL5  
30 were induced in mice immunized with HIS-mIL5wt and HIS-mIL5.1 protein material. First, it was determined if antibodies against the HIS-mIL5wt immunization material were elicited by testing dilutions of antisera from individual mice on ELISA plates coated with the HIS-mIL5wt material. It was found that

already by bleed one, all mice had developed high-titered antibody responses against the HIS-mIL5wt material (E1320, expressed from Drosophila S2 cells and purified) which was estimated to be approximately 95% pure.

5

This result is not a firm confirmation that the antisera cross-reacts with murine IL5. In this setup, reactivities would also be detected against impurities from the Drosophila S2 cells, the S2 medium (which contain e.g. BSA from fetal calf serum, the HIS-tag as well as denatured mIL5 B cell epitopes. To demonstrate, that the antibodies induced contain reactivities against native murine IL5, the sera were tested in ELISA plates coated with mIL5 purchased from R&D systems. This material (R&D cat. no. 405-ML) is biologically active, contains no HIS-tag, is expressed in the baculovirus Sf21 system, is also very pure (97 %), and can be purchased free of carrier-protein (BSA). Pooled sera from both immunisation groups reacted with the purchased mIL5 coated on ELISA plates, whereas sera from PBS immunised mice did not. This was shown when testing sera from bleed 3 obtained at day 75, 11 days after the 4<sup>th</sup> immunization, but also sera from bleed 1 and 2 reacts with the purchased mIL5 in a similar setup. In order to exclude signals from cross-reaction with the BSA carrier, the experiments were repeated for bleeds 1 and 2 using carrier-free versions of the purchased mIL5 material and BSA-free ELISA buffers, and still high anti-mIL5 responses are seen.

To further confirm that the induced antisera cross-react with native mIL5, a competitive ELISA was set up. This ELISA tests the ability of the different antisera to inhibit the interaction between soluble native murine IL5 and monoclonal antibodies TRFK4 or TRFK5, which are both neutralizing antibodies. Dilution series of antiserum pools were preincubated with soluble native mIL5 and the samples were added to ELISA plates



coated with catching antibody TRFK5. Bound murine IL5 (which was not absorbed by the antisera) was next visualised using layers of biotinylated TRFK4 and subsequently horseradish peroxidase labeled streptavidin. An anti-mIL5 positive and an anti-mIL5 negative antiserum from DNA vaccinated mice were included as controls. It was demonstrated that antisera from both HIS-mIL5wt and HIS-mIL5.1 immunized mice could inhibit the interaction between soluble mIL5 and TRFK4 or TRFK5.

10 Based on the above-referenced it is concluded that mIL5 specific autoantibodies are induced in mice immunized with either the HIS-mIL5wt or the HIS-mIL5.1 protein preparations (in 100% of the mice tested). In other words, B cell tolerance to mIL5 can be broken using recombinant HIS-tagged versions of both wild type and AutoVac murine IL5. A plausible explanation for the observation that B cell tolerance is broken to the wild type protein is that the HIS-tag in these mice functions as a "foreign" immunogenic T helper epitope. Another explanation could be that the administration of Complete Freund's Adjuvant breaks B cell tolerance to mIL5. These hypotheses can be tested using non-HIS tagged antigens and/or alternative, less strong adjuvants such as AdjuPhos.

25 Further characterization of the antibody responses in mice immunized with mIL5 AutoVac proteins:

ELISA experiments were set up in order to determine whether antibodies specific for the inserted T helper epitope could be detected in sera from mIL5 protein immunised mice. For each immunisation group, antisera (bleed 2) were pooled and tested for reactivity against synthetic P2 peptide which had been immobilised in AquaBind microtiter plates. Anti-HIS-mIL5.1 antiserum contained reactivity against the inserted P2 peptide, whereas neither anti-HIS-mIL5wt or anti-PBS/CFA reacted with the peptide.

It was also tested whether the the anti-HIS-mILwt and anti-HIS-mIL5.1 antisera contained reactivity against the 15-mer HIS-tag (UNIZYME HIS-tag, SEQ ID NO: 25) that is fused to the N-terminal of both the wild type and AutoVac mIL5 proteins. The peptide was synthesized and covalently immobilized in AquaBind microtiter plates, and pooled antisera from each immunization group (bleeds 1, 2 and 3) were tested for reactivity against the bound peptide. Antisera from all protein immunized mice reacted with the synthetic HIS-tag peptide.

It was also tested whether the anti-HIS-mIL5wt and anti-HIS-mIL5.1 antisera was reactive with components from the S2 Drosophila cells or culture medium. ELISA plates coated with BSA (a major medium component) or S2-background preparation (generated by subjecting culture supernatant from Her2 expressing Drosophila S2 cells to a purification scheme similar to that of the mIL5 purification procedure). The results of these analyses demonstrated that whereas the anti-BSA responses were very low, the reactions with the S2-background material were pronounced.

#### Eosinophil Counts in BALF:

To determine if the anti-IL5 antibodies in vaccinated mice could down-regulate the *in vivo* activity of IL5, we induced IL5-dependent eosinophilia in the lungs of the vaccinated mice. Eosinophils were induced by challenging sensitized mice with OVA intranasally. High numbers of eosinophils were induced in control OVA mice and mice vaccinated with UniHis-mIL5.1, but not in Uni-His-mIL5 or PBS vaccinated mice. The discrepancy of eosinophil numbers between control groups (OVA and PBS) and experimental groups (UniHis-mIL5 and UniHis-mIL5.1), and the positive results from the DNA vaccinated mice reported above, led us to believe that the groups may have

been switched. However, no attempts to demonstrate a switch supported this interpretation. The protein vaccinations are being repeated in an identical setup to clarify this controversy.

5

### Discussion:

The ability of both the UniHis-mIL5 and UniHis-mIL5.1 proteins to induce antibodies that cross-react with wildtype murine IL5 was clearly demonstrated. Whether the ability of the UniHis-mIL5 protein to bypass immunological tolerance is due to the UniHis-tag, or some other reason (e.g. CFA adjuvant) remains to be clarified. It was surprising to see that only the UniHis-mIL5 construct was able to down-regulate the endogenous *in vivo* activity of mIL5 in an eosinophilia model. This inability of antisera generated from UniHis-mIL5.1 protein vaccination to inhibit eosinophilia, and its ability to inhibit eosinophilia via DNA vaccinations suggests that a technical mistake may have occurred in this experiment. This is also supported by the unusual finding of PBS vaccination inhibiting eosinophilia. This most likely explanation is that these two groups (PBS and UniHis-mIL5.1) were switched.

### 25 LIST OF REFERENCES

Akutsu I. *et al.*, 1995, *Immunol. Lett.*, 45: 109-116.  
Alexander A.G. *et al.*, 1994, *Thorax*, 49(12): 1231-1233.

30 Azuma C. *et al.*, 1986, *Nucleic Acid Res.* 1986, 14(22): 9149-9158.

Barata L.T. *et al.*, 1998, *J. Allergy and Clin. Immunol.*, 101: 222-230.

35

Baumann M.A. *et al.*, 1997, *Methods*, 11: 88-97

Callard R.E. & Gearing A.J.H., 'IL-5', Cytokine Facts Book 1994, Academic Press.

Campbell H.D. et al., 1988, Eur. J. Biochem., 174: 345-352.

5

Chand N. et al., 1992, Eur. J. Immunol., 21: 121-123.

Coefrier E. et al., 1994, Br. J. Pharmacol., 113(3): 749-56.

10 Coffman R.L. et al., 1989, Science, 245: 308-310.

Corrigan C.J. & Kay A.B., 1996, Eur. Resp. J., 9, suppl. 22: 72s-78s.

15 Cousins D.J. et al., 1994, Am. J. Resp. Crit. Care. Med., 150: S50-S53.

Danzig M. et al., 1997, Allergy, 52(8): 787-794.

20 Dickason R.R. et al., 1994, Cytokine, 6(6): 647-656.

Dickason R.R. et al., 1996a, Nature, 379: 652-655.

Dickason R.R. et al., 1996b, J. Mol. Med., 74(9), 535-546

25

Egan R.W. et al., 1995, Int. Arch. Allergy Immunol., 107: 321-322.

Foster P.S. et al., 1996, J. Exp. Med., 183: 195-201.

30

Graber P. et al., 1993, Eur. J. Biochem., 212(3): 751-755.

Graber P. et al., 1995, J. Biol. Chem., 270(26): 15762-15769.

35 Hamelmann E. et al., 1997, Am. J. Crit. Care Med., 155(3): 819-825.

Huston M.M. et al., 1996, J. Immunol., 156(4): 1392-1401.

40 Karlen S. et al., 1998, Int. Rev. Immunol., 16(3-4): 227-247.

Kodama S. et al., 1993, Eur. J. Biochem., 211(3): 903-908.

Kopf M. et al., 1996, Immunity, 4: 15-24.

45

Kung T.T. et al., 1995, Am. J. Respir. Cell. Mol. Biol., 13: 360-365.

Lee N.A. et al., 1997a, J. Immunol., 158: 1332-1344.

50

Lee J.J. et al., 1997b, J. Exp. Med. 1997b, 185(12): 2143-2156.

Lopez A.F. et al., 1992, Immunology Today, 13: 495-500.

5

Mauser P.J. et al., 1993, Am. Rev. Respir. Dis., 148: 1623-1627.

Mauser P.J. et al., 1995, Am. J. Respir. Crit. Care Med., 10 152(2): 467-472.

Milburn M.V. et al., 1993, Nature, 363: 172-176.

Mori A. et al., 1997, J. Allergy Clin. Immunol., 100(6) Pt 2: 15 S56-64.

Moxham J. & Costello J.F., 'Respiratory diseases', chapt. 14, Textbook of Medicine, Churchill Livingstone 1990, Ed. Souhami R.L. and Moxham J.

20

Nagai H. et al., 1993a, Ann. N.Y. Acad.Sci., 91-96.

Nagai H. et al., 1993b, Life Sciences, 53: PL 243-247.

Ohashi Y. et al., 1998, Scand. J. Immunol, 47: 596-602.

Ortega D. & Busse W.W., 'Asthma: Pathogenesis and treatment', chapt. 28, Allergy, W.B. Saunders Company 1997, Ed. Kaplan A.P.

30

Proudfoot A.E. et al., 1990, Biochem J., 270(2): 357-361.

Proudfoot A.E. et al., 1996, J. Protein Chem., 15(5): 491-499.

Rose K. et al., 1992, Biochem J, 286(Pt 3): 825-828.

Sanderson C.J., 1992, Blood, 79(12): 3101-3109.

Sher A. et al., 1990, J. Immunol., 145: 3911-3916.

40

Takatsu K. et al., Interleukin-5, Growth Factors and Cytokines in Health and Disease 1997, vol.2A, JAI Press Inc., Ed. Le-roith D. & Bondy C.

Tanabe T. et al., 1987, J. Biol. Chem., 262: 16580-16584.

Tavernier J. et al., 1989, DNA, 8(7), 491-501.

Tominaga A. et al., 1990, J. Immunol., 144(4): 1345-1352.

50

Tominaga A. et al., 1991, J. Exp. Med., 173(2): 429-437.

- Underwood D.C. et al., 1996, Am. J. Resp. Crit. Care Med.,  
154: 850-857.
- 5 van Oosterhout A.J.M. et al., 1993, Am. Rev. Resp. Dis., 147:  
548-552.
- van Oosterhout A.J.M. et al., 1995, Am. J. Respir. Crit. Care  
Med., 151: 177-183.
- 10 Villinger F. et al., 1995, J. Immunol., 155: 3946-3954.
- Wang P. et al., 1998, J. Immunol., 160: 4427-4432.
- 15 Yamaguchi Y. et al., 1991, Blood, 78(10): 2542-2547.

1

## Amended claims

1. A method for *in vivo* down-regulation of interleukin 5 (IL5) activity in an animal, including a human being, the method comprising effecting presentation to the animal's immune system of an immunogenically effective amount of
- 5 at least one IL5 polypeptide autologous in the animal or a subsequence thereof which has been formulated so that immunization of the animal with the autologous IL5 polypeptide or subsequence thereof induces production by the animal of antibodies against the IL5 polypeptide, and/or
- 10 - at least one IL5 analogue wherein is introduced at least one modification in the amino acid sequence of the animal's autologous IL5 polypeptide which has as a result that immunization of the animal with the analogue induces production of antibodies in the animal against the animal's autologous IL5 polypeptide.
- 15 2. The method according to claim 1, wherein is presented an IL5 analogue with at least one modification of the IL5 amino acid sequence.
3. The method according to claim 2, wherein the modification has as a result that a substantial fraction of IL5 B-cell epitopes are preserved and that
- 20 - at least one foreign T helper lymphocyte epitope ( $T_H$  epitope) is introduced, and/or
- at least one first moiety is introduced which effects targeting of the modified molecule to an antigen presenting cell (APC) or a B-lymphocyte, and/or
- at least one second moiety is introduced which stimulates the immune system, and/or
- 25 - at least one third moiety is introduced which optimizes presentation of the modified IL5 polypeptide to the immune system.
4. The method according to claim 3, wherein the modification includes introduction as side groups, by covalent or non-covalent binding to suitable chemical groups in IL5 or a subsequence thereof, of the foreign  $T_H$  epitope and/or of the first and/or of the second and/or of the third moiety.
- 30 5. The method according to claim 3 or 4, wherein the modification includes amino acid substitution and/or deletion and/or insertion and/or addition.
- 35 6. The method according to claim 5, wherein the modification results in the provision of a fusion polypeptide.

AMENDED SHEET

CLAIM/23058PC01

2

7. The method according to claim 5 or 6, wherein introduction of the amino acid substitution and/or deletion and/or insertion and/or addition results in a substantial preservation of the overall tertiary structure of IL5.

5

8. The method according to any one of claims 2-7, wherein the modification includes duplication of at least one IL5 B-cell epitope and/or introduction of a hapten.

9. The method according to any one of claims 3-8, wherein the foreign T-cell epitope is immunodominant in the animal.

10. The method according to any one of claims 3-9, wherein the foreign T-cell epitope is promiscuous.

11. The method according to claim 10, wherein the at least one foreign T-cell epitope is selected from a natural promiscuous T-cell epitope and an artificial MHC-II binding peptide sequence.

12. The method according to claim 11, wherein the natural T-cell epitope is selected from a Tetanus toxoid epitope such as P2 or P30, a diphtheria toxoid epitope, an influenza virus hemagglutinin epitope, and a *P. falciparum* CS epitope.

13. The method according to any one of claims 3-12, wherein the first moiety is a substantially specific binding partner for a B-lymphocyte specific surface antigen or for an APC specific surface antigen, such as a hapten or a carbohydrate for which there is a receptor on the B-lymphocyte or the APC.

14. The method according to any one of claims 3-13, wherein the second moiety is selected from a cytokine, a hormone, and a heat-shock protein.

30

15. The method according to claim 6, wherein the cytokine is selected from, or is an effective part of, interferon  $\gamma$  (IFN- $\gamma$ ), Flt3L, interleukin 1 (IL-1), interleukin 2 (IL-2), interleukin 4 (IL-4), interleukin 6 (IL-6), interleukin 12 (IL-12), interleukin 13 (IL-13), interleukin 15 (IL-15), and granulocyte-macrophage colony stimulating factor (GM-CSF), and the heat-shock protein is selected from, or is an effective part of any of, HSP70, HSP90, HSC70, GRP94, and calreticulin (CRT).

AMENDED SHEET

CLAIM/23058PC01



3

16. The method according to any one of claims 3-15, wherein the third moiety is of lipid nature, such as a palmitoyl group, a myristyl group, a farnesyl group, a geranyl-geranyl group, a GPI-anchor, and an N-acyl diglyceride group.

5 17. The method according to any one of the preceding claims, wherein the IL5 polypeptide has been modified in at least one of loops 1-3 or in the amino acid residues C-terminal to helix D, said loops and said helix D corresponding to those shown in Fig. 3 for human and murine IL5.

10 18. The method according to claim 17, wherein the IL5 polypeptide is a human IL5 polypeptide.

15 19. The method according to claim 18, wherein the human IL5 polypeptide has been modified by substituting at least one amino acid sequence in SEQ ID NO: 1 with at least one amino acid sequence of equal or different length thereby giving rise to a foreign T<sub>H</sub> epitope, wherein substituted amino acid residues are selected from the group consisting of residues 87-90, residues 88-91, residues 32-43, residues 33-43, residues 59-64, residues 86-91, and residues 110-113.

20 20. The method according to any one of the preceding claims, wherein presentation to the immune system is effected by having at least two copies of the IL5 polypeptide, the subsequence thereof or the modified IL5 polypeptide covalently or non-covalently linked to a carrier molecule capable of effecting presentation of multiple copies of antigenic determinants.

25

21. The method according to any the preceding claims, wherein the IL5 polypeptide, the subsequence thereof, or the modified IL5 polypeptide has been formulated with an adjuvant which facilitates breaking of autotolerance to autoantigens.

30 22. The method according to claim 21, wherein the adjuvant is selected from the group consisting of an immune targeting adjuvant; an immune modulating adjuvant such as a toxin, a cytokine and a mycobacterial derivative; an oil formulation; a polymer; a micelle forming adjuvant; a saponin; an immunostimulating complex matrix (an ISCOM matrix); a particle; DDA; aluminium adjuvants; DNA adjuvants;  $\gamma$ -inulin; and an encapsulating  
35 adjuvant.

CLAIM/23058PC01

AMENDED SHEET

4

23. The method according to any one of the preceding claims, wherein an effective amount of the IL5 polypeptide or the IL5 analogue is administered to the animal via a route selected from the parenteral route such as the intradermal, the subdermal, the intracutaneous, the subcutaneous, and the intramuscular routes; the peritoneal route; the oral route; the buccal route; the sublingual route; the epidural route; the spinal route; the anal route; and the intracranial route.

24. The method according to claim 23, wherein the effective amount is between 0.5 µg and 2,000 µg of the IL5 polypeptide, the subsequence thereof or the analogue thereof.

10

25. The method according to claim 23 or 24, which includes at least one administration of the IL5 polypeptide or analogue per year, such as at least 2, at least 3, at least 4, at least 6, and at least 12 administrations per year.

15 26. The method according to any one of claims 23-25, wherein the IL5 polypeptide or analogue is contained in a virtual lymph node (VLN) device.

27. The method according to any one of claims 1-20, wherein presentation of modified IL5 to the immune system is effected by introducing nucleic acid(s) encoding the modified IL5 into the animal's cells and thereby obtaining *in vivo* expression by the cells of the nucleic acid(s) introduced.

28. The method according to claim 27, wherein the nucleic acid(s) introduced is/are selected from naked DNA, DNA formulated with charged or uncharged lipids, DNA formulated in liposomes, DNA included in a viral vector, DNA formulated with a transfection-facilitating protein or polypeptide, DNA formulated with a targeting protein or polypeptide, DNA formulated with Calcium precipitating agents, DNA coupled to an inert carrier molecule, DNA encapsulated in chitin or chitosan, and DNA formulated with an adjuvant such as the adjuvants defined in claim 22.

30

29. The method according to claim 27 or 28, wherein the nucleic acids are administered intraarterially, intravenously, or by the routes defined in claim 23.

30. The method according to claim 28 or 29, wherein the nucleic acid(s) is/are contained in a VLN device.

AMENDED SHEET

CLAIM/23058PC01

5

31. The method according to any one of claims 28-30, which includes at least one administration of the nucleic acids per year, such as at least 2, at least 3, at least 4, at least 6, and at least 12 administrations per year

5 32. A method for treating and/or preventing and/or ameliorating asthma or other chronic allergic conditions characterized by eosinophilia, the method comprising down-regulating IL5 activity according to the method of any one of claims 1-31 to such an extent that the number of eosinophil cells, either systemically or locally at the disease focus, is significantly reduced, such as a reduction of at least 20%.

10

33. An IL5 analogue which is derived from an animal IL5 polypeptide wherein is introduced a modification which has as a result that immunization of the animal with the analogue induces production of antibodies against the IL5 polypeptide, and wherein the modification involves amino acid substitution and/or insertion and/or deletion to any one of  
15 loops 1-3 or C-terminally to helix D in IL5.

34. An IL5 analogue according to claim 33, wherein the modification is as defined in any one of claims 2-20.

20 35. An immunogenic composition comprising an immunogenically effective amount of an IL5 polypeptide autologous in an animal, said IL5 polypeptide being formulated together with an immunologically acceptable adjuvant so as to break the animal's autotolerance towards the IL5 polypeptide, the composition further comprising a pharmaceutically and immunologically acceptable carrier and/or vehicle.

25

36. An immunogenic composition comprising an immunogenically effective amount of an IL5 analogue according to claim 33 or 34, the composition further comprising a pharmaceutically and immunologically acceptable carrier and/or vehicle and optionally an adjuvant.

30

37. An immunogenic composition according to Claim 35 or 36, wherein the adjuvant is selected from the group consisting of the adjuvants of claim 22.

38. A nucleic acid fragment which encodes an IL5 analogue according to claim 33 or 34.

35

39. A vector carrying the nucleic acid fragment according to claim 38.

AMENDED SHEET

CLAIM/23058PC01

6

40. The vector according to claim 39 which is capable of autonomous replication.

41. The vector according to claim 39 or 40 which is selected from the group consisting of a plasmid, a phage, a cosmid, a mini-chromosome, and a virus.

5

42. The vector according to any one of claims 39-41, comprising, in the 5'→3' direction and in operable linkage, a promoter for driving expression of the nucleic acid fragment according to claim 38, optionally a nucleic acid sequence encoding a leader peptide enabling secretion of or integration into the membrane of the polypeptide fragment, the nucleic acid fragment according to claim 38, and optionally a terminator.

43. The vector according to any one of claims 39-42 which, when introduced into a host cell, is integrated in the host cell genome.

44. The vector according to any one of claims 39-42 which, when introduced into a host cell, is not capable of being integrated in the host cell genome.

45. The vector according to any one of claims 39-44, wherein the promoter drives expression in a eukaryotic cell and/or in a prokaryotic cell.

20

46. A transformed cell carrying the vector of any one of claims 39-45.

47. The transformed cell according to claim 46 which is capable of replicating the nucleic acid fragment according to claim 38.

25

48. The transformed cell according to claim 47, which is a microorganism selected from a bacterium, a yeast, a protozoan, or a cell derived from a multicellular organism selected from a fungus, an insect cell such as an S<sub>2</sub> or an SF cell, a plant cell, and a mammalian cell.

30

49. The transformed cell according to claim 48 which is a bacterium of the genus *Escherichia*, *Bacillus*, *Salmonella*, or *Mycobacterium*.

50. The transformed cell according to claim 52, which is selected from the group consisting of an *E. coli* cell, and a non-pathogenic *Mycobacterium* cell such as *M. bovis* BCG.

AMENDED SHEET

CLAIM/23056PC01

7

51. The transformed cell according to any one of claims 46-50, which expresses the nucleic acid fragment according to claim 38.

52. The transformed cell according to claim 55, which secretes or carries on its surface,  
5 the IL5 analogue according to claim 33 or 34.

53. The method according to any one of claims 1-20, wherein presentation to the immune system is effected by administering a non-pathogenic microorganism or virus which is carrying a nucleic acid fragment which encodes and expresses the IL5 polypeptide or  
10 analogue.

54. The method according to claim 53, wherein the virus is a non-virulent pox virus such as a vaccinia virus.

15 55. The method according to claim 54, wherein the microorganism is a bacterium, such as a bacterium defined in claim 49 or 50.

56. The method according to any one of claims 53-55, wherein the non-pathogenic microorganism or virus is administered one single time to the animal.  
20

57. A composition for inducing production of antibodies against IL5, the composition comprising

- a nucleic acid fragment according to claim 38 or a vector according to any one of claims 39-45, and
- 25 - a pharmaceutically and immunologically acceptable carrier and/or vehicle and/or adjuvant.

58. The composition according to claim 57, wherein the nucleic acid fragment is formulated according to claim 28 or 30.  
30

59. A stable cell line which carries the vector according to any one of claims 39-45 and which expresses the nucleic acid fragment according to claim 38, and which optionally secretes or carries the IL5 analogue according to claim 33 or 34 on its surface.

35 60. A method for the preparation of the cell according to any one of claims 46-52, the method comprising transforming a host cell with the nucleic acid fragment according to claim 38 or with the vector according to any one of claims 39-45.

CLAIM/23058PC01

AMENDED SHEET

61. A method for the identification of a modified IL5 polypeptide which is capable of inducing antibodies against unmodified IL5 in an animal species where the unmodified IL5 polypeptide is a self-protein, the method comprising

5

- preparing, by means of peptide synthesis or genetic engineering techniques, a set of mutually distinct modified IL5 polypeptides wherein amino acids have been added to, inserted in, deleted from, or substituted into the amino acid sequence of an IL5 polypeptide of the animal species thereby giving rise to amino acid sequences in the set which comprise T-cell epitopes which are foreign to the animal species, or preparing a set of nucleic acid fragments encoding the set of mutually distinct modified IL5 polypeptides,
- testing members of the set for their ability to induce production of antibodies by the animal species against the unmodified IL5, and
- 15 - identifying and optionally isolating the member(s) of the set which significantly induces antibody production against unmodified IL5 in the animal species, or identifying and optionally isolating the polypeptide expression products encoded by members of the set of nucleic acid fragments which significantly induces antibody production against unmodified IL5 in the animal species.

20

62. A method for the preparation of an immunogenic composition comprising at least one modified IL5 polypeptide which is capable of inducing antibodies against unmodified IL5 in an animal species where the unmodified IL5 polypeptide is a self-protein, the method comprising

25

- preparing, by means of peptide synthesis or genetic engineering techniques, a set of mutually distinct modified IL5 polypeptides wherein amino acids have been added to, inserted in, deleted from, or substituted into the amino acid sequence of an IL5 polypeptide of the animal species thereby giving rise to amino acid sequences in the set comprising T-cell epitopes which are foreign to the animal,
- 30 - testing members of the set for their ability to induce production of antibodies by the animal species against the unmodified IL5, and
- admixing the member(s) of the set which significantly induces production of antibodies in the animal species which are reactive with IL5 with a pharmaceutically and immunologically acceptable carrier and/or vehicle, optionally in combination with at least one pharmaceutically and immunologically acceptable adjuvant.

35

AMENDED SHEET

CLAIM/23058PC01

Friday 27 of Apr 2001, PV&amp;P 33639600

-&gt;+49 89 2399 4465

Page 24 of 28

9

63. The method according to claim 61 or 62, wherein preparation of the members of the set comprises preparation of mutually distinct nucleic acid sequences, each sequence being a nucleic acid sequence according to claim 38, insertion of the nucleic acid sequences into appropriate expression vectors, transformation of suitable host cells with the vectors, and expression of the nucleic acid sequences, optionally followed by isolation of the expression products.
64. The method according to claim 63, wherein the preparation of the nucleic acid sequences and/or the vectors is achieved by the aid of a molecular amplification technique such as PCR, or by the aid of nucleic acid synthesis.
65. Use of IL5 or a subsequence thereof for the preparation of an immunogenic composition comprising an adjuvant for down-regulating IL5 activity in an animal.
66. Use of IL5 or a subsequence thereof for the preparation of an immunogenic composition comprising an adjuvant for the treatment, prophylaxis or amelioration of asthma or other chronic allergic conditions.
67. Use of an IL5 analogue for the preparation of an immunogenic composition optionally comprising an adjuvant for down-regulating IL5 activity in an animal.
68. Use of an IL5 analogue for the preparation of an immunogenic composition optionally comprising an adjuvant for the treatment, prophylaxis or amelioration of asthma or other chronic allergic conditions.

CLAIM/23058PC01

AMENDED SHEET

1/7

		20
Ile-Pro-Thr-Glu-Ile-Pro-Thr-Ser-Ala-Leu-Val-Lys-Glu-Thr-Leu-Ala-Leu-Leu-Ser-Thr-		
* * Met	Met Thr Val	Ala
		40
His-Arg-Thr-Leu-Leu-Ile-Ala-Asn-Glu-Thr-Leu-Arg-Ile-Pro-Val-Pro-Val-His-Lys-Asn-		
Ala	Thr Ser Met Leu Thr	
		60
His-Gln-Leu-Cys-Thr-Glu-Glu-Ile-Phe-Gln-Gly-Ile-Gly-Thr-Leu-Glu-Ser-Gln-Thr-Val-		
Ile Gly	Leu Asp Ile Lys Asn	
		80
Gln-Gly-Gly-Thr-Val-Glu-Arg-Leu-Phe-Lys-Asn-Leu-Ser-Leu-Ile-Lys-Lys-Tyr-Ile-Asp-		
Arg	Met Gln	
		100
Gly-Gln-Lys-Lys-Lys-Cys-Gly-Glu-Glu-Arg-Arg-Val-Asn-Gln-Phe-Leu-Asp-Tyr-Leu-		
Arg	Glu Thr Arg	
		110
Gln-Glu-Phe-Leu-Gly-Val-Met-Asn-Thr-Glu-Trp-Ile-Ile-Glu-Ser		
	Ser Ala Met Gly	
		115

Fig. 1



2/7




Fig. 2A

256777-000000

09/980916

3/7




Fig. 2B

0980916-021902

4/7




Fig. 2C

205 T20 " 9 T608650

5/7




Fig. 3

6/7




Fig. 4




Fig. 5

## BIRCH, STEWART, KOLASCH &amp; BIRCH, LLP

P.O. Box 747 • Falls Church, Virginia 22040-0747  
 Telephone: (703) 205-8000 • Facsimile (703) 205-8050

PLEASE NOTE:  
 YOU MUST  
 COMPLETE THE  
 FOLLOWING

## COMBINED DECLARATION AND POWER OF ATTORNEY FOR PATENT AND DESIGN APPLICATIONS

As a below named inventor, I hereby declare that: my residence, post office address and citizenship are as stated next to my name; that I verily believe that I am the original, first and sole inventor (if only one inventor is named below) or an original, first and joint inventor (if plural inventors are named below) of the subject matter which is claimed and for which a patent is sought on the invention entitled:

Insert Title:

### METHOD FOR DOWN-REGULATING IL5 ACTIVITY

Fill in Appropriate Information - For Use Without Specification Attached: the specification of which is attached hereto. If not attached hereto, the specification was filed on October 23, 2001 as United States Application Number \_\_\_\_\_; and amended on \_\_\_\_\_ (if applicable) and/or the specification was filed on \_\_\_\_\_ as PCT International Application Number \_\_\_\_\_; and was amended under PCT Article 19 on \_\_\_\_\_ (if applicable)

I hereby state that I have reviewed and understand the contents of the above-identified specification, including the claims, as amended by any amendment referred to above.

I acknowledge the duty to disclose information which is material to patentability as defined in Title 37, Code of Federal Regulations, §1.56.

I do not know and do not believe the same was ever known or used in the United States of America before my or our invention thereof, or patented or described in any printed publication in any country before my or our invention thereof or more than one year prior to this application, that the same was not in public use or on sale in the United States of America more than one year prior to this application, that the invention has not been patented or made the subject of an inventor's certificate issued before the date of this application in any country foreign to the United States of America on an application filed by me or my legal representative or assigns more than twelve months (six months for designs) prior to this application, and that no application for patent or inventor's certificate on this invention has been filed in any country foreign to the United States of America prior to this application by me or my legal representatives or assigns, except as follows.

I hereby claim foreign priority benefits under Title 35, United States Code, §119(a)-(d) of any foreign application(s) for patent or inventor's certificate listed below and have also identified below any foreign application for patent or inventor's certificate having a filing date before that of the application on which priority is claimed:

### Prior Foreign Application(s)

### Priority Claimed

Insert Priority Information: (if appropriate)

PA 1999 00552 (Number)	Denmark (Country)	April 23, 1999 (Month/Day/Year Filed)	<input checked="" type="checkbox"/> Yes	<input type="checkbox"/> No
_____ (Number)	_____ (Country)	_____ (Month/Day/Year Filed)	<input type="checkbox"/> Yes	<input type="checkbox"/> No
_____ (Number)	_____ (Country)	_____ (Month/Day/Year Filed)	<input type="checkbox"/> Yes	<input type="checkbox"/> No
_____ (Number)	_____ (Country)	_____ (Month/Day/Year Filed)	<input type="checkbox"/> Yes	<input type="checkbox"/> No

I hereby claim the benefit under Title 35, United States Code, §119(e) of any United States provisional applications(s) listed below.

Insert Provisional Application(s): (if any)

60/132,811 (Application Number)	May 6, 1999 (Filing Date)
_____ (Application Number)	_____ (Filing Date)

All Foreign Applications, if any, for any Patent or Inventor's Certificate Filed More than 12 Months (6 Months for Designs) Prior to the Filing Date of This Application:

Country	Application Number	Date of Filing (Month/Day/Year)
---------	--------------------	---------------------------------

Insert Requested Information: (if appropriate)

_____	_____	_____
_____	_____	_____

I hereby claim the benefit under Title 35, United States Code, §120 of any United States and/or PCT application(s) listed below and, insofar as the subject matter of each of the claims of this application is not disclosed in the prior United States and/or PCT application in the manner provided by the first paragraph of Title 35, United States Code, §112, I acknowledge the duty to disclose information which is material to the patentability as defined in Title 37, Code of Federal Regulations, §1.56 which became available between the filing date of the prior application and the national or PCT international filing date of this application

Insert Prior U.S. Application(s): (if any)

_____ (Application Number)	_____ (Filing Date)	_____ (Status - patented, pending, abandoned)
-------------------------------	------------------------	--

I hereby appoint the practitioners at **CUSTOMER NO. 2292** as my attorneys or agents to prosecute this application and/or an international application based on this application and to transact all business in the United States Patent and Trademark Office connected therewith and in connection with the resulting patent based on instructions received from the entity who first sent the application papers to the practitioners, unless the inventor(s) or assignee provides said practitioners with a written notice to the contrary:

Send Correspondence to:

**BIRCH, STEWART, KOLASCH & BIRCH, LLP** or **CUSTOMER NO. 2292**

P.O. Box 747 • Falls Church, Virginia 22040-0747

Telephone: (703) 205-8000 • Facsimile: (703) 205-8050

PLEASE NOTE:  
YOU MUST  
COMPLETE  
THE  
FOLLOWING:

Full Name of First  
or Sole Inventor  
Insert Name of  
Inventor  
Insert Date This  
Document is Signed

Insert Residence  
Insert Citizenship

Insert Post Office  
Address

Full Name of Second  
Inventor, if any:  
see above

Full Name of Third  
Inventor, if any:  
see above

Full Name of Fourth  
Inventor, if any:  
see above

Full Name of Fifth  
Inventor, if any:  
see above

Full Name of Sixth  
Inventor, if any:  
see above

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patent issued thereon

GIVEN NAME/FAMILY NAME	INVENTOR'S SIGNATURE	DATE*
Steen KLYSNER	<i>Steen Kly</i>	3/12/01
Residence (City, State & Country)	CITIZENSHIP	
Hørsholm, Denmark <i>DKX</i>	Danish	
MAILING ADDRESS (Complete Street Address including City, State & Country)		
C/O Kogle Allé 6, DK-2970 Hørsholm Denmark		
GIVEN NAME/FAMILY NAME	INVENTOR'S SIGNATURE	DATE*
Residence (City, State & Country)	CITIZENSHIP	
MAILING ADDRESS (Complete Street Address including City, State & Country)		
GIVEN NAME/FAMILY NAME	INVENTOR'S SIGNATURE	DATE*
Residence (City, State & Country)	CITIZENSHIP	
MAILING ADDRESS (Complete Street Address including City, State & Country)		
GIVEN NAME/FAMILY NAME	INVENTOR'S SIGNATURE	DATE*
Residence (City, State & Country)	CITIZENSHIP	
MAILING ADDRESS (Complete Street Address including City, State & Country)		
GIVEN NAME/FAMILY NAME	INVENTOR'S SIGNATURE	DATE*
Residence (City, State & Country)	CITIZENSHIP	
MAILING ADDRESS (Complete Street Address including City, State & Country)		
GIVEN NAME/FAMILY NAME	INVENTOR'S SIGNATURE	DATE*
Residence (City, State & Country)	CITIZENSHIP	
MAILING ADDRESS (Complete Street Address including City, State & Country)		



<210> 2

[illegible]

<211> 126  
 <212> PRT  
 <213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:Human IL5  
 modified by substitution with tetanus toxoid P2  
 epitope

<220>

<221> MUTAGEN

<222> (32)..(46)

<223> Tetanus toxoid P2 epitope (SEQ ID NO: 23)

<220>

<221> SIMILAR

<222> (1)..(86)

<223> Identical to residues 1-86 in SEQ ID NO: 1

<220>

<221> SIMILAR

<222> (102)..(126)

<223> Identical to residues 91-115 in SEQ ID NO: 1

<400> 2

Ile Pro Thr Glu Ile Pro Thr Ser Ala Leu Val Lys Glu Thr Leu Ala  
 1 5 10 15

Leu Leu Ser Thr His Arg Thr Leu Leu Ile Ala Asn Glu Thr Leu Arg  
 20 25 30

Ile Pro Val Pro Val His Lys Asn His Gln Leu Cys Thr Glu Glu Ile  
 35 40 45

Phe Gln Gly Ile Gly Thr Leu Glu Ser Gln Thr Val Gln Gly Gly Thr  
 50 55 60

Val Glu Arg Leu Phe Lys Asn Leu Ser Leu Ile Lys Lys Tyr Ile Asp  
 65 70 75 80

Gly Gln Lys Lys Lys Cys Gln Tyr Ile Lys Ala Asn Ser Lys Phe Ile  
 85 90 95

Gly Ile Thr Glu Leu Arg Arg Val Asn Gln Phe Leu Asp Tyr Leu Gln  
 100 105 110

Glu Phe Leu Gly Val Met Asn Thr Glu Trp Ile Ile Glu Ser  
 115 120 125

<210> 3

<211> 118

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:Human IL5  
 modified by substitution with tetanus toxoid P2  
 epitope

09980916-024900

<220>  
 <221> MUTAGEN  
 <222> (32)..(46)  
 <223> Tetanus toxoid P2 epitope (SEQ ID NO: 23)

<220>  
 <221> SIMILAR  
 <222> (1)..(31)  
 <223> Identical to residues 1-31 in SEQ ID NO: 1

<220>  
 <221> SIMILAR  
 <222> (47)..(118)  
 <223> Identical to residues 44-115 in SEQ ID NO: 1

<400> 3  
 Ile Pro Thr Glu Ile Pro Thr Ser Ala Leu Val Lys Glu Thr Leu Ala  
           1                  5                  10                  15  
 Leu Leu Ser Thr His Arg Thr Leu Leu Ile Ala Asn Glu Thr Leu Gln  
                   20                  25                  30  
 Tyr Ile Lys Ala Asn Ser Lys Phe Ile Gly Ile Thr Glu Leu Cys Thr  
           35                  40                  45  
 Glu Glu Ile Phe Gln Gly Ile Gly Thr Leu Glu Ser Gln Thr Val Gln  
           50                  55                  60  
 Gly Gly Thr Val Glu Arg Leu Phe Lys Asn Leu Ser Leu Ile Lys Lys  
           65                  70                  75                  80  
 Tyr Ile Asp Gly Gln Lys Lys Lys Cys Gly Glu Glu Arg Arg Arg Val  
                   85                  90                  95  
 Asn Gln Phe Leu Asp Tyr Leu Gln Glu Phe Leu Gly Val Met Asn Thr  
                   100                  105                  110  
 Glu Trp Ile Ile Glu Ser  
           115

<210> 4  
 <211> 124  
 <212> PRT  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: Human IL5  
           modified by substitution with tetanus toxoid P2  
           epitope

<220>  
 <221> MUTAGEN  
 <222> (59)..(73)  
 <223> Tetanus toxoid P2 epitope (SEQ ID NO: 23)

<220>  
 <221> SIMILAR  
 <222> (1)..(58)  
 <223> Identical to residues 1-58 in SEQ ID NO: 1

0065058 WO 00/65058

&lt;220&gt;

&lt;221&gt; SIMILAR

&lt;222&gt; (74)..(124)

&lt;223&gt; Identical to residues 65-115 in SEQ ID NO: 1

&lt;400&gt; 4

Ile	Pro	Thr	Glu	Ile	Pro	Thr	Ser	Ala	Leu	Val	Lys	Glu	Thr	Leu	Ala
1				5					10					15	

Leu	Leu	Ser	Thr	His	Arg	Thr	Leu	Leu	Ile	Ala	Asn	Glu	Thr	Leu	Arg
			20					25					30		

Ile	Pro	Val	Pro	Val	His	Lys	Asn	His	Gln	Leu	Cys	Thr	Glu	Glu	Ile
		35					40					45			

Phe	Gln	Gly	Ile	Gly	Thr	Leu	Glu	Ser	Gln	Gln	Tyr	Ile	Lys	Ala	Asn
	50					55					60				

Ser	Lys	Phe	Ile	Gly	Ile	Thr	Glu	Leu	Val	Glu	Arg	Leu	Phe	Lys	Asn
65					70					75					80

Leu	Ser	Leu	Ile	Lys	Lys	Tyr	Ile	Asp	Gly	Gln	Lys	Lys	Lys	Cys	Gly
				85					90					95	

Glu	Glu	Arg	Arg	Arg	Val	Asn	Gln	Phe	Leu	Asp	Tyr	Leu	Gln	Glu	Phe
			100					105					110		

Leu	Gly	Val	Met	Asn	Thr	Glu	Trp	Ile	Ile	Glu	Ser
		115					120				

&lt;210&gt; 5

&lt;211&gt; 124

&lt;212&gt; PRT

&lt;213&gt; Artificial Sequence

&lt;220&gt;

<223> Description of Artificial Sequence: Human IL5  
modified by substitution with tetanus toxoid P2  
epitope

&lt;220&gt;

&lt;221&gt; MUTAGEN

&lt;222&gt; (86)..(100)

&lt;223&gt; Tetanus toxoid P2 epitope (SEQ ID NO: 23)

&lt;220&gt;

&lt;221&gt; SIMILAR

&lt;222&gt; (1)..(85)

&lt;223&gt; Identical to residues 1-85 in SEQ ID NO: 1

&lt;220&gt;

&lt;221&gt; SIMILAR

&lt;222&gt; (101)..(124)

&lt;223&gt; Identical to residues 90-115 in SEQ ID NO: 1

&lt;400&gt; 5

Ile	Pro	Thr	Glu	Ile	Pro	Thr	Ser	Ala	Leu	Val	Lys	Glu	Thr	Leu	Ala
1				5					10					15	

20051220 9T600650

```

Leu Leu Ser Thr His Arg Thr Leu Leu Ile Ala Asn Glu Thr Leu Arg
      20                      25                      30

Ile Pro Val Pro Val His Lys Asn His Gln Leu Cys Thr Glu Glu Ile
      35                      40                      45

Phe Gln Gly Ile Gly Thr Leu Glu Ser Gln Thr Val Gln Gly Gly Thr
      50                      55                      60

Val Glu Arg Leu Phe Lys Asn Leu Ser Leu Ile Lys Lys Tyr Ile Asp
      65                      70                      75                      80

Gly Gln Lys Lys Lys Gln Tyr Ile Lys Ala Asn Ser Lys Phe Ile Gly
      85                      90                      95

Ile Thr Glu Leu Arg Val Asn Gln Phe Leu Asp Tyr Leu Gln Glu Phe
      100                     105                     110

Leu Gly Val Met Asn Thr Glu Trp Ile Ile Glu Ser
      115                     120

```

```

<210> 6
<211> 126
<212> PRT
<213> Artificial Sequence

```

```

<220>
<223> Description of Artificial Sequence:Human IL5
      modified by substitution with tetanus toxoid P2
      epitope

```

```

<220>
<221> MUTAGEN
<222> (110)..(124)
<223> Tetanus toxoid P2 epitope (SEQ ID NO: 23)

```

```

<220>
<221> SIMILAR
<222> (1)..(109)
<223> Identical to residues 1-109 in SEQ ID NO: 1

```

```

<220>
<221> SIMILAR
<222> (125)..(126)
<223> Identical to residues 114-115 in SEQ ID NO: 1

```

```

<400> 6
Ile Pro Thr Glu Ile Pro Thr Ser Ala Leu Val Lys Glu Thr Leu Ala
  1                      5                      10                      15

Leu Leu Ser Thr His Arg Thr Leu Leu Ile Ala Asn Glu Thr Leu Arg
      20                      25                      30

Ile Pro Val Pro Val His Lys Asn His Gln Leu Cys Thr Glu Glu Ile
      35                      40                      45

Phe Gln Gly Ile Gly Thr Leu Glu Ser Gln Thr Val Gln Gly Gly Thr
      50                      55                      60

```

05580516 034190

Val Glu Arg Leu Phe Lys Asn Leu Ser Leu Ile Lys Lys Tyr Ile Asp  
65 70 75 80

Gly Gln Lys Lys Lys Cys Gly Glu Glu Arg Arg Arg Val Asn Gln Phe  
85 90 95

Leu Asp Tyr Leu Gln Glu Phe Leu Gly Val Met Asn Thr Gln Tyr Ile  
100 105 110

Lys Ala Asn Ser Lys Phe Ile Gly Ile Thr Glu Leu Glu Ser  
115 120 125

<210> 7  
<211> 132  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Human IL5  
modified by substitution with tetanus toxoid P30  
epitope

<220>  
<221> MUTAGEN  
<222> (87)..(107)  
<223> Tetanus toxoid P30 epitope (SEQ ID NO: 24)

<220>  
<221> SIMILAR  
<222> (1)..(86)  
<223> Identical to residues 1-86 in SEQ ID NO: 1

<220>  
<221> SIMILAR  
<222> (108)..(132)  
<223> Identical to residues 91-115 in SEQ ID NO: 1

<400> 7  
Ile Pro Thr Glu Ile Pro Thr Ser Ala Leu Val Lys Glu Thr Leu Ala  
1 5 10 15

Leu Leu Ser Thr His Arg Thr Leu Leu Ile Ala Asn Glu Thr Leu Arg  
20 25 30

Ile Pro Val Pro Val His Lys Asn His Gln Leu Cys Thr Glu Glu Ile  
35 40 45

Phe Gln Gly Ile Gly Thr Leu Glu Ser Gln Thr Val Gln Gly Gly Thr  
50 55 60

Val Glu Arg Leu Phe Lys Asn Leu Ser Leu Ile Lys Lys Tyr Ile Asp  
65 70 75 80

Gly Gln Lys Lys Lys Cys Phe Asn Asn Phe Thr Val Ser Phe Trp Leu  
85 90 95

206722 97608650

```
<210> 8
<211> 124
<212> PRT
<213> Artificial Sequence
```

<220>  
<223> Description of Artificial Sequence:Human IL5  
modified by substitution with tetanus toxoid P30  
epitope

```
<220>
<221> MUTAGEN
<222> (32)..(52)
<223> Tetanus toxoid P30 epitope (SEQ ID NO: 24)
```

```
<220>  
<221> SIMILAR  
<222> (1)..(31)  
<223> Identical to residues 1-31 in SEQ ID NO: 1
```

```
<220>  
<221> SIMILAR  
<222> (53)..(124)  
<223> Identical to residues 44-115 in SEQ ID NO: 1
```

```

<400> 8
Ile Pro Thr Glu Ile Pro Thr Ser Ala Leu Val Lys Glu Thr Leu Ala
  1          5          10          15

Leu Leu Ser Thr His Arg Thr Leu Leu Ile Ala Asn Glu Thr Leu Phe
          20          25          30

Asn Asn Phe Thr Val Ser Phe Trp Leu Arg Val Pro Lys Val Ser Ala
          35          40          45

Ser His Leu Glu Cys Thr Glu Glu Ile Phe Gln Gly Ile Gly Thr Leu
  50          55          60

Glu Ser Gln Thr Val Gln Gly Gly Thr Val Glu Arg Leu Phe Lys Asn
  65          70          75          80

Leu Ser Leu Ile Lys Lys Tyr Ile Asp Gly Gln Lys Lys Lys Cys Gly
          85          90          95

Glu Glu Arg Arg Arg Val Asn Gln Phe Leu Asp Tyr Leu Gln Glu Phe
          100          105          110

Leu Gly Val Met Asn Thr Glu Trp Ile Ile Glu Ser
  115          120

```

<210> 9  
 <211> 130  
 <212> PRT  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: Human IL5  
 modified by substitution with tetanus toxoid P30  
 epitope

<220>  
 <221> MUTAGEN  
 <222> (59)..(79)  
 <223> Tetanus toxoid P30 epitope (SEQ ID NO: 24)

<220>  
 <221> SIMILAR  
 <222> (1)..(58)  
 <223> Identical to residues 1-58 in SEQ ID NO: 1

<220>  
 <221> SIMILAR  
 <222> (80)..(130)  
 <223> Identical to residues 65-115 in SEQ ID NO: 1

<400> 9  
 Ile Pro Thr Glu Ile Pro Thr Ser Ala Leu Val Lys Glu Thr Leu Ala  
           1                  5                  10                  15  
 Leu Leu Ser Thr His Arg Thr Leu Leu Ile Ala Asn Glu Thr Leu Arg  
                   20                  25                  30  
 Ile Pro Val Pro Val His Lys Asn His Gln Leu Cys Thr Glu Glu Ile  
           35                  40                  45  
 Phe Gln Gly Ile Gly Thr Leu Glu Ser Gln Phe Asn Asn Phe Thr Val  
           50                  55                  60  
 Ser Phe Trp Leu Arg Val Pro Lys Val Ser Ala Ser His Leu Glu Val  
           65                  70                  75                  80  
 Glu Arg Leu Phe Lys Asn Leu Ser Leu Ile Lys Lys Tyr Ile Asp Gly  
                   85                  90                  95  
 Gln Lys Lys Lys Cys Gly Glu Glu Arg Arg Arg Val Asn Gln Phe Leu  
           100                  105                  110  
 Asp Tyr Leu Gln Glu Phe Leu Gly Val Met Asn Thr Glu Trp Ile Ile  
           115                  120                  125  
 Glu Ser  
           130

<210> 10  
 <211> 132  
 <212> PRT

09980916 "021902"



<220>

 $\langle 220 \rangle$ 

<222> (110) .. (130)

<220>

<222> (1) .. (129)

 $\langle 220 \rangle$ 

<222> (131) .. (132)

<400> 10

<210> 11

<212> PRT

**<220>**

<223> Description of Artificial Sequence: Human IL5 modified by substitution with tetanus toxoid P2 and P30 epitopes

<220>  
 <221> MUTAGEN  
 <222> (86)..(100)  
 <223> Tetanus toxoid P2 epitope (SEQ ID NO: 23)

<220>  
 <221> MUTAGEN  
 <222> (119)..(139)  
 <223> Tetanus toxoid P30 epitope (SEQ ID NO: 24)

<220>  
 <221> SIMILAR  
 <222> (1)..(85)  
 <223> Identical to residues 1-85 in SEQ ID NO: 1

<220>  
 <221> SIMILAR  
 <222> (101)..(118)  
 <223> Identical to residues 92-109 in SEQ ID NO: 1

<220>  
 <221> SIMILAR  
 <222> (140)..(141)  
 <223> Identical to residues 114-115 in SEQ ID NO: 1

<400> 11  
 Ile Pro Thr Glu Ile Pro Thr Ser Ala Leu Val Lys Glu Thr Leu Ala  
           1                  5                  10                  15  
 Leu Leu Ser Thr His Arg Thr Leu Leu Ile Ala Asn Glu Thr Leu Arg  
                   20                  25                  30  
 Ile Pro Val Pro Val His Lys Asn His Gln Leu Cys Thr Glu Glu Ile  
           35                  40                  45  
 Phe Gln Gly Ile Gly Thr Leu Glu Ser Gln Thr Val Gln Gly Gly Thr  
           50                  55                  60  
 Val Glu Arg Leu Phe Lys Asn Leu Ser Leu Ile Lys Lys Tyr Ile Asp  
           65                  70                  75                  80  
 Gly Gln Lys Lys Lys Gln Tyr Ile Lys Ala Asn Ser Lys Phe Ile Gly  
                   85                  90                  95  
 Ile Thr Glu Leu Arg Val Asn Gln Phe Leu Asp Tyr Leu Gln Glu Phe  
           100                  105                  110  
 Leu Gly Val Met Asn Thr Phe Asn Asn Phe Thr Val Ser Phe Trp Leu  
           115                  120                  125  
 Arg Val Pro Lys Val Ser Ala Ser His Leu Glu Glu Ser  
           130                  135                  140

<210> 12  
 <211> 113  
 <212> PRT  
 <213> Mus musculus

09930916 "021902"

11

<220>  
 <221> DISULFID  
 <222> {42}  
 <223> Interchain disulphide bond to Cys-84 in SEQ ID  
 NO:12

<220>  
 <221> DISULFID  
 <222> {84}  
 <223> Interchain disulphide bond to Cys-42 in SEQ ID  
 NO:12

<400> 12  
 Met Glu Ile Pro Met Ser Thr Val Val Lys Glu Thr Leu Ala Leu Leu  
 1 5 10 15  
 Ser Ala His Arg Ala Leu Leu Thr Ser Asn Glu Thr Met Arg Leu Pro  
 20 25 30  
 Val Pro Thr His Lys Asn His Gln Leu Cys Ile Gly Glu Ile Phe Gln  
 35 40 45  
 Gly Leu Asp Ile Leu Lys Asp Gln Thr Val Arg Gly Gly Thr Val Met  
 50 55 60  
 Arg Leu Phe Gln Asn Leu Ser Leu Ile Lys Lys Tyr Ile Asp Arg Gln  
 65 70 75 80  
 Glu Lys Lys Cys Gly Glu Glu Arg Arg Arg Thr Arg Gln Phe Leu Asp  
 85 90 95  
 Tyr Leu Gln Glu Phe Leu Gly Ser Met Asn Thr Ala Ala Ile Ile Glu  
 100 105 110  
 Gly

<210> 13  
 <211> 124  
 <212> PRT  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: Murine IL5  
 modified by substitution with tetanus toxoid P2  
 epitope

<220>  
 <221> MUTAGEN  
 <222> (85)..(99)  
 <223> Tetanus toxoid P2 epitope (SEQ ID NO: 23)

<220>  
 <221> SIMILAR  
 <222> (1)..(84)  
 <223> Identical to residues 1-84 in SEQ ID NO: 12

<220>  
 <221> SIMILAR

09980916-02490

<223> Identical to residues 89-113 in SEQ ID NO: 12

Met Glu Ile Pro Met Ser Thr Val Val Lys Glu Thr Leu Ala Leu Leu  
1 5 10 15

Ser Ala His Arg Ala Leu Leu Thr Ser Asn Glu Thr Met Arg Leu Pro  
20 25 30

Val Pro Thr His Lys Asn His Gln Leu Cys Ile Gly Glu Ile Phe Gln  
35 40 45

Gly Leu Asp Ile Leu Lys Asp Gln Thr Val Arg Gly Gly Thr Val Met  
50 55 60

Arg Leu Phe Gln Asn Leu Ser Leu Ile Lys Lys Tyr Ile Asp Arg Gln  
65 70 75 80

Glu Lys Lys Cys Gln Tyr Ile Lys Ala Asn Ser Lys Phe Ile Gly Ile  
85 90 95

Thr Glu Leu Arg Arg Thr Arg Gln Phe Leu Asp Tyr Leu Gln Glu Phe  
100 105 110

Leu Gly Ser Met Asn Thr Ala Ala Ile Ile Glu Gly  
115 120

<211> 116

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Murine IL5  
modified by substitution with tetanus toxoid P2  
epitope

<220>

&lt;221&gt; MUTAGEN

<222> (30) .. (44)

<223> Tetanus toxoid P2 epitope (SEQ ID NO: 23)

**<220>**

<221> SIMILAR

<222> (1) . . (29)

<223> Identical to residues 1-29 in SEQ ID NO: 12

**<220>**

<221> SIMILAR

<222> (45) .. (116)

<223> Identical to residues 42-113 in SEQ ID NO: 12

<400> 14

Met Glu Ile Pro Met Ser Thr Val Val Lys Glu Thr Leu Ala Leu Leu  
1 5 10 15

Ser Ala His Arg Ala Leu Leu Thr Ser Asn Glu Thr Met Gln Tyr Ile  
20 25 30

Lys Ala Asn Ser Lys Phe Ile Gly Ile Thr Glu Leu Cys Ile Gly Glu  
           35                          40                          45  
 Ile Phe Gln Gly Leu Asp Ile Leu Lys Asp Gln Thr Val Arg Gly Gly  
           50                          55                          60  
 Thr Val Met Arg Leu Phe Gln Asn Leu Ser Leu Ile Lys Lys Tyr Ile  
           65                          70                          75                          80  
 Asp Arg Gln Glu Lys Lys Cys Gly Glu Glu Arg Arg Arg Thr Arg Gln  
                           85                          90                          95  
 Phe Leu Asp Tyr Leu Gln Glu Phe Leu Gly Ser Met Asn Thr Ala Ala  
                           100                          105                          110  
 Ile Ile Glu Gly  
           115

<210> 15  
 <211> 122  
 <212> PRT  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: Murine IL5  
       modified by substitution with tetanus toxoid P2  
       epitope

<220>  
 <221> MUTAGEN  
 <222> (57)..(71)  
 <223> Tetanus toxoid P2 epitope (SEQ ID NO: 23)

<220>  
 <221> SIMILAR  
 <222> (1)..(56)  
 <223> Identical to residues 1-56 in SEQ ID NO: 12

<220>  
 <221> SIMILAR  
 <222> (72)..(122)  
 <223> Identical to residues 63-113 in SEQ ID NO: 12

<400> 15  
 Met Glu Ile Pro Met Ser Thr Val Val Lys Glu Thr Leu Ala Leu Leu  
   1                          5                          10                          15  
 Ser Ala His Arg Ala Leu Leu Thr Ser Asn Glu Thr Met Arg Leu Pro  
           20                          25                          30  
 Val Pro Thr His Lys Asn His Gln Leu Cys Ile Gly Glu Ile Phe Gln  
           35                          40                          45  
 Gly Leu Asp Ile Leu Lys Asp Gln Gln Tyr Ile Lys Ala Asn Ser Lys  
           50                          55                          60  
 Phe Ile Gly Ile Thr Glu Leu Val Met Arg Leu Phe Gln Asn Leu Ser  
           65                          70                          75                          80

20051220 " 9T608550

14

Leu Ile Lys Lys Tyr Ile Asp Arg Gln Glu Lys Lys Cys Gly Glu Glu  
                             85                            90                            95

Arg Arg Arg Thr Arg Gln Phe Leu Asp Tyr Leu Gln Glu Phe Leu Gly  
                             100                            105                            110

Ser Met Asn Thr Ala Ala Ile Ile Glu Gly  
                             115                            120

<210> 16  
 <211> 122  
 <212> PRT  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: Murine IL5  
       modified by substitution with tetanus toxoid P2  
       epitope

<220>  
 <221> MUTAGEN  
 <222> (84)..(98)  
 <223> Tetanus toxoid P2 epitope (SEQ ID NO: 23)

<220>  
 <221> SIMILAR  
 <222> (1)..(83)  
 <223> Identical to residues 1-83 in SEQ ID NO: 12

<220>  
 <221> SIMILAR  
 <222> (99)..(122)  
 <223> Identical to residues 90-113 in SEQ ID NO: 12

<400> 16  
 Met Glu Ile Pro Met Ser Thr Val Val Lys Glu Thr Leu Ala Leu Leu  
       1                            5                            10                            15

Ser Ala His Arg Ala Leu Leu Thr Ser Asn Glu Thr Met Arg Leu Pro  
                             20                            25                            30

Val Pro Thr His Lys Asn His Gln Leu Cys Ile Gly Glu Ile Phe Gln  
                             35                            40                            45

Gly Leu Asp Ile Leu Lys Asp Gln Thr Val Arg Gly Gly Thr Val Met  
                             50                            55                            60

Arg Leu Phe Gln Asn Leu Ser Leu Ile Lys Lys Tyr Ile Asp Arg Gln  
                             65                            70                            75                            80

Glu Lys Lys Gln Tyr Ile Lys Ala Asn Ser Lys Phe Ile Gly Ile Thr  
                             85                            90                            95

Glu Leu Arg Thr Arg Gln Phe Leu Asp Tyr Leu Gln Glu Phe Leu Gly  
                             100                            105                            110

Ser Met Asn Thr Ala Ala Ile Ile Glu Gly  
                             115                            120

"Gibco" 91606660

<210> 17  
 <211> 124  
 <212> PRT  
 <213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Murine IL5  
 modified by substitution with tetanus toxoid P2  
 epitope

<220>

<221> MUTAGEN

<222> (108)..(122)

<223> Tetanus toxoid P2 epitope (SEQ ID NO: 23)

<220>

<221> SIMILAR

<222> (1)..(107)

<223> Identical to residues 1-107 in SEQ ID NO: 12

<220>

<221> SIMILAR

<222> (123)..(124)

<223> Identical to residues 112-113 in SEQ ID NO: 12

<400> 17

Met	Glu	Ile	Pro	Met	Ser	Thr	Val	Val	Lys	Glu	Thr	Leu	Ala	Leu	Leu
1				5					10					15	
Ser	Ala	His	Arg	Ala	Leu	Leu	Thr	Ser	Asn	Glu	Thr	Met	Arg	Leu	Pro
			20					25					30		
Val	Pro	Thr	His	Lys	Asn	His	Gln	Leu	Cys	Ile	Gly	Glu	Ile	Phe	Gln
			35				40					45			
Gly	Leu	Asp	Ile	Leu	Lys	Asp	Gln	Thr	Val	Arg	Gly	Gly	Thr	Val	Met
	50				55						60				
Arg	Leu	Phe	Gln	Asn	Leu	Ser	Leu	Ile	Lys	Lys	Tyr	Ile	Asp	Arg	Gln
	65				70					75					80
Glu	Lys	Lys	Cys	Gly	Glu	Glu	Arg	Arg	Arg	Thr	Arg	Gln	Phe	Leu	Asp
				85					90					95	
Tyr	Leu	Gln	Glu	Phe	Leu	Gly	Ser	Met	Asn	Thr	Gln	Tyr	Ile	Lys	Ala
			100					105					110		
Asn	Ser	Lys	Phe	Ile	Gly	Ile	Thr	Glu	Leu	Glu	Gly				
			115				120								

<210> 18  
 <211> 130  
 <212> PRT  
 <213> Artificial Sequence

<220>

2005-09-09 10:00:00

16

<223> Description of Artificial Sequence: Murine IL5  
modified by substitution with tetanus toxoid P30  
epitope

<220>

<221> MUTAGEN

<222> (85)..(105)

<223> Tetanus toxoid P2 epitope (SEQ ID NO: 24)

<220>

<221> SIMILAR

<222> (1)..(84)

<223> Identical to residues 1-84 in SEQ ID NO: 12

<220>

<221> SIMILAR

<222> (106)..(130)

<223> Identical to residues 89-113 in SEQ ID NO: 12

<400> 18

Met	Glu	Ile	Pro	Met	Ser	Thr	Val	Val	Lys	Glu	Thr	Leu	Ala	Leu	Leu
1				5					10					15	

Ser	Ala	His	Arg	Ala	Leu	Leu	Thr	Ser	Asn	Glu	Thr	Met	Arg	Leu	Pro
			20					25					30		

Val	Pro	Thr	His	Lys	Asn	His	Gln	Leu	Cys	Ile	Gly	Glu	Ile	Phe	Gln
		35					40					45			

Gly	Leu	Asp	Ile	Leu	Lys	Asp	Gln	Thr	Val	Arg	Gly	Gly	Thr	Val	Met
	50					55					60				

Arg	Leu	Phe	Gln	Asn	Leu	Ser	Leu	Ile	Lys	Lys	Tyr	Ile	Asp	Arg	Gln
	65				70					75					80

Glu	Lys	Lys	Cys	Phe	Asn	Asn	Phe	Thr	Val	Ser	Phe	Trp	Leu	Arg	Val
				85					90					95	

Pro	Lys	Val	Ser	Ala	Ser	His	Leu	Glu	Arg	Arg	Thr	Arg	Gln	Phe	Leu
			100					105					110		

Asp	Tyr	Leu	Gln	Glu	Phe	Leu	Gly	Ser	Met	Asn	Thr	Ala	Ala	Ile	Ile
		115					120						125		

Glu	Gly
	130

<210> 19

<211> 122

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Murine IL5  
modified by substitution with tetanus toxoid P30  
epitope

<220>

<221> MUTAGEN

09580916-004502



<220>

<221> SIMILAR

<222> (78)..(128)

<223> Identical to residues 63-113 in SEQ ID NO: 12

<400> 20

Met Glu Ile Pro Met Ser Thr Val Val Lys Glu Thr Leu Ala Leu Leu  
1 5 10 15

Ser Ala His Arg Ala Leu Leu Thr Ser Asn Glu Thr Met Arg Leu Pro  
20 25 30

Val Pro Thr His Lys Asn His Gln Leu Cys Ile Gly Glu Ile Phe Gln  
35 40 45

Gly Leu Asp Ile Leu Lys Asp Gln Phe Asn Asn Phe Thr Val Ser Phe  
50 55 60

Trp Leu Arg Val Pro Lys Val Ser Ala Ser His Leu Glu Val Met Arg  
65 70 75 80

Leu Phe Gln Asn Leu Ser Leu Ile Lys Lys Tyr Ile Asp Arg Gln Glu  
85 90 95

Lys Lys Cys Gly Glu Glu Arg Arg Arg Thr Arg Gln Phe Leu Asp Tyr  
100 105 110

Leu Gln Glu Phe Leu Gly Ser Met Asn Thr Ala Ala Ile Ile Glu Gly  
115 120 125

<210> 21

<211> 130

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Murine IL5  
modified by substitution with tetanus toxoid P30  
epitope

<220>

<221> MUTAGEN

<222> (108)..(128)

<223> Tetanus toxoid P30 epitope (SEQ ID NO: 24)

<220>

<221> SIMILAR

<222> (1)..(107)

<223> Identical to residues 1-107 in SEQ ID NO: 12

<220>

<221> SIMILAR

<222> (129)..(130)

<223> Identical to residues 112-113 in SEQ ID NO: 12

<400> 21

0060916-021902

19

Met Glu Ile Pro Met Ser Thr Val Val Lys Glu Thr Leu Ala Leu Leu  
 1 5 10 15

Ser Ala His Arg Ala Leu Leu Thr Ser Asn Glu Thr Met Arg Leu Pro  
 20 25 30

Val Pro Thr His Lys Asn His Gln Leu Cys Ile Gly Glu Ile Phe Gln  
 35 40 45

Gly Leu Asp Ile Leu Lys Asp Gln Thr Val Arg Gly Gly Thr Val Met  
 50 55 60

Arg Leu Phe Gln Asn Leu Ser Leu Ile Lys Lys Tyr Ile Asp Arg Gln  
 65 70 75 80

Glu Lys Lys Cys Gly Glu Glu Arg Arg Arg Thr Arg Gln Phe Leu Asp  
 85 90 95

Tyr Leu Gln Glu Phe Leu Gly Ser Met Asn Thr Phe Asn Asn Phe Thr  
 100 105 110

Val Ser Phe Trp Leu Arg Val Pro Lys Val Ser Ala Ser His Leu Glu  
 115 120 125

Glu Gly  
 130

<210> 22  
 <211> 139  
 <212> PRT  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: Murine IL5  
 modified by substitution with tetanus toxoid P2  
 and P30 epitopes

<220>  
 <221> MUTAGEN  
 <222> (84)..(98)  
 <223> Tetanus toxoid P2 epitope (SEQ ID NO: 23)

<220>  
 <221> MUTAGEN  
 <222> (117)..(137)  
 <223> Tetanus toxoid P30 epitope (SEQ ID NO: 24)

<220>  
 <221> SIMILAR  
 <222> (1)..(83)  
 <223> Identical to residues 1-83 in SEQ ID NO: 12

<220>  
 <221> SIMILAR  
 <222> (99)..(116)  
 <223> Identical to residues 90-109 in SEQ ID NO: 12

<220>

00590016 004300

<222> (138) .. (139)

<400> 22

<211> 45

```
<220>  
<221> CDS  
<222> (1)..(45)
```

<400> 25  
atg aaa cac caa cac caa cat caa cat caa cat caa cat caa caa 45  
Met Lys His Gln His Gln His Gln His Gln His Gln His Gln Gln  
1 5 10 15

<400> 26  
Met Lys His Gln His Gln His Gln His Gln His Gln His Gln Gln  
1 5 10 15

<220>  
<223> Description of Artificial Sequence: Human IL-5  
modified by substitution with tetanus toxoid  
epitope

```
<220>
<221> mutation
<222> (262)..(306)
<223> Tetanus toxoid P2 epitope
```

```
<220>
<221> misc_feature
<222> (307)..(378)
<223> DNA encoding amino acids 92-115 of human IL5
```

```

<400> 27
atc ccc aca gaa att ccc aca agt gca ttg gtg aaa gag acc ttg gca 48
Ile Pro Thr Glu Ile Pro Thr Ser Ala Leu Val Lys Glu Thr Leu Ala
      1             5             10             15

```

ctg ctt tct act cat cga act ctg ctg ata gcc aat gag act ctc cgg 96  
Leu Leu Ser Thr His Arg Thr Leu Leu Ile Ala Asn Glu Thr Leu Arg  
20 25 30

att cct gtt cct gta cat aaa aat cac caa ctg tgc act gaa gaa atc 144  
Ile Pro Val Pro Val His Lys Asn His Gln Leu Cys Thr Glu Glu Ile  
35 40 45

ttt cag gga ata ggc aca ctc gag agt caa act gtg caa ggg ggt act 192  
Phe Gln Gly Ile Gly Thr Leu Glu Ser Gln Thr Val Gln Gly Gly Thr  
50 55 60

gtg gaa aga cta ttc aaa aac ttg tcc tta ata aag aaa tac atc gat 240  
Val Glu Arg Leu Phe Lys Asn Leu Ser Leu Ile Lys Lys Tyr Ile Asp  
65 70 75 80

ggc caa aaa aaa aag tgt gga cag tac atc aag gcc aac tcc aag ttc 288  
Gly Gln Lys Lys Lys Cys Gly Gln Tyr Ile Lys Ala Asn Ser Lys Phe  
85 90 95

atc ggc atc acc gag ctg aga gta aac caa ttc cta gac tat ctg cag 336  
Ile Gly Ile Thr Glu Leu Arg Val Asn Gln Phe Leu Asp Tyr Leu Gln  
100 105 110

gag ttt ctt ggt gta atg aac acc gag tgg ata ata gaa agt tga 381  
Glu Phe Leu Gly Val Met Asn Thr Glu Trp Ile Ile Glu Ser  
115 120 125

<223> Description of Artificial Sequence: Human IL-5  
modified by substitution with tetanus toxoid  
epitope

Ile Pro Thr Glu Ile Pro Thr Ser Ala Leu Val Lys Glu Thr Leu Ala  
1 5 10 15

Leu Leu Ser Thr His Arg Thr Leu Leu Ile Ala Asn Glu Thr Leu Arg  
20 25 30

Ile Pro Val Pro Val His Lys Asn His Gln Leu Cys Thr Glu Glu Ile  
35 40 45

Phe Gln Gly Ile Gly Thr Leu Glu Ser Gln Thr Val Gln Gly Gly Thr  
50 55 60

Val Glu Arg Leu Phe Lys Asn Leu Ser Leu Ile Lys Lys Tyr Ile Asp  
65 70 75 80

Gly Gln Lys Lys Lys Cys Gly Gln Tyr Ile Lys Ala Asn Ser Lys Phe  
85 90 95

Ile Gly Ile Thr Glu Leu Arg Val Asn Gln Phe Leu Asp Tyr Leu Gln  
100 105 110

Glu Phe Leu Gly Val Met Asn Thr Glu Trp Ile Ile Glu Ser  
115 120 125

[illegible]

<210> 29  
 <211> 375  
 <212> DNA  
 <213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Human IL-5  
 modified by substitution with tetanus toxoid  
 epitope

<220>

<221> CDS

<222> (1)..(375)

<220>

<221> mutation

<222> (94)..(156)

<223> Tetanus toxoid P30 epitope

<220>

<221> misc\_feature

<222> (1)..(93)

<223> DNA encoding amino acids 1-31 of human IL5

<220>

<221> misc\_feature

<222> (157)..(372)

<223> DNA encoding amino acids 44-115 of human IL5

<400> 29

atc ccc aca gaa att ccc aca agt gca ttg gtg aaa gag acc ttg gca	48
Ile Pro Thr Glu Ile Pro Thr Ser Ala Leu Val Lys Glu Thr Leu Ala	
1 5 10 15	
ctg ctt tct act cat cga act ctg ctg ata gcc aat gag act ctc ttc	96
Leu Leu Ser Thr His Arg Thr Leu Leu Ile Ala Asn Glu Thr Leu Phe	
20 25 30	
aac aac ttc acc gtg agc ttc tgg ctg cgc gtg cct aag gtg agc gcc	144
Asn Asn Phe Thr Val Ser Phe Trp Leu Arg Val Pro Lys Val Ser Ala	
35 40 45	
agc cac ctg gag tgc act gaa gaa atc ttt cag gga ata ggc aca ctc	192
Ser His Leu Glu Cys Thr Glu Glu Ile Phe Gln Gly Ile Gly Thr Leu	
50 55 60	
gag agt caa act gtg caa ggg ggt act gtg gaa aga cta ttc aaa aac	240
Glu Ser Gln Thr Val Gln Gly Gly Thr Val Glu Arg Leu Phe Lys Asn	
65 70 75 80	
ttg tcc tta ata aag aaa tac atc gat ggc caa aaa aaa aag tgt gga	288
Leu Ser Leu Ile Lys Lys Tyr Ile Asp Gly Gln Lys Lys Lys Cys Gly	
85 90 95	
gaa gaa aga cgg aga gta aac caa ttc cta gac tat ctg cag gag ttt	336
Glu Glu Arg Arg Arg Val Asn Gln Phe Leu Asp Tyr Leu Gln Glu Phe	
100 105 110	

0900916 091502

24

ctt ggt gta atg aac acc gag tgg ata ata gaa agt tga  
 Leu Gly Val Met Asn Thr Glu Trp Ile Ile Glu Ser  
           115                                  120

375

&lt;210&gt; 30

&lt;211&gt; 124

&lt;212&gt; PRT

&lt;213&gt; Artificial Sequence

<223> Description of Artificial Sequence: Human Il-5  
 modified by substitution with tetanus toxoid  
 epitope

&lt;400&gt; 30

Ile Pro Thr Glu Ile Pro Thr Ser Ala Leu Val Lys Glu Thr Leu Ala  
       1                                  5                                  10                                  15

Leu Leu Ser Thr His Arg Thr Leu Leu Ile Ala Asn Glu Thr Leu Phe  
                                   20                                  25                                  30

Asn Asn Phe Thr Val Ser Phe Trp Leu Arg Val Pro Lys Val Ser Ala  
                                   35                                  40                                  45

Ser His Leu Glu Cys Thr Glu Glu Ile Phe Gln Gly Ile Gly Thr Leu  
                                   50                                  55                                  60

Glu Ser Gln Thr Val Gln Gly Gly Thr Val Glu Arg Leu Phe Lys Asn  
                                   65                                  70                                  75                                  80

Leu Ser Leu Ile Lys Lys Tyr Ile Asp Gly Gln Lys Lys Lys Cys Gly  
                                   85                                  90                                  95

Glu Glu Arg Arg Arg Val Asn Gln Phe Leu Asp Tyr Leu Gln Glu Phe  
                                   100                                  105                                  110

Leu Gly Val Met Asn Thr Glu Trp Ile Ile Glu Ser  
                                   115                                  120

&lt;210&gt; 31

&lt;211&gt; 393

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

<223> Description of Artificial Sequence: Human Il-5  
 modified by substitution with tetanus toxoid  
 epitope

&lt;220&gt;

&lt;221&gt; CDS

&lt;222&gt; (1)..(393)

&lt;220&gt;

&lt;221&gt; mutation

&lt;222&gt; (175)..(237)

&lt;223&gt; Tetanus toxoid P30 epitope

&lt;220&gt;

05080916 004902



<221> misc\_feature

<222> (1)..(174)

<223> DNA encoding amino acids 1-58 of human IL5

<220>

<221> misc\_feature

<222> (238)..(390)

<223> DNA encoding amino acids 65-115 of human IL5

<400> 31

atc	ccc	aca	gaa	att	ccc	aca	agt	gca	ttg	gtg	aaa	gag	acc	ttg	gca	48
Ile	Pro	Thr	Glu	Ile	Pro	Thr	Ser	Ala	Leu	Val	Lys	Glu	Thr	Leu	Ala	
1				5				10						15		

ctg	ctt	tct	act	cat	cga	act	ctg	ctg	ata	gcc	aat	gag	act	ctc	cgg	96
Leu	Leu	Ser	Thr	His	Arg	Thr	Leu	Leu	Ile	Ala	Asn	Glu	Thr	Leu	Arg	
			20				25					30				

att	cct	gtt	cct	gta	cat	aaa	aat	cac	caa	ctg	tgc	act	gaa	gaa	atc	144
Ile	Pro	Val	Pro	Val	His	Lys	Asn	His	Gln	Leu	Cys	Thr	Glu	Glu	Ile	
		35				40					45					

ttt	cag	gga	ata	ggc	aca	ctc	gag	agt	caa	ttc	aac	aac	ttc	acc	gtg	192
Phe	Gln	Gly	Ile	Gly	Thr	Leu	Glu	Ser	Gln	Phe	Asn	Asn	Phe	Thr	Val	
	50					55					60					

agc	ttc	tgg	ctg	cgc	gtg	cct	aag	gtg	agc	gcc	agc	cac	ctg	gag	gtg	240
Ser	Phe	Trp	Leu	Arg	Val	Pro	Lys	Val	Ser	Ala	Ser	His	Leu	Glu	Val	
65					70					75				80		

gaa	aga	cta	ttc	aaa	aac	ttg	tcc	tta	ata	aag	aaa	tac	atc	gat	ggc	288
Glu	Arg	Leu	Phe	Lys	Asn	Leu	Ser	Leu	Ile	Lys	Lys	Tyr	Ile	Asp	Gly	
				85				90						95		

caa	aaa	aaa	aag	tgt	gga	gaa	gaa	aga	cgg	aga	gta	aac	caa	ttc	cta	336
Gln	Lys	Lys	Lys	Cys	Gly	Glu	Glu	Arg	Arg	Arg	Val	Asn	Gln	Phe	Leu	
			100					105					110			

gac	tat	ctg	cag	gag	ttt	ctt	ggt	gta	atg	aac	acc	gag	tgg	ata	ata	384
Asp	Tyr	Leu	Gln	Glu	Phe	Leu	Gly	Val	Met	Asn	Thr	Glu	Trp	Ile	Ile	
		115					120					125				

gaa	agt	tga														393
Glu	Ser															
		130														

<210> 32

<211> 130

<212> PRT

<213> Artificial Sequence

<223> Description of Artificial Sequence: Human Il-5  
modified by substitution with tetanus toxoid  
epitope

<400> 32

Ile	Pro	Thr	Glu	Ile	Pro	Thr	Ser	Ala	Leu	Val	Lys	Glu	Thr	Leu	Ala	
1				5				10						15		

Leu	Leu	Ser	Thr	His	Arg	Thr	Leu	Leu	Ile	Ala	Asn	Glu	Thr	Leu	Arg	
			20				25						30			

"02390" 916 05530 05530

[illegible]

```
<210> 33
<211> 375
<212> DNA
<213> Artificial Sequence
```

<220>  
<223> Description of Artificial Sequence: Human Il-5  
modified by substitution with tetanus toxoid  
epitope

```
<220>
<221> CDS
<222> (1) .. (375)
```

```
<220>
<221> mutation
<222> (175)..(219)
<223> Tetanus toxoid P2 epitope
```

```
<220>  
<221> misc_feature  
<222> (1)..(174)  
<223> DNA encoding amino acids 1-58 of human IL5
```

```
<220>  
<221> misc_feature  
<222> (220)..(372)  
<223> DNA encoding amino acids 65-115 of human IL5
```

```

<400> 33
atc ccc aca gaa att ccc aca agt gca ttg gtg aaa gag acc ttg gca 48
Ile Pro Thr Glu Ile Pro Thr Ser Ala Leu Val Lys Glu Thr Leu Ala
      1                      5                      10                      15

ctg ctt tct act cat cga act ctg ctg ata gcc aat gag act ctc cgg 96
Leu Leu Ser Thr His Arg Thr Leu Leu Ile Ala Asn Glu Thr Leu Arg
      20                      25                      30

```

Parameter	Value
Initial temperature (K)	300
Final temperature (K)	1000
Time step (s)	0.01
Number of time steps	10000
Initial velocity (m/s)	0
Final velocity (m/s)	10
Time step (s)	0.01
Number of time steps	10000
Initial position (m)	0
Final position (m)	10
Time step (s)	0.01
Number of time steps	10000
Initial energy (J)	0
Final energy (J)	10
Time step (s)	0.01
Number of time steps	10000
Initial mass (kg)	1
Final mass (kg)	1
Time step (s)	0.01
Number of time steps	10000
Initial pressure (Pa)	101325
Final pressure (Pa)	101325
Time step (s)	0.01
Number of time steps	10000
Initial density (kg/m³)	1.225
Final density (kg/m³)	1.225
Time step (s)	0.01
Number of time steps	10000
Initial viscosity (Pa·s)	1.81e-05
Final viscosity (Pa·s)	1.81e-05
Time step (s)	0.01
Number of time steps	10000
Initial thermal conductivity (W/m·K)	0.0257
Final thermal conductivity (W/m·K)	0.0257
Time step (s)	0.01
Number of time steps	10000
Initial specific heat capacity (J/kg·K)	1000
Final specific heat capacity (J/kg·K)	1000
Time step (s)	0.01
Number of time steps	10000
Initial Prandtl number	0.7
Final Prandtl number	0.7
Time step (s)	0.01
Number of time steps	10000
Initial Schmidt number	0.7
Final Schmidt number	0.7
Time step (s)	0.01
Number of time steps	10000
Initial Lewis number	0.7
Final Lewis number	0.7
Time step (s)	0.01
Number of time steps	10000
Initial Peclet number	1000
Final Peclet number	1000
Time step (s)	0.01
Number of time steps	10000
Initial Fourier number	0.01
Final Fourier number	0.01
Time step (s)	0.01
Number of time steps	10000
Initial Biot number	1000
Final Biot number	1000
Time step (s)	0.01
Number of time steps	10000
Initial Stefan number	1000
Final Stefan number	1000
Time step (s)	0.01
Number of time steps	10000
Initial Rayleigh number	1000
Final Rayleigh number	1000
Time step (s)	0.01
Number of time steps	10000
Initial Grashof number	1000
Final Grashof number	1000
Time step (s)	0.01
Number of time steps	10000
Initial Prandtl number	0.7
Final Prandtl number	0.7
Time step (s)	0.01
Number of time steps	10000
Initial Schmidt number	0.7
Final Schmidt number	0.7
Time step (s)	0.01
Number of time steps	10000
Initial Lewis number	0.7
Final Lewis number	0.7
Time step (s)	0.01
Number of time steps	10000
Initial Peclet number	1000
Final Peclet number	1000
Time step (s)	0.01
Number of time steps	10000
Initial Fourier number	0.01
Final Fourier number	0.01
Time step (s)	0.01
Number of time steps	10000
Initial Biot number	1000
Final Biot number	1000
Time step (s)	0.01
Number of time steps	10000
Initial Stefan number	1000
Final Stefan number	1000
Time step (s)	0.01
Number of time steps	10000
Initial Rayleigh number	1000
Final Rayleigh number	1000
Time step (s)	0.01
Number of time steps	10000
Initial Grashof number	1000
Final Grashof number	1000
Time step (s)	0.01
Number of time steps	10000
Initial Prandtl number	0.7
Final Prandtl number	0.7
Time step (s)	0.01
Number of time steps	10000
Initial Schmidt number	0.7
Final Schmidt number	0.7
Time step (s)	0.01
Number of time steps	10000
Initial Lewis number	0.7
Final Lewis number	0.7
Time step (s)	0.01
Number of time steps	10000
Initial Peclet number	1000
Final Peclet number	1000
Time step (s)	0.01
Number of time steps	10000
Initial Fourier number	0.01
Final Fourier number	0.01
Time step (s)	0.01
Number of time steps	10000
Initial Biot number	1000
Final Biot number	1000
Time step (s)	0.01
Number of time steps	10000
Initial Stefan number	1000
Final Stefan number	1000
Time step (s)	0.01
Number of time steps	10000
Initial Rayleigh number	1000
Final Rayleigh number	1000
Time step (s)	0.01
Number of time steps	10000
Initial Grashof number	1000
Final Grashof number	1000
Time step (s)	0.01
Number of time steps	10000
Initial Prandtl number	0.7

Leu Gly Val Met Asn Thr Glu Trp Ile Ile Glu Ser  
115 120

<210> 35  
 <211> 357  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: Human Il-5  
 modified by substitution with tetanus toxoid  
 epitope

<220>  
 <221> CDS  
 <222> (1)..(357)

<220>  
 <221> mutation  
 <222> (94)..(138)  
 <223> Tetanus toxoid P2 epitope

<220>  
 <221> misc\_feature  
 <222> (1)..(93)  
 <223> DNA encoding amino acids 1-31 of human IL5

<220>  
 <221> misc\_feature  
 <222> (139)..(354)  
 <223> DNA encoding amino acids 44-115 of human IL5

<400> 35  
 atc ccc aca gaa att ccc aca agt gca ttg gtg aaa gag acc ttg gca 48  
 Ile Pro Thr Glu Ile Pro Thr Ser Ala Leu Val Lys Glu Thr Leu Ala  
 1 5 10 15  
 ctg ctt tct act cat cga act ctg ctg ata gcc aat gag act ctc cag 96  
 Leu Leu Ser Thr His Arg Thr Leu Leu Ile Ala Asn Glu Thr Leu Gln  
 20 25 30  
 tac atc aag gcc aac tcc aag ttc atc ggc atc acc gag ctg tgc act 144  
 Tyr Ile Lys Ala Asn Ser Lys Phe Ile Gly Ile Thr Glu Leu Cys Thr  
 35 40 45  
 gaa gaa atc ttt cag gga ata ggc aca ctc gag agt caa act gtg caa 192  
 Glu Glu Ile Phe Gln Gly Ile Gly Thr Leu Glu Ser Gln Thr Val Gln  
 50 55 60  
 ggg ggt act gtg gaa aga cta ttc aaa aac ttg tcc tta ata aag aaa 240  
 Gly Gly Thr Val Glu Arg Leu Phe Lys Asn Leu Ser Leu Ile Lys Lys  
 65 70 75 80  
 tac atc gat ggc caa aaa aaa aag tgt gga gaa gaa aga cgg aga gta 288  
 Tyr Ile Asp Gly Gln Lys Lys Lys Cys Gly Glu Glu Arg Arg Arg Val  
 85 90 95  
 aac caa ttc cta gac tat ctg cag gag ttt ctt ggt gta atg aac acc 336  
 Asn Gln Phe Leu Asp Tyr Leu Gln Glu Phe Leu Gly Val Met Asn Thr  
 100 105 110

20051201 91608650

gag tgg ata ata gaa agt tga  
 Glu Trp Ile Ile Glu Ser  
 115

357

<210> 36  
 <211> 118  
 <212> PRT  
 <213> Artificial Sequence  
 <223> Description of Artificial Sequence: Human Il-5  
 modified by substitution with tetanus toxoid  
 epitope

<400> 36  
 Ile Pro Thr Glu Ile Pro Thr Ser Ala Leu Val Lys Glu Thr Leu Ala  
 1 5 10 15  
 Leu Leu Ser Thr His Arg Thr Leu Leu Ile Ala Asn Glu Thr Leu Gln  
 20 25 30  
 Tyr Ile Lys Ala Asn Ser Lys Phe Ile Gly Ile Thr Glu Leu Cys Thr  
 35 40 45  
 Glu Glu Ile Phe Gln Gly Ile Gly Thr Leu Glu Ser Gln Thr Val Gln  
 50 55 60  
 Gly Gly Thr Val Glu Arg Leu Phe Lys Asn Leu Ser Leu Ile Lys Lys  
 65 70 75 80  
 Tyr Ile Asp Gly Gln Lys Lys Lys Cys Gly Glu Glu Arg Arg Arg Val  
 85 90 95  
 Asn Gln Phe Leu Asp Tyr Leu Gln Glu Phe Leu Gly Val Met Asn Thr  
 100 105 110  
 Glu Trp Ile Ile Glu Ser  
 115

<210> 37  
 <211> 375  
 <212> DNA  
 <213> Artificial Sequence  
 <220>  
 <223> Description of Artificial Sequence: Human Il-5  
 modified by substitution with tetanus toxoid  
 epitope  
 <220>  
 <221> CDS  
 <222> (1)..(375)  
 <220>  
 <221> mutation  
 <222> (256)..(300)  
 <223> Tetanus toxoid P2 epitope  
 <220>

```
<221> misc_feature
<222> (1)..(255)
<223> DNA encoding amino acids 1-85 of human IL5

<220>
<221> misc_feature
<222> (301)..(372)
<223> DNA encoding amino acids 92-115 of human IL5
```

<400> 37																	
atc	ccc	aca	gaa	att	ccc	aca	agt	gca	ttg	gtg	aaa	gag	acc	ttg	gca	48	
Ile	Pro	Thr	Glu	Ile	Pro	Thr	Ser	Ala	Leu	Val	Lys	Glu	Thr	Leu	Ala		
1				5					10					15			
ctg	ctt	tct	act	cat	cga	act	ctg	ctg	ata	gcc	aat	gag	act	ctc	cgg	96	
Leu	Leu	Ser	Thr	His	Arg	Thr	Leu	Leu	Ile	Ala	Asn	Glu	Thr	Leu	Arg		
			20					25					30				
att	cct	ggt	cct	gta	cat	aaa	aat	cac	caa	ctg	tgc	act	gaa	gaa	atc	144	
Ile	Pro	Val	Pro	Val	His	Lys	Asn	His	Gln	Leu	Cys	Thr	Glu	Glu	Ile		
		35					40					45					
ttt	cag	gga	ata	ggc	aca	ctc	gag	agt	caa	act	gtg	caa	ggg	ggt	act	192	
Phe	Gln	Gly	Ile	Gly	Thr	Leu	Glu	Ser	Gln	Thr	Val	Gln	Gly	Gly	Thr		
	50					55					60						
gtg	gaa	aga	cta	ttc	aaa	aac	ttg	tcc	tta	ata	aag	aaa	tac	atc	gat	240	
Val	Glu	Arg	Leu	Phe	Lys	Asn	Leu	Ser	Leu	Ile	Lys	Lys	Tyr	Ile	Asp		
65					70					75					80		
ggc	caa	aaa	aaa	aag	cag	tac	atc	aag	gcc	aac	tcc	aag	ttc	atc	ggc	288	
Gly	Gln	Lys	Lys	Lys	Gln	Tyr	Ile	Lys	Ala	Asn	Ser	Lys	Phe	Ile	Gly		
				85					90					95			
atc	acc	gag	ctg	aga	gta	aac	caa	ttc	cta	gac	tat	ctg	cag	gag	ttt	336	
Ile	Thr	Glu	Leu	Arg	Val	Asn	Gln	Phe	Leu	Asp	Tyr	Leu	Gln	Glu	Phe		
			100					105					110				
ctt	ggt	gta	atg	aac	acc	gag	tgg	ata	ata	gaa	agt	tga				375	
Leu	Gly	Val	Met	Asn	Thr	Glu	Trp	Ile	Ile	Glu	Ser						
		115					120										

```
<210> 38
<211> 124
<212> PRT
<213> Artificial Sequence
<223> Description of Artificial Sequence: Human Il-5
        modified by substitution with tetanus toxoid
        epitope
```

```

<400> 38
Ile Pro Thr Glu Ile Pro Thr Ser Ala Leu Val Lys Glu Thr Leu Ala
  1             5             10             15

Leu Leu Ser Thr His Arg Thr Leu Leu Ile Ala Asn Glu Thr Leu Arg
      20             25             30

Ile Pro Val Pro Val His Lys Asn His Gln Leu Cys Thr Glu Glu Ile
      35             40             45

```

[illegible]

31

Phe Gln Gly Ile Gly Thr Leu Glu Ser Gln Thr Val Gln Gly Gly Thr  
 50 55 60  
 Val Glu Arg Leu Phe Lys Asn Leu Ser Leu Ile Lys Lys Tyr Ile Asp  
 65 70 75 80  
 Gly Gln Lys Lys Lys Gln Tyr Ile Lys Ala Asn Ser Lys Phe Ile Gly  
 85 90 95  
 Ile Thr Glu Leu Arg Val Asn Gln Phe Leu Asp Tyr Leu Gln Glu Phe  
 100 105 110  
 Leu Gly Val Met Asn Thr Glu Trp Ile Ile Glu Ser  
 115 120

&lt;210&gt; 39

&lt;211&gt; 399

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

<223> Description of Artificial Sequence: Human IL-5  
 modified by substitution with tetanus toxoid  
 epitope

&lt;220&gt;

&lt;221&gt; CDS

&lt;222&gt; (1)..(399)

&lt;220&gt;

&lt;221&gt; mutation

&lt;222&gt; (262)..(324)

&lt;223&gt; Tetanus toxoid P30 epitope

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;222&gt; (1)..(261)

&lt;223&gt; DNA encoding amino acids 1-87 of human IL5

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;222&gt; (325)..(396)

&lt;223&gt; DNA encoding amino acids 92-115 of human IL5

&lt;400&gt; 39

atc ccc aca gaa att ccc aca agt gca ttg gtg aaa gag acc ttg gca 48  
 Ile Pro Thr Glu Ile Pro Thr Ser Ala Leu Val Lys Glu Thr Leu Ala  
 1 5 10 15

ctg ctt tct act cat cga act ctg ctg ata gcc aat gag act ctc cgg 96  
 Leu Leu Ser Thr His Arg Thr Leu Leu Ile Ala Asn Glu Thr Leu Arg  
 20 25 30

att cct gtt cct gta cat aaa aat cac caa ctg tgc act gaa gaa atc 144  
 Ile Pro Val Pro Val His Lys Asn His Gln Leu Cys Thr Glu Glu Ile  
 35 40 45

"CTG" 9750045 001000

ttt cag gga ata ggc aca ctc gag agt caa act gtg caa ggg ggt act 192  
Phe Gln Gly Ile Gly Thr Leu Glu Ser Gln Thr Val Gln Gly Gly Thr  
50 55 60

gtg gaa aga cta ttc aaa aac ttg tcc tta ata aag aaa tac atc gat 240  
Val Glu Arg Leu Phe Lys Asn Leu Ser Leu Ile Lys Lys Tyr Ile Asp  
65 70 75 80

ggc caa aaa aaa aag tgt gga ttc aac aac ttc acc gtg agc ttc tgg 288  
Gly Gln Lys Lys Lys Cys Gly Phe Asn Asn Phe Thr Val Ser Phe Trp  
85 90 95

ctg cgc gtg cct aag gtg agc gcc agc cac ctg gag aga gta aac caa 336  
Leu Arg Val Pro Lys Val Ser Ala Ser His Leu Glu Arg Val Asn Gln  
100 105 110

ttc cta gac tat ctg cag gag ttt ctt ggt gta atg aac acc gag tgg 384  
Phe Leu Asp Tyr Leu Gln Glu Phe Leu Gly Val Met Asn Thr Glu Trp  
115 120 125

```

ata ata gaa agt tga
Ile Ile Glu Ser
130

```

<211> 132

<212> PRT

<213> Artificial Sequence

<223> Description of Artificial Sequence: Human Il-5  
modified by substitution with tetanus toxoid  
epitope

<400> 40

<400> 40  
Ile Pro Thr Glu Ile Pro Thr Ser Ala Leu Val Lys Glu Thr Leu Ala  
1 5 10 15

Leu Leu Ser Thr His Arg Thr Leu Leu Ile Ala Asn Glu Thr Leu Arg  
20 25 30

Ile Pro Val Pro Val His Lys Asn His Gln Leu Cys Thr Glu Glu Ile  
35 40 45

Phe Gln Gly Ile Gly Thr Leu Glu Ser Gln Thr Val Gln Gly Gly Thr  
50 55 60

Val Glu Arg Leu Phe Lys Asn Leu Ser Leu Ile Lys Lys Tyr Ile Asp  
65 70 75 80

Gly Gln Lys Lys Lys Cys Gly Phe Asn Asn Phe Thr Val Ser Phe Trp  
85 90 95

Leu Arg Val Pro Lys Val Ser Ala Ser His Leu Glu Arg Val Asn Gln  
100 105 110

Phe Leu Asp Tyr Leu Gln Glu Phe Leu Gly Val Met Asn Thr Glu Trp  
115 120 125

Ile Ile Glu Ser  
130

[illegible]



<210> 41  
 <211> 393  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: Human IL-5  
 modified by substitution with tetanus toxoid  
 epitope

<220>  
 <221> CDS  
 <222> (1)..(393)

<220>  
 <221> mutation  
 <222> (256)..(318)  
 <223> Tetanus toxoid P30 epitope

<220>  
 <221> misc\_feature  
 <222> (1)..(255)  
 <223> DNA encoding amino acids 1-85 of human IL5

<220>  
 <221> misc\_feature  
 <222> (319)..(390)  
 <223> DNA encoding amino acids 92-115 of human IL5

<400> 41  
 atc ccc aca gaa att ccc aca agt gca ttg gtg aaa gag acc ttg gca 48  
 Ile Pro Thr Glu Ile Pro Thr Ser Ala Leu Val Lys Glu Thr Leu Ala  
 1 5 10 15  
 ctg ctt tct act cat cga act ctg ctg ata gcc aat gag act ctc cgg 96  
 Leu Leu Ser Thr His Arg Thr Leu Leu Ile Ala Asn Glu Thr Leu Arg  
 20 25 30  
 att cct gtt cct gta cat aaa aat cac caa ctg tgc act gaa gaa atc 144  
 Ile Pro Val Pro Val His Lys Asn His Gln Leu Cys Thr Glu Glu Ile  
 35 40 45  
 ttt cag gga ata ggc aca ctc gag agt caa act gtg caa ggg ggt act 192  
 Phe Gln Gly Ile Gly Thr Leu Glu Ser Gln Thr Val Gln Gly Gly Thr  
 50 55 60  
 gtg gaa aga cta ttc aaa aac ttg tcc tta ata aag aaa tac atc gat 240  
 Val Glu Arg Leu Phe Lys Asn Leu Ser Leu Ile Lys Lys Tyr Ile Asp  
 65 70 75 80  
 ggc caa aaa aaa aag ttc aac aac ttc acc gtg agc ttc tgg ctg cgc 288  
 Gly Gln Lys Lys Lys Phe Asn Asn Phe Thr Val Ser Phe Trp Leu Arg  
 85 90 95  
 gtg cct aag gtg agc gcc agc cac ctg gag aga gta aac caa ttc cta 336  
 Val Pro Lys Val Ser Ala Ser His Leu Glu Arg Val Asn Gln Phe Leu  
 100 105 110

"GATEWAY" 97000000

34

gac tat ctg cag gag ttt ctt ggt gta atg aac acc gag tgg ata ata 384  
 Asp Tyr Leu Gln Glu Phe Leu Gly Val Met Asn Thr Glu Trp Ile Ile  
           115                          120                          125

gaa agt tga 393  
 Glu Ser  
       130

&lt;210&gt; 42

&lt;211&gt; 130

&lt;212&gt; PRT

&lt;213&gt; Artificial Sequence

<223> Description of Artificial Sequence: Human Il-5  
 modified by substitution with tetanus toxoid  
 epitope

&lt;400&gt; 42

Ile Pro Thr Glu Ile Pro Thr Ser Ala Leu Val Lys Glu Thr Leu Ala  
       1                          5                          10                          15

Leu Leu Ser Thr His Arg Thr Leu Leu Ile Ala Asn Glu Thr Leu Arg  
                           20                          25                          30

Ile Pro Val Pro Val His Lys Asn His Gln Leu Cys Thr Glu Glu Ile  
                           35                          40                          45

Phe Gln Gly Ile Gly Thr Leu Glu Ser Gln Thr Val Gln Gly Gly Thr  
       50                          55                          60

Val Glu Arg Leu Phe Lys Asn Leu Ser Leu Ile Lys Lys Tyr Ile Asp  
       65                          70                          75                          80

Gly Gln Lys Lys Lys Phe Asn Asn Phe Thr Val Ser Phe Trp Leu Arg  
                           85                          90                          95

Val Pro Lys Val Ser Ala Ser His Leu Glu Arg Val Asn Gln Phe Leu  
                           100                          105                          110

Asp Tyr Leu Gln Glu Phe Leu Gly Val Met Asn Thr Glu Trp Ile Ile  
                           115                          120                          125

Glu Ser  
       130

&lt;210&gt; 43

&lt;211&gt; 444

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

<223> Description of Artificial Sequence: Human Il-5  
 modified by substitution with tetanus toxoid  
 epitopes

&lt;220&gt;

&lt;221&gt; CDS

&lt;222&gt; (1)..(444)

"05T00" 9T600650

<220>  
 <221> mutation  
 <222> (262)..(306)  
 <223> Tetanus toxoid P2 epitope

<220>  
 <221> mutation  
 <222> (307)..(369)  
 <223> Tetanus toxoid P30 epitope

<220>  
 <221> misc\_feature  
 <222> (1)..(261)  
 <223> DNA encoding amino acids 1-87 of human IL5

<220>  
 <221> misc\_feature  
 <222> (370)..(441)  
 <223> DNA encoding amino acids 92-115 of human IL5

<400> 43  
 atc ccc aca gaa att ccc aca agt gca ttg gtg aaa gag acc ttg gca 48  
 ile pro thr glu ile pro thr ser ala leu val lys glu thr leu ala  
 1 5 10 15  
 ctg ctt tct act cat cga act ctg ctg ata gcc aat gag act ctc cgg 96  
 leu leu ser thr his arg thr leu leu ile ala asn glu thr leu arg  
 20 25 30  
 att cct gtt cct gta cat aaa aat cac caa ctg tgc act gaa gaa atc 144  
 ile pro val pro val his lys asn his gln leu cys thr glu glu ile  
 35 40 45  
 ttt cag gga ata ggc aca ctc gag agt caa act gtg caa ggg ggt act 192  
 phe gln gly ile gly thr leu glu ser gln thr val gln gly gly thr  
 50 55 60  
 gtg gaa aga cta ttc aaa aac ttg tcc tta ata aag aaa tac atc gat 240  
 val glu arg leu phe lys asn leu ser leu ile lys lys tyr ile asp  
 65 70 75 80  
 ggc caa aaa aaa aag tgt gga cag tac atc aag gcc aac tcc aag ttc 288  
 gly gln lys lys lys cys gly gln tyr ile lys ala asn ser lys phe  
 85 90 95  
 atc ggc atc acc gag ctg ttc aac aac ttc acc gtg agc ttc tgg ctg 336  
 ile gly ile thr glu leu phe asn asn phe thr val ser phe trp leu  
 100 105 110  
 cgc gtg cct aag gtg agc gcc agc cac ctg gag aga gta aac caa ttc 384  
 arg val pro lys val ser ala ser his leu glu arg val asn gln phe  
 115 120 125  
 cta gac tat ctg cag gag ttt ctt ggt gta atg aac acc gag tgg ata 432  
 leu asp tyr leu gln glu phe leu gly val met asn thr glu trp ile  
 130 135 140  
 ata gaa agt tga 444  
 ile glu ser  
 145

0590916 021506

<210> 44  
 <211> 147  
 <212> PRT  
 <213> Artificial Sequence  
 <223> Description of Artificial Sequence: Human IL-5  
 modified by substitution with tetanus toxoid  
 epitopes

<400> 44  
 Ile Pro Thr Glu Ile Pro Thr Ser Ala Leu Val Lys Glu Thr Leu Ala  
           1                          5                          10                          15  
 Leu Leu Ser Thr His Arg Thr Leu Leu Ile Ala Asn Glu Thr Leu Arg  
                           20                          25                          30  
 Ile Pro Val Pro Val His Lys Asn His Gln Leu Cys Thr Glu Glu Ile  
                           35                          40                          45  
 Phe Gln Gly Ile Gly Thr Leu Glu Ser Gln Thr Val Gln Gly Gly Thr  
           50                          55                          60  
 Val Glu Arg Leu Phe Lys Asn Leu Ser Leu Ile Lys Lys Tyr Ile Asp  
           65                          70                          75                          80  
 Gly Gln Lys Lys Lys Cys Gly Gln Tyr Ile Lys Ala Asn Ser Lys Phe  
                           85                          90                          95  
 Ile Gly Ile Thr Glu Leu Phe Asn Asn Phe Thr Val Ser Phe Trp Leu  
                           100                          105                          110  
 Arg Val Pro Lys Val Ser Ala Ser His Leu Glu Arg Val Asn Gln Phe  
           115                          120                          125  
 Leu Asp Tyr Leu Gln Glu Phe Leu Gly Val Met Asn Thr Glu Trp Ile  
           130                          135                          140  
 Ile Glu Ser  
 145

<210> 45  
 <211> 375  
 <212> DNA  
 <213> Artificial Sequence  
 <220>  
 <223> Description of Artificial Sequence: Murine IL-5  
 modified by substitution with tetanus toxoid  
 epitope

<220>  
 <221> CDS  
 <222> (1)..(375)  
 <220>  
 <221> mutation  
 <222> (256)..(300)  
 <223> Tetanus toxoid P2 epitope

09960916-024902

<220>  
 <221> misc\_feature  
 <222> (1)..(255)  
 <223> DNA encoding amino acids 1-85 in murine IL5

<220>  
 <221> misc\_feature  
 <222> (301)..(375)  
 <223> DNA encoding amino acids 90-113 of murine IL5

<400> 45  
 atg gag att ccc atg agc aca gtg gtg aaa gag acc ttg aca cag ctg 48  
 Met Glu Ile Pro Met Ser Thr Val Val Lys Glu Thr Leu Thr Gln Leu  
 1 5 10 15  
 tcc gct cac cga gct ctg ttg aca agc aat gag acg atg agg ctt cct 96  
 Ser Ala His Arg Ala Leu Leu Thr Ser Asn Glu Thr Met Arg Leu Pro  
 20 25 30  
 gtc cct act cat aaa aat cac cag cta tgc att gga gag atc ttt cag 144  
 Val Pro Thr His Lys Asn His Gln Leu Cys Ile Gly Glu Ile Phe Gln  
 35 40 45  
 ggg cta gac ata ctg aag aat caa act gtc cgt ggg ggt acc gtg gaa 192  
 Gly Leu Asp Ile Leu Lys Asn Gln Thr Val Arg Gly Gly Thr Val Glu  
 50 55 60  
 atg cta ttc caa aac ctg tca tta ata aag aaa tac atc gat aga caa 240  
 Met Leu Phe Gln Asn Leu Ser Leu Ile Lys Lys Tyr Ile Asp Arg Gln  
 65 70 75 80  
 aaa gag aag tgt ggc cag tac atc aaa gct aac tcc aaa ttc atc ggt 288  
 Lys Glu Lys Cys Gly Gln Tyr Ile Lys Ala Asn Ser Lys Phe Ile Gly  
 85 90 95  
 atc acc gag ctg agg acg agg cag ttc ctg gat tat ctg cag gag ttc 336  
 Ile Thr Glu Leu Arg Thr Arg Gln Phe Leu Asp Tyr Leu Gln Glu Phe  
 100 105 110  
 ctt ggt gtg atg agt aca gag tgg gca atg gaa ggc taa 375  
 Leu Gly Val Met Ser Thr Glu Trp Ala Met Glu Gly  
 115 120

<210> 46  
 <211> 124  
 <212> PRT  
 <213> Artificial Sequence  
 <223> Description of Artificial Sequence: Murine Il-5  
 modified by substitution with tetanus toxoid  
 epitope

<400> 46  
 Met Glu Ile Pro Met Ser Thr Val Val Lys Glu Thr Leu Thr Gln Leu  
 1 5 10 15  
 Ser Ala His Arg Ala Leu Leu Thr Ser Asn Glu Thr Met Arg Leu Pro  
 20 25 30  
 Val Pro Thr His Lys Asn His Gln Leu Cys Ile Gly Glu Ile Phe Gln  
 35 40 45

09980915 "021902

38

Gly Leu Asp Ile Leu Lys Asn Gln Thr Val Arg Gly Gly Thr Val Glu  
 50 55 60

Met Leu Phe Gln Asn Leu Ser Leu Ile Lys Lys Tyr Ile Asp Arg Gln  
 65 70 75 80

Lys Glu Lys Cys Gly Gln Tyr Ile Lys Ala Asn Ser Lys Phe Ile Gly  
 85 90 95

Ile Thr Glu Leu Arg Thr Arg Gln Phe Leu Asp Tyr Leu Gln Glu Phe  
 100 105 110

Leu Gly Val Met Ser Thr Glu Trp Ala Met Glu Gly  
 115 120

<210> 47  
 <211> 369  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: Murine Il-5  
 modified by substitution with tetanus toxoid  
 epitope

<220>  
 <221> CDS  
 <222> (1)..(369)

<220>  
 <221> mutation  
 <222> (88)..(150)  
 <223> Tetanus toxoid P30 epitope

<220>  
 <221> misc\_feature  
 <222> (1)..(87)  
 <223> DNA encoding amino acids 1-29 of murine IL5

<220>  
 <221> misc\_feature  
 <222> (151)..(366)  
 <223> DNA encoding amino acids 42-113 of murine IL5

<400> 47  
 atg gag att ccc atg agc aca gtg gtg aaa gag acc ttg aca cag ctg 48  
 Met Glu Ile Pro Met Ser Thr Val Val Lys Glu Thr Leu Thr Gln Leu  
 1 5 10 15

tcc gct cac cga gct ctg ttg aca agc aat gag acg atg ttc aac aac 96  
 Ser Ala His Arg Ala Leu Leu Thr Ser Asn Glu Thr Met Phe Asn Asn  
 20 25 30

ttc acc gtg agc ttc tgg ctg cgc gtg ccc aag gtg agc gcc agc cac 144  
 Phe Thr Val Ser Phe Trp Leu Arg Val Pro Lys Val Ser Ala Ser His  
 35 40 45

0065058 WO 00/65058

[illegible]

```
<210> 48
<211> 122
<212> PRT
<213> Artificial Sequence
<223> Description of Artificial Sequence: Murine Il-5
modified by substitution with tetanus toxoid
epitope
```

```

<400> 48
Met Glu Ile Pro Met Ser Thr Val Val Lys Glu Thr Leu Thr Gln Leu
  1           5          10           15

Ser Ala His Arg Ala Leu Leu Thr Ser Asn Glu Thr Met Phe Asn Asn
          20          25          30

Phe Thr Val Ser Phe Trp Leu Arg Val Pro Lys Val Ser Ala Ser His
          35          40          45

Leu Glu Cys Ile Gly Glu Ile Phe Gln Gly Leu Asp Ile Leu Lys Asn
  50           55           60

Gln Thr Val Arg Gly Gly Thr Val Glu Met Leu Phe Gln Asn Leu Ser
  65           70           75           80

Leu Ile Lys Lys Tyr Ile Asp Arg Gln Lys Glu Lys Cys Gly Glu Glu
          85          90          95

Arg Arg Arg Thr Arg Gln Phe Leu Asp Tyr Leu Gln Glu Phe Leu Gly
          100         105         110

Val Met Ser Thr Glu Trp Ala Met Glu Gly
      115          120

```

```
<210> 49
<211> 387
<212> DNA
<213> Artificial Sequence
```

General Information	
Author(s)	W. H. R. ...
Title	...
Journal	...
Volume	...
Issue	...
Pages	...
Year	...
DOI	...
Keywords	...
Abstract	...
Introduction	...
Methods	...
Results	...
Discussion	...
Conclusion	...
References	...
Appendix	...
Supplementary Material	...
Notes	...
Footnote	...
Page Number	...

<220>  
 <223> Description of Artificial Sequence: Murine Il-5  
 modified by substitution with tetanus toxoid  
 epitope

<220>  
 <221> CDS  
 <222> (1)..(387)

<220>  
 <221> mutation  
 <222> (169)..(231)  
 <223> Tetanus toxoid P2 epitope

<220>  
 <221> misc\_feature  
 <222> (1)..(168)  
 <223> DNA encoding amino acids 1-56 of murine IL5

<220>  
 <221> misc\_feature  
 <222> (232)..(384)  
 <223> DNA encoding amino acids 63-113 of murine IL5

<400> 49  
 atg gag att ccc atg agc aca gtg gtg aaa gag acc ttg aca cag ctg 48  
 Met Glu Ile Pro Met Ser Thr Val Val Lys Glu Thr Leu Thr Gln Leu  
 1 5 10 15  
 tcc gct cac cga gct ctg ttg aca agc aat gag acg atg agg ctt cct 96  
 Ser Ala His Arg Ala Leu Leu Thr Ser Asn Glu Thr Met Arg Leu Pro  
 20 25 30  
 gtc cct act cat aaa aat cac cag cta tgc att gga gag atc ttt cag 144  
 Val Pro Thr His Lys Asn His Gln Leu Cys Ile Gly Glu Ile Phe Gln  
 35 40 45  
 ggg cta gac ata ctg aag aat caa ttc aac aac ttc acc gtg agc ttc 192  
 Gly Leu Asp Ile Leu Lys Asn Gln Phe Asn Asn Phe Thr Val Ser Phe  
 50 55 60  
 tgg ctg cgc gtg ccc aag gtg agc gcc agc cac ctg gag gtg gaa atg 240  
 Trp Leu Arg Val Pro Lys Val Ser Ala Ser His Leu Glu Val Glu Met  
 65 70 75 80  
 cta ttc caa aac ctg tca tta ata aag aaa tac atc gat aga caa aaa 288  
 Leu Phe Gln Asn Leu Ser Leu Ile Lys Lys Tyr Ile Asp Arg Gln Lys  
 85 90 95  
 gag aag tgt ggc gag gag aga cgg agg acg agg cag ttc ctg gat tat 336  
 Glu Lys Cys Gly Glu Glu Arg Arg Thr Arg Gln Phe Leu Asp Tyr  
 100 105 110  
 ctg cag gag ttc ctt ggt gtg atg agt aca gag tgg gca atg gaa ggc 384  
 Leu Gln Glu Phe Leu Gly Val Met Ser Thr Glu Trp Ala Met Glu Gly  
 115 120 125  
 taa 387



<210> 50  
 <211> 128  
 <212> PRT  
 <213> Artificial Sequence  
 <223> Description of Artificial Sequence: Murine Il-5  
 modified by substitution with tetanus toxoid  
 epitope

<400> 50  
 Met Glu Ile Pro Met Ser Thr Val Val Lys Glu Thr Leu Thr Gln Leu  
 1 5 10 15  
 Ser Ala His Arg Ala Leu Leu Thr Ser Asn Glu Thr Met Arg Leu Pro  
 20 25 30  
 Val Pro Thr His Lys Asn His Gln Leu Cys Ile Gly Glu Ile Phe Gln  
 35 40 45  
 Gly Leu Asp Ile Leu Lys Asn Gln Phe Asn Asn Phe Thr Val Ser Phe  
 50 55 60  
 Trp Leu Arg Val Pro Lys Val Ser Ala Ser His Leu Glu Val Glu Met  
 65 70 75 80  
 Leu Phe Gln Asn Leu Ser Leu Ile Lys Lys Tyr Ile Asp Arg Gln Lys  
 85 90 95  
 Glu Lys Cys Gly Glu Glu Arg Arg Arg Thr Arg Gln Phe Leu Asp Tyr  
 100 105 110  
 Leu Gln Glu Phe Leu Gly Val Met Ser Thr Glu Trp Ala Met Glu Gly  
 115 120 125

<210> 51  
 <211> 351  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: Murine Il-5  
 modified by substitution with tetanus toxoid  
 epitope

<220>  
 <221> CDS  
 <222> (1)..(351)

<220>  
 <221> mutation  
 <222> (88)..(132)  
 <223> Tetanus toxoid P2 epitope

<220>  
 <221> misc\_feature  
 <222> (1)..(87)  
 <223> DNA encoding amino acids 1-29 of murine IL5

<220>

&lt;221&gt; misc\_feature

&lt;222&gt; (133)..(348)

&lt;223&gt; DNA encoding amino acids 42-113 of murine IL5

&lt;400&gt; 51

atg	gag	att	ccc	atg	agc	aca	gtg	gtg	aaa	gag	acc	ttg	aca	cag	ctg	48
Met	Glu	Ile	Pro	Met	Ser	Thr	Val	Val	Lys	Glu	Thr	Leu	Thr	Gln	Leu	
1				5					10					15		

tcc	gct	cac	cga	gct	ctg	ttg	aca	agc	aat	gag	acg	atg	cag	tac	atc	96
Ser	Ala	His	Arg	Ala	Leu	Leu	Thr	Ser	Asn	Glu	Thr	Met	Gln	Tyr	Ile	
			20					25					30			

aaa	gct	aac	tcc	aaa	ttc	atc	ggt	atc	acc	gag	ctg	tgc	att	gga	gag	144
Lys	Ala	Asn	Ser	Lys	Phe	Ile	Gly	Ile	Thr	Glu	Leu	Cys	Ile	Gly	Glu	
		35					40					45				

atc	ttt	cag	ggg	cta	gac	ata	ctg	aag	aat	caa	act	gtc	cgt	ggg	ggt	192
Ile	Phe	Gln	Gly	Leu	Asp	Ile	Leu	Lys	Asn	Gln	Thr	Val	Arg	Gly	Gly	
	50					55					60					

acc	gtg	gaa	atg	cta	ttc	caa	aac	ctg	tca	tta	ata	aag	aaa	tac	atc	240
Thr	Val	Glu	Met	Leu	Phe	Gln	Asn	Leu	Ser	Leu	Ile	Lys	Lys	Tyr	Ile	
	65				70					75					80	

gat	aga	caa	aaa	gag	aag	tgt	ggc	gag	gag	aga	cgg	agg	acg	agg	cag	288
Asp	Arg	Gln	Lys	Glu	Lys	Cys	Gly	Glu	Glu	Arg	Arg	Arg	Thr	Arg	Gln	
				85				90						95		

ttc	ctg	gat	tat	ctg	cag	gag	ttc	ctt	ggt	gtg	atg	agt	aca	gag	tgg	336
Phe	Leu	Asp	Tyr	Leu	Gln	Glu	Phe	Leu	Gly	Val	Met	Ser	Thr	Glu	Trp	
		100						105						110		

gca	atg	gaa	ggc	taa												351
Ala	Met	Glu	Gly													
		115														

&lt;210&gt; 52

&lt;211&gt; 116

&lt;212&gt; PRT

&lt;213&gt; Artificial Sequence

<223> Description of Artificial Sequence: Murine Il-5  
modified by substitution with tetanus toxoid  
epitope

&lt;400&gt; 52

Met	Glu	Ile	Pro	Met	Ser	Thr	Val	Val	Lys	Glu	Thr	Leu	Thr	Gln	Leu	
1				5					10					15		

Ser	Ala	His	Arg	Ala	Leu	Leu	Thr	Ser	Asn	Glu	Thr	Met	Gln	Tyr	Ile	
			20					25					30			

Lys	Ala	Asn	Ser	Lys	Phe	Ile	Gly	Ile	Thr	Glu	Leu	Cys	Ile	Gly	Glu	
		35					40					45				

Ile	Phe	Gln	Gly	Leu	Asp	Ile	Leu	Lys	Asn	Gln	Thr	Val	Arg	Gly	Gly	
	50					55					60					

Thr	Val	Glu	Met	Leu	Phe	Gln	Asn	Leu	Ser	Leu	Ile	Lys	Lys	Tyr	Ile	
	65				70					75					80	

05980916 031300

Asp Arg Gln Lys Glu Lys Cys Gly Glu Glu Arg Arg Arg Thr Arg Gln  
                             85                            90                            95

Phe Leu Asp Tyr Leu Gln Glu Phe Leu Gly Val Met Ser Thr Glu Trp  
                             100                            105                            110

Ala Met Glu Gly  
                             115

<210> 53  
 <211> 369  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: Murine Il-5  
           modified by substitution with tetanus toxoid  
           epitope

<220>  
 <221> CDS  
 <222> (1)..(369)

<220>  
 <221> mutation  
 <222> (250)..(294)  
 <223> Tetanus toxoid P2 epitope

<220>  
 <221> misc\_feature  
 <222> (1)..(249)  
 <223> DNA encoding amino acids 1-83 of murine IL5

<220>  
 <221> misc\_feature  
 <222> (295)..(366)  
 <223> DNA encoding amino acids 90-113 of murine IL5

<400> 53  
 atg gag att ccc atg agc aca gtg gtg aaa gag acc ttg aca cag ctg 48  
 Met Glu Ile Pro Met Ser Thr Val Val Lys Glu Thr Leu Thr Gln Leu  
   1                            5                            10                            15

tcc gct cac cga gct ctg ttg aca agc aat gag acg atg agg ctt cct 96  
 Ser Ala His Arg Ala Leu Leu Thr Ser Asn Glu Thr Met Arg Leu Pro  
                             20                            25                            30

gtc cct act cat aaa aat cac cag cta tgc att gga gag atc ttt cag 144  
 Val Pro Thr His Lys Asn His Gln Leu Cys Ile Gly Glu Ile Phe Gln  
                             35                            40                            45

ggg cta gac ata ctg aag aat caa act gtc cgt ggg ggt acc gtg gaa 192  
 Gly Leu Asp Ile Leu Lys Asn Gln Thr Val Arg Gly Gly Thr Val Glu  
                             50                            55                            60

atg cta ttc caa aac ctg tca tta ata aag aaa tac atc gat aga caa 240  
 Met Leu Phe Gln Asn Leu Ser Leu Ile Lys Lys Tyr Ile Asp Arg Gln  
   65                            70                            75                            80

09980916 "021902"

44

```

aaa gag aag cag tac atc aag gcc aac tcc aag ttc atc ggc atc acc 288
Lys Glu Lys Gln Tyr Ile Lys Ala Asn Ser Lys Phe Ile Gly Ile Thr
      85                      90                      95

gag ctg agg acg agg cag ttc ctg gat tat ctg cag gag ttc ctt ggt 336
Glu Leu Arg Thr Arg Gln Phe Leu Asp Tyr Leu Gln Glu Phe Leu Gly
      100                      105                      110

gtg atg agt aca gag tgg gca atg gaa ggc taa 369
Val Met Ser Thr Glu Trp Ala Met Glu Gly
      115                      120

```

<210> 54  
 <211> 122  
 <212> PRT  
 <213> Artificial Sequence  
 <223> Description of Artificial Sequence: Murine Il-5  
 modified by substitution with tetanus toxoid

epitope

```

<400> 54
Met Glu Ile Pro Met Ser Thr Val Val Lys Glu Thr Leu Thr Gln Leu
 1              5              10              15

Ser Ala His Arg Ala Leu Leu Thr Ser Asn Glu Thr Met Arg Leu Pro
      20              25              30

Val Pro Thr His Lys Asn His Gln Leu Cys Ile Gly Glu Ile Phe Gln
      35              40              45

Gly Leu Asp Ile Leu Lys Asn Gln Thr Val Arg Gly Gly Thr Val Glu
      50              55              60

Met Leu Phe Gln Asn Leu Ser Leu Ile Lys Lys Tyr Ile Asp Arg Gln
      65              70              75              80

Lys Glu Lys Gln Tyr Ile Lys Ala Asn Ser Lys Phe Ile Gly Ile Thr
      85              90              95

Glu Leu Arg Thr Arg Gln Phe Leu Asp Tyr Leu Gln Glu Phe Leu Gly
      100              105              110

Val Met Ser Thr Glu Trp Ala Met Glu Gly
      115              120

```

<210> 55  
 <211> 393  
 <212> DNA  
 <213> Artificial Sequence  
  
 <220>  
 <223> Description of Artificial Sequence: Murine Il-5  
 modified by substitution with tetanus toxoid  
 epitope  
  
 <220>

2005-12-01 10:00:00

<221> CDS  
<222> (1)..(393)

<220>  
<221> mutation  
<222> (256)..(318)  
<223> Tetanus toxoid P30 epitope

<220>  
<221> misc\_feature  
<222> (1)..(255)  
<223> DNA encoding amino acids 1-85 of murine IL5

<220>  
<221> misc\_feature  
<222> (319)..(390)  
<223> DNA encoding amino acids 90-113 of murine IL5

<400> 55  
atg gag att ccc atg agc aca gtg gtg aaa gag acc ttg aca cag ctg 48  
Met Glu Ile Pro Met Ser Thr Val Val Lys Glu Thr Leu Thr Gln Leu  
1 5 10 15  
  
tcc gct cac cga gct ctg ttg aca agc aat gag acg atg agg ctt cct 96  
Ser Ala His Arg Ala Leu Leu Thr Ser Asn Glu Thr Met Arg Leu Pro  
20 25 30  
  
gtc cct act cat aaa aat cac cag cta tgc att gga gag atc ttt cag 144  
Val Pro Thr His Lys Asn His Gln Leu Cys Ile Gly Glu Ile Phe Gln  
35 40 45  
  
ggg cta gac ata ctg aag aat caa act gtc cgt ggg ggt acc gtg gaa 192  
Gly Leu Asp Ile Leu Lys Asn Gln Thr Val Arg Gly Gly Thr Val Glu  
50 55 60  
  
atg cta ttc caa aac ctg tca tta ata aag aaa tac atc gat aga caa 240  
Met Leu Phe Gln Asn Leu Ser Leu Ile Lys Lys Tyr Ile Asp Arg Gln  
65 70 75 80  
  
aaa gag aag tgt ggc ttc aac aac ttc acc gtg agc ttc tgg ctg cgc 288  
Lys Glu Lys Cys Gly Phe Asn Asn Phe Thr Val Ser Phe Trp Leu Arg  
85 90 95  
  
gtg ccc aag gtg agc gcc agc cac ctg gag agg acg agg cag ttc ctg 336  
Val Pro Lys Val Ser Ala Ser His Leu Glu Arg Thr Arg Gln Phe Leu  
100 105 110  
  
gat tat ctg cag gag ttc ctt ggt gtg atg agt aca gag tgg gca atg 384  
Asp Tyr Leu Gln Glu Phe Leu Gly Val Met Ser Thr Glu Trp Ala Met  
115 120 125  
  
gaa ggc taa 393  
Glu Gly  
130

<210> 56  
<211> 130  
<212> PRT  
<213> Artificial Sequence  
<223> Description of Artificial Sequence: Murine Il-5

0900946 02192

46

modified by substitution with tetanus toxoid  
epitope

&lt;400&gt; 56

Met Glu Ile Pro Met Ser Thr Val Val Lys Glu Thr Leu Thr Gln Leu  
1 5 10 15

Ser Ala His Arg Ala Leu Leu Thr Ser Asn Glu Thr Met Arg Leu Pro  
20 25 30

Val Pro Thr His Lys Asn His Gln Leu Cys Ile Gly Glu Ile Phe Gln  
35 40 45

Gly Leu Asp Ile Leu Lys Asn Gln Thr Val Arg Gly Gly Thr Val Glu  
50 55 60

Met Leu Phe Gln Asn Leu Ser Leu Ile Lys Lys Tyr Ile Asp Arg Gln  
65 70 75 80

Lys Glu Lys Cys Gly Phe Asn Asn Phe Thr Val Ser Phe Trp Leu Arg  
85 90 95

Val Pro Lys Val Ser Ala Ser His Leu Glu Arg Thr Arg Gln Phe Leu  
100 105 110

Asp Tyr Leu Gln Glu Phe Leu Gly Val Met Ser Thr Glu Trp Ala Met  
115 120 125

Glu Gly  
130

&lt;210&gt; 57

&lt;211&gt; 387

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

<223> Description of Artificial Sequence: Murine IL-5  
modified by substitution with tetanus toxoid  
epitope

&lt;220&gt;

&lt;221&gt; CDS

&lt;222&gt; (1)..(387)

&lt;220&gt;

&lt;221&gt; mutation

&lt;222&gt; (250)..(312)

&lt;223&gt; Tetanus toxoid P30 epitope

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;222&gt; (1)..(249)

&lt;223&gt; DNA encoding amino acids 1-83 of murine IL5

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;222&gt; (313)..(384)

&lt;223&gt; DNA encoding amino acids 90-113 of murine IL5

09000916 001490

<400> 57  
 atg gag att ccc atg agc aca gtg gtg aaa gag acc ttg aca cag ctg 48  
 Met Glu Ile Pro Met Ser Thr Val Val Lys Glu Thr Leu Thr Gln Leu  
 1 5 10 15

tcc gct cac cga gct ctg ttg aca agc aat gag acg atg agg ctt cct 96  
 Ser Ala His Arg Ala Leu Leu Thr Ser Asn Glu Thr Met Arg Leu Pro  
 20 25 30

gtc cct act cat aaa aat cac cag cta tgc att gga gag atc ttt cag 144  
 Val Pro Thr His Lys Asn His Gln Leu Cys Ile Gly Glu Ile Phe Gln  
 35 40 45

ggg cta gac ata ctg aag aat caa act gtc cgt ggg ggt acc gtg gaa 192  
 Gly Leu Asp Ile Leu Lys Asn Gln Thr Val Arg Gly Gly Thr Val Glu  
 50 55 60

atg cta ttc caa aac ctg tca tta ata aag aaa tac atc gat aga caa 240  
 Met Leu Phe Gln Asn Leu Ser Leu Ile Lys Lys Tyr Ile Asp Arg Gln  
 65 70 75 80

aaa gag aag ttc aac aac ttc acc gtg agc ttc tgg ctg cgc gtg ccc 288  
 Lys Glu Lys Phe Asn Asn Phe Thr Val Ser Phe Trp Leu Arg Val Pro  
 85 90 95

aag gtg agc gcc agc cac ctg gag agg acg agg cag ttc ctg gat tat 336  
 Lys Val Ser Ala Ser His Leu Glu Arg Thr Arg Gln Phe Leu Asp Tyr  
 100 105 110

ctg cag gag ttc ctt ggt gtg atg agt aca gag tgg gca atg gaa ggc 384  
 Leu Gln Glu Phe Leu Gly Val Met Ser Thr Glu Trp Ala Met Glu Gly  
 115 120 125

taa 387

<210> 58  
 <211> 128  
 <212> PRT  
 <213> Artificial Sequence  
 <223> Description of Artificial Sequence: Murine Il-5  
 modified by substitution with tetanus toxoid  
 epitope

<400> 58  
 Met Glu Ile Pro Met Ser Thr Val Val Lys Glu Thr Leu Thr Gln Leu  
 1 5 10 15

Ser Ala His Arg Ala Leu Leu Thr Ser Asn Glu Thr Met Arg Leu Pro  
 20 25 30

Val Pro Thr His Lys Asn His Gln Leu Cys Ile Gly Glu Ile Phe Gln  
 35 40 45

Gly Leu Asp Ile Leu Lys Asn Gln Thr Val Arg Gly Gly Thr Val Glu  
 50 55 60

Met Leu Phe Gln Asn Leu Ser Leu Ile Lys Lys Tyr Ile Asp Arg Gln  
 65 70 75 80

0065058 WO 00/65058

Lys Glu Lys Phe Asn Asn Phe Thr Val Ser Phe Trp Leu Arg Val Pro  
85 90 95

Lys Val Ser Ala Ser His Leu Glu Arg Thr Arg Gln Phe Leu Asp Tyr  
100 105 110

Leu Gln Glu Phe Leu Gly Val Met Ser Thr Glu Trp Ala Met Glu Gly  
115 120 125

<210> 59

<211> 438

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Murine Il-5  
modified by substitution with tetanus toxoid  
epitopes

<220>

<221> CDS

<222> (1)..(438)

<220>

<221> mutation

<222> (256)..(300)

<223> Tetanus toxoid P2 epitope

<220>

<221> mutation

<222> (301)..(363)

<223> Tetanus toxoid P30 epitope

<220>

<221> misc\_feature

<222> (1)..(255)

<223> DNA encoding amino acids 1-85 of murine IL5

<220>

<221> misc\_feature

<222> (364)..(435)

<223> DNA encoding amino acids 90-113 of murine IL5

<400> 59

atg gag att ccc atg agc aca gtg gtg aaa gag acc ttg aca cag ctg 48  
Met Glu Ile Pro Met Ser Thr Val Val Lys Glu Thr Leu Thr Gln Leu  
1 5 10 15

tcc gct cac cga gct ctg ttg aca agc aat gag acg atg agg ctt cct 96  
Ser Ala His Arg Ala Leu Leu Thr Ser Asn Glu Thr Met Arg Leu Pro  
20 25 30

gtc cct act cat aaa aat cac cag cta tgc att gga gag atc ttt cag 144  
Val Pro Thr His Lys Asn His Gln Leu Cys Ile Gly Glu Ile Phe Gln  
35 40 45

0998094 0045720 9T608660



ggg cta gac ata ctg aag aat caa act gtc cgt ggg ggt acc gtg gaa 192  
Gly Leu Asp Ile Leu Lys Asn Gln Thr Val Arg Gly Gly Thr Val Glu  
50 55 60

atg cta ttc caa aac ctg tca tta ata aag aaa tac atc gat aga caa 240  
Met Leu Phe Gln Asn Leu Ser Leu Ile Lys Lys Tyr Ile Asp Arg Gln  
65 70 75 80

aaa gag aag tgt ggc cag tac atc aag gcc aac tcc aag ttc atc ggc 288  
Lys Glu Lys Cys Gly Gln Tyr Ile Lys Ala Asn Ser Lys Phe Ile Gly  
85 90 95

atc acc gag ctg ttc aac aac ttc acc gtg agc ttc tgg ctg cgc gtg 336  
Ile Thr Glu Leu Phe Asn Asn Phe Thr Val Ser Phe Trp Leu Arg Val  
100 105 110

ccc aag gtg agc gcc agc cac ctg gag agg acg agg cag ttc ctg gat 384  
Pro Lys Val Ser Ala Ser His Leu Glu Arg Thr Arg Gln Phe Leu Asp  
115 120 125

tat ctg cag gag ttc ctt ggt gtg atg agt aca gag tgg gca atg gaa 432  
Tyr Leu Gln Glu Phe Leu Gly Val Met Ser Thr Glu Trp Ala Met Glu  
130 135 140

ggc taa 438  
Gly  
145

<210> 60

<211> 145

<212> PRT

<213> Artificial Sequence

<223> Description of Artificial Sequence: Murine Il-5  
modified by substitution with tetanus toxoid  
epitopes

<400> 60

Met Glu Ile Pro Met Ser Thr Val Val Lys Glu Thr Leu Thr Gln Leu  
1 5 10 15

Ser Ala His Arg Ala Leu Leu Thr Ser Asn Glu Thr Met Arg Leu Pro  
20 25 30

Val Pro Thr His Lys Asn His Gln Leu Cys Ile Gly Glu Ile Phe Gln  
35 40 45

Gly Leu Asp Ile Leu Lys Asn Gln Thr Val Arg Gly Gly Thr Val Glu  
50 55 60

Met Leu Phe Gln Asn Leu Ser Leu Ile Lys Lys Tyr Ile Asp Arg Gln  
65 70 75 80

Lys Glu Lys Cys Gly Gln Tyr Ile Lys Ala Asn Ser Lys Phe Ile Gly  
85 90 95

Ile Thr Glu Leu Phe Asn Asn Phe Thr Val Ser Phe Trp Leu Arg Val  
100 105 110

Pro Lys Val Ser Ala Ser His Leu Glu Arg Thr Arg Gln Phe Leu Asp  
115 120 125

Socioeconomic characteristics	
Age (years)	18-24
Gender	Male
Marital status	Married
Education (years)	12-15
Occupation	Unemployed
Income (USD/month)	100-150
Health insurance	Private
Smoking status	Smoker
Alcohol consumption	Regular
Exercise frequency	Low
Dietary habits	High fat
Stress level	High
Family size	Large
Urban/rural	Urban
Religion	Christian
Political affiliation	Conservative
Travel history	International
Comorbidities	Hypertension
Medication use	Antidepressants
Substance use	Heroin
Legal status	Arrested
Employment status	Unemployed
Home ownership	Rent
Neighborhood safety	Low
Access to healthcare	Limited
Health literacy	Low
Health beliefs	Fatalistic
Health behaviors	Risky
Health status	Chronic
Healthcare utilization	Low
Healthcare costs	High
Healthcare access	Barriers
Healthcare quality	Poor
Healthcare satisfaction	Low
Healthcare trust	Low
Healthcare engagement	Low
Healthcare compliance	Low
Healthcare adherence	Low
Healthcare utilization	Low
Healthcare costs	High
Healthcare access	Barriers
Healthcare quality	Poor
Healthcare satisfaction	Low
Healthcare trust	Low
Healthcare engagement	Low
Healthcare compliance	Low
Healthcare adherence	Low
Healthcare utilization	Low
Healthcare costs	High
Healthcare access	Barriers
Healthcare quality	Poor
Healthcare satisfaction	Low
Healthcare trust	Low
Healthcare engagement	Low
Healthcare compliance	Low
Healthcare adherence	Low
Healthcare utilization	Low
Healthcare costs	High
Healthcare access	Barriers
Healthcare quality	Poor
Healthcare satisfaction	Low
Healthcare trust	Low
Healthcare engagement	Low
Healthcare compliance	Low
Healthcare adherence	Low
Healthcare utilization	Low
Healthcare costs	High
Healthcare access	Barriers
Healthcare quality	Poor
Healthcare satisfaction	Low
Healthcare trust	Low
Healthcare engagement	Low
Healthcare compliance	Low
Healthcare adherence	Low
Healthcare utilization	Low
Healthcare costs	High
Healthcare access	Barriers
Healthcare quality	Poor
Healthcare satisfaction	Low
Healthcare trust	Low
Healthcare engagement	Low
Healthcare compliance	Low
Healthcare adherence	Low
Healthcare utilization	Low
Healthcare costs	High
Healthcare access	Barriers
Healthcare quality	Poor
Healthcare satisfaction	Low
Healthcare trust	Low
Healthcare engagement	Low
Healthcare compliance	Low
Healthcare adherence	Low
Healthcare utilization	Low
Healthcare costs	High
Healthcare access	Barriers
Healthcare quality	Poor
Healthcare satisfaction	Low
Healthcare trust	Low
Healthcare engagement	Low
Healthcare compliance	Low
Healthcare adherence	Low
Healthcare utilization	Low
Healthcare costs	High
Healthcare access	Barriers
Healthcare quality	Poor
Healthcare satisfaction	Low
Healthcare trust	Low
Healthcare engagement	Low
Healthcare compliance	Low
Healthcare adherence	Low
Healthcare utilization	Low
Healthcare costs	High
Healthcare access	Barriers
Healthcare quality	Poor
Healthcare satisfaction	Low
Healthcare trust	Low
Healthcare engagement	Low
Healthcare compliance	Low
Healthcare adherence	Low
Healthcare utilization	Low
Healthcare costs	High
Healthcare access	Barriers
Healthcare quality	Poor
Healthcare satisfaction	Low
Healthcare trust	Low
Healthcare engagement	Low
Healthcare compliance	Low
Healthcare adherence	Low
Healthcare utilization	Low
Healthcare costs	High
Healthcare access	Barriers
Healthcare quality	Poor
Healthcare satisfaction	Low
Healthcare trust	Low
Healthcare engagement	Low
Healthcare compliance	Low
Healthcare adherence	Low
Healthcare utilization	Low
Healthcare costs	High
Healthcare access	Barriers
Healthcare quality	Poor
Healthcare satisfaction	Low
Healthcare trust	Low
Healthcare engagement	Low
Healthcare compliance	Low
Healthcare adherence	Low
Healthcare utilization	Low
Healthcare costs	High
Healthcare access	Barriers
Healthcare quality	Poor
Healthcare satisfaction	Low
Healthcare trust	Low
Healthcare engagement	Low
Healthcare compliance	Low
Healthcare adherence	Low
Healthcare utilization	Low
Healthcare costs	High
Healthcare access	Barriers
Healthcare quality	Poor
Healthcare satisfaction	Low
Healthcare trust	Low
Healthcare engagement	Low
Healthcare compliance	Low
Healthcare adherence	Low
Healthcare utilization	Low
Healthcare costs	High
Healthcare access	Barriers
Healthcare quality	Poor
Healthcare satisfaction	Low
Healthcare trust	Low
Healthcare engagement	Low
Healthcare compliance	Low
Healthcare adherence	Low
Healthcare utilization	Low
Healthcare costs	High

Tyr Leu Gln Glu Phe Leu Gly Val Met Ser Thr Glu Trp Ala Met Glu  
130 135 140

Gly  
145

```
<210> 61
<211> 57
<212> DNA
<213> Homo sapiens
```

```
<220>  
<221> CDS  
<222> (1)..(57)  
<223> DNA encoding natural human IL5 leader sequence
```

```

<400> 61
atg agg atg ctt ctg cat ttg agt ttg ctg gct ctt gga gct gcc tac 48
Met Arg Met Leu Leu His Leu Ser Leu Leu Ala Leu Gly Ala Ala Tyr
      1              5              10              15

gtg tat gcc 57
Val Tyr Ala

```

```
<210> 62
<211> 19
<212> PRT
<213> Homo sapiens
```

```
<400> 62
Met Arg Met Leu Leu His Leu Ser Leu Leu Ala Leu Gly Ala Ala Tyr
  1                      5                      10                      15
Val Tyr Ala
```

```
<210> 63
<211> 60
<212> DNA
<213> Mus musculus
```

```
<220>
<221> CDS
<222> {1}..(60)
<223> DNA encoding natural murine IL5 leader sequence
```

```

<400> 63
atg aga agg atg ctt ctg cac ttg agt gtt ctg act ctc agc tgt gtc 48
Met Arg Arg Met Leu Leu His Leu Ser Val Leu Thr Leu Ser Cys Val
      1              5              10              15

tgg gcc act gcc
Trp Ala Thr Ala
      20

```

```
<210> 64
<211> 19
<212> PRT
<213> Mus musculus
```

```

<400> 64
Met Arg Arg Met Leu Leu His Leu Ser Val Leu Thr Leu Ser Cys Val
  1                               5                10                15
Trp Ala Thr Ala
                20

```

```
<210> 65
<211> 13
<212> PRT
<213> Artificial Sequence
```

```
<220>
<223> Description of Artificial Sequence: Promiscuous T
        helper epitope
```

<400> 65  
Ala Lys Phe Val Ala Ala Trp Thr Leu Lys Ala Ala Ala  
1 5 10

SEQUENCE LISTING

<110> M&E Biotech A/S  
Klysner, Steen

<120> Method For Down-Regulating IL5 Activity

<130> 3631-0112P

<140>

<141>

<160> 65

<170> PatentIn Ver. 2.1

<210> 1

<211> 115

<212> PRT

<213> Homo sapiens

<220>

<221> DISULFID

<222> (44)

<223> Interchain disulphide bond to Cys-86 in SEQ ID  
NO:1

<220>

<221> DISULFID

<222> (86)

<223> Interchain disulphide bond to Cys-44 in SEQ ID  
NO:1

<400> 1

Ile Pro Thr Glu Ile Pro Thr Ser Ala Leu Val Lys Glu Thr Leu Ala  
1 5 10 15

Leu Leu Ser Thr His Arg Thr Leu Leu Ile Ala Asn Glu Thr Leu Arg  
20 25 30

Ile Pro Val Pro Val His Lys Asn His Gln Leu Cys Thr Glu Glu Ile  
35 40 45

Phe Gln Gly Ile Gly Thr Leu Glu Ser Gln Thr Val Gln Gly Gly Thr  
50 55 60

Val Glu Arg Leu Phe Lys Asn Leu Ser Leu Ile Lys Lys Tyr Ile Asp  
65 70 75 80

Gly Gln Lys Lys Lys Cys Gly Glu Glu Arg Arg Arg Val Asn Gln Phe  
85 90 95

Leu Asp Tyr Leu Gln Glu Phe Leu Gly Val Met Asn Thr Glu Trp Ile  
100 105 110

Ile Glu Ser  
115

09980916.021902

<210> 2  
 <211> 126  
 <212> PRT  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: Human IL5  
 modified by substitution with tetanus toxoid P2  
 epitope

<220>  
 <221> MUTAGEN  
 <222> (32)..(46)  
 <223> Tetanus toxoid P2 epitope (SEQ ID NO: 23)

<220>  
 <221> SIMILAR  
 <222> (1)..(86)  
 <223> Identical to residues 1-86 in SEQ ID NO: 1

<220>  
 <221> SIMILAR  
 <222> (102)..(126)  
 <223> Identical to residues 91-115 in SEQ ID NO: 1

<400> 2  
 Ile Pro Thr Glu Ile Pro Thr Ser Ala Leu Val Lys Glu Thr Leu Ala  
           1                  5                  10                  15  
 Leu Leu Ser Thr His Arg Thr Leu Leu Ile Ala Asn Glu Thr Leu Arg  
                   20                  25                  30  
 Ile Pro Val Pro Val His Lys Asn His Gln Leu Cys Thr Glu Glu Ile  
                   35                  40                  45  
 Phe Gln Gly Ile Gly Thr Leu Glu Ser Gln Thr Val Gln Gly Gly Thr  
           50                  55                  60  
 Val Glu Arg Leu Phe Lys Asn Leu Ser Leu Ile Lys Lys Tyr Ile Asp  
           65                  70                  75                  80  
 Gly Gln Lys Lys Lys Cys Gln Tyr Ile Lys Ala Asn Ser Lys Phe Ile  
                   85                  90                  95  
 Gly Ile Thr Glu Leu Arg Arg Val Asn Gln Phe Leu Asp Tyr Leu Gln  
           100                  105                  110  
 Glu Phe Leu Gly Val Met Asn Thr Glu Trp Ile Ile Glu Ser  
           115                  120                  125

<210> 3  
 <211> 118

20050609 16:05:56

<212> PRT  
<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:Human IL5  
modified by substitution with tetanus toxoid P2  
epitope

<220>

<221> MUTAGEN

<222> (32)..(46)

<223> Tetanus toxoid P2 epitope (SEQ ID NO: 23)

<220>

<221> SIMILAR

<222> (1)..(31)

<223> Identical to residues 1-31 in SEQ ID NO: 1

<220>

<221> SIMILAR

<222> (47)..(118)

<223> Identical to residues 44-115 in SEQ ID NO: 1

<400> 3

Ile Pro Thr Glu Ile Pro Thr Ser Ala Leu Val Lys Glu Thr Leu Ala  
1 5 10 15

Leu Leu Ser Thr His Arg Thr Leu Leu Ile Ala Asn Glu Thr Leu Gln  
20 25 30

Tyr Ile Lys Ala Asn Ser Lys Phe Ile Gly Ile Thr Glu Leu Cys Thr  
35 40 45

Glu Glu Ile Phe Gln Gly Ile Gly Thr Leu Glu Ser Gln Thr Val Gln  
50 55 60

Gly Gly Thr Val Glu Arg Leu Phe Lys Asn Leu Ser Leu Ile Lys Lys  
65 70 75 80

Tyr Ile Asp Gly Gln Lys Lys Lys Cys Gly Glu Glu Arg Arg Arg Val  
85 90 95

Asn Gln Phe Leu Asp Tyr Leu Gln Glu Phe Leu Gly Val Met Asn Thr  
100 105 110

Glu Trp Ile Ile Glu Ser  
115

<210> 4

<211> 124

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:Human IL5

03980916 "021902

modified by substitution with tetanus toxoid P2  
epitope

<220>  
<221> MUTAGEN  
<222> (59)..(73)  
<223> Tetanus toxoid P2 epitope (SEQ ID NO:23)

<220>  
<221> SIMILAR  
<222> (1)..(58)  
<223> Identical to residues 1-58 in SEQ ID NO: 1

<220>  
<221> SIMILAR  
<222> (74)..(124)  
<223> Identical to residues 65-115 in SEQ ID NO: 1

<400> 4  
Ile Pro Thr Glu Ile Pro Thr Ser Ala Leu Val Lys Glu Thr Leu Ala  
1 5 10 15  
Leu Leu Ser Thr His Arg Thr Leu Leu Ile Ala Asn Glu Thr Leu Arg  
20 25 30  
Ile Pro Val Pro Val His Lys Asn His Gln Leu Cys Thr Glu Glu Ile  
35 40 45  
Phe Gln Gly Ile Gly Thr Leu Glu Ser Gln Gln Tyr Ile Lys Ala Asn  
50 55 60  
Ser Lys Phe Ile Gly Ile Thr Glu Leu Val Glu Arg Leu Phe Lys Asn  
65 70 75 80  
Leu Ser Leu Ile Lys Lys Tyr Ile Asp Gly Gln Lys Lys Lys Cys Gly  
85 90 95  
Glu Glu Arg Arg Arg Val Asn Gln Phe Leu Asp Tyr Leu Gln Glu Phe  
100 105 110  
Leu Gly Val Met Asn Thr Glu Trp Ile Ile Glu Ser  
115 120

<210> 5  
<211> 124  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence:Human IL5  
modified by substitution with tetanus toxoid P2  
epitope

<220>  
<221> MUTAGEN

00000916-021902

<222> (86)..(100)

<223> Tetanus toxoid P2 epitope (SEQ ID NO: 23)

<220>

<221> SIMILAR

<222> (1)..(85)

<223> Identical to residues 1-85 in SEQ ID NO: 1

<220>

<221> SIMILAR

<222> (101)..(124)

<223> Identical to residues 90-115 in SEQ ID NO: 1

<400> 5

Ile Pro Thr Glu Ile Pro Thr Ser Ala Leu Val Lys Glu Thr Leu Ala  
1 5 10 15

Leu Leu Ser Thr His Arg Thr Leu Leu Ile Ala Asn Glu Thr Leu Arg  
20 25 30

Ile Pro Val Pro Val His Lys Asn His Gln Leu Cys Thr Glu Glu Ile  
35 40 45

Phe Gln Gly Ile Gly Thr Leu Glu Ser Gln Thr Val Gln Gly Gly Thr  
50 55 60

Val Glu Arg Leu Phe Lys Asn Leu Ser Leu Ile Lys Lys Tyr Ile Asp  
65 70 75 80

Gly Gln Lys Lys Lys Gln Tyr Ile Lys Ala Asn Ser Lys Phe Ile Gly  
85 90 95

Ile Thr Glu Leu Arg Val Asn Gln Phe Leu Asp Tyr Leu Gln Glu Phe  
100 105 110

Leu Gly Val Met Asn Thr Glu Trp Ile Ile Glu Ser  
115 120

<210> 6

<211> 126

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Human IL5  
modified by substitution with tetanus toxoid P2  
epitope

<220>

<221> MUTAGEN

<222> (110)..(124)

<223> Tetanus toxoid P2 epitope (SEQ ID NO: 23)

<220>

<221> SIMILAR

00990916-021902



<222> (1)..(109)

<223> Identical to residues 1-109 in SEQ ID NO: 1

<220>

<221> SIMILAR

<222> (125)..(126)

<223> Identical to residues 114-115 in SEQ ID NO: 1

<400> 6

Ile Pro Thr Glu Ile Pro Thr Ser Ala Leu Val Lys Glu Thr Leu Ala  
1 5 10 15

Leu Leu Ser Thr His Arg Thr Leu Leu Ile Ala Asn Glu Thr Leu Arg  
20 25 30

Ile Pro Val Pro Val His Lys Asn His Gln Leu Cys Thr Glu Glu Ile  
35 40 45

Phe Gln Gly Ile Gly Thr Leu Glu Ser Gln Thr Val Gln Gly Gly Thr  
50 55 60

Val Glu Arg Leu Phe Lys Asn Leu Ser Leu Ile Lys Lys Tyr Ile Asp  
65 70 75 80

Gly Gln Lys Lys Lys Cys Gly Glu Glu Arg Arg Arg Val Asn Gln Phe  
85 90 95

Leu Asp Tyr Leu Gln Glu Phe Leu Gly Val Met Asn Thr Gln Tyr Ile  
100 105 110

Lys Ala Asn Ser Lys Phe Ile Gly Ile Thr Glu Leu Glu Ser  
115 120 125

<210> 7

<211> 132

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:Human IL5  
modified by substitution with tetanus toxoid P30  
epitope

<220>

<221> MUTAGEN

<222> (87)..(107)

<223> Tetanus toxoid P30 epitope (SEQ ID NO: 24)

<220>

<221> SIMILAR

<222> (1)..(86)

<223> Identical to residues 1-86 in SEQ ID NO: 1

<220>

<221> SIMILAR

00980916 "02100"

<222> (108)..(132)

<223> Identical to residues 91-115 in SEQ ID NO: 1

<400> 7

Ile Pro Thr Glu Ile Pro Thr Ser Ala Leu Val Lys Glu Thr Leu Ala  
1 5 10 15

Leu Leu Ser Thr His Arg Thr Leu Leu Ile Ala Asn Glu Thr Leu Arg  
20 25 30

Ile Pro Val Pro Val His Lys Asn His Gln Leu Cys Thr Glu Glu Ile  
35 40 45

Phe Gln Gly Ile Gly Thr Leu Glu Ser Gln Thr Val Gln Gly Gly Thr  
50 55 60

Val Glu Arg Leu Phe Lys Asn Leu Ser Leu Ile Lys Lys Tyr Ile Asp  
65 70 75 80

Gly Gln Lys Lys Lys Cys Phe Asn Asn Phe Thr Val Ser Phe Trp Leu  
85 90 95

Arg Val Pro Lys Val Ser Ala Ser His Leu Glu Arg Arg Val Asn Gln  
100 105 110

Phe Leu Asp Tyr Leu Gln Glu Phe Leu Gly Val Met Asn Thr Glu Trp  
115 120 125

Ile Ile Glu Ser  
130

<210> 8

<211> 124

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Human IL5  
modified by substitution with tetanus toxoid P30  
epitope

<220>

<221> MUTAGEN

<222> (32)..(52)

<223> Tetanus toxoid P30 epitope (SEQ ID NO: 24)

<220>

<221> SIMILAR

<222> (1)..(31)

<223> Identical to residues 1-31 in SEQ ID NO: 1

<220>

<221> SIMILAR

<222> (53)..(124)

<223> Identical to residues 44-115 in SEQ ID NO: 1

205720-9750550

&lt;400&gt; 8

Ile Pro Thr Glu Ile Pro Thr Ser Ala Leu Val Lys Glu Thr Leu Ala  
 1 5 10 15

Leu Leu Ser Thr His Arg Thr Leu Leu Ile Ala Asn Glu Thr Leu Phe  
 20 25 30

Asn Asn Phe Thr Val Ser Phe Trp Leu Arg Val Pro Lys Val Ser Ala  
 35 40 45

Ser His Leu Glu Cys Thr Glu Glu Ile Phe Gln Gly Ile Gly Thr Leu  
 50 55 60

Glu Ser Gln Thr Val Gln Gly Gly Thr Val Glu Arg Leu Phe Lys Asn  
 65 70 75 80

Leu Ser Leu Ile Lys Lys Tyr Ile Asp Gly Gln Lys Lys Lys Cys Gly  
 85 90 95

Glu Glu Arg Arg Arg Val Asn Gln Phe Leu Asp Tyr Leu Gln Glu Phe  
 100 105 110

Leu Gly Val Met Asn Thr Glu Trp Ile Ile Glu Ser  
 115 120

&lt;210&gt; 9

&lt;211&gt; 130

&lt;212&gt; PRT

&lt;213&gt; Artificial Sequence

&lt;220&gt;

<223> Description of Artificial Sequence:Human IL5  
 modified by substitution with tetanus toxoid P30  
 epitope

&lt;220&gt;

&lt;221&gt; MUTAGEN

&lt;222&gt; (59)..(79)

&lt;223&gt; Tetanus toxoid P30 epitope (SEQ ID NO: 24)

&lt;220&gt;

&lt;221&gt; SIMILAR

&lt;222&gt; (1)..(58)

&lt;223&gt; Identical to residues 1-58 in SEQ ID NO: 1

&lt;220&gt;

&lt;221&gt; SIMILAR

&lt;222&gt; (80)..(130)

&lt;223&gt; Identical to residues 65-115 in SEQ ID NO: 1

&lt;400&gt; 9

Ile Pro Thr Glu Ile Pro Thr Ser Ala Leu Val Lys Glu Thr Leu Ala  
 1 5 10 15

2005120"9T608550

Leu Leu Ser Thr His Arg Thr Leu Leu Ile Ala Asn Glu Thr Leu Arg  
                   20                  25                  30  
 Ile Pro Val Pro Val His Lys Asn His Gln Leu Cys Thr Glu Glu Ile  
                   35                  40                  45  
 Phe Gln Gly Ile Gly Thr Leu Glu Ser Gln Phe Asn Asn Phe Thr Val  
                   50                  55                  60  
 Ser Phe Trp Leu Arg Val Pro Lys Val Ser Ala Ser His Leu Glu Val  
                   65                  70                  75                  80  
 Glu Arg Leu Phe Lys Asn Leu Ser Leu Ile Lys Lys Tyr Ile Asp Gly  
                   85                  90                  95  
 Gln Lys Lys Lys Cys Gly Glu Glu Arg Arg Arg Val Asn Gln Phe Leu  
                   100                  105                  110  
 Asp Tyr Leu Gln Glu Phe Leu Gly Val Met Asn Thr Glu Trp Ile Ile  
                   115                  120                  125  
 Glu Ser  
                   130

<210> 10  
 <211> 132  
 <212> PRT  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: Human IL5  
       modified by substitution with tetanus toxoid P30  
       epitope

<220>  
 <221> MUTAGEN  
 <222> (110)..(130)  
 <223> Tetanus toxoid P30 epitope (SEQ ID NO: 24)

<220>  
 <221> SIMILAR  
 <222> (1)..(129)  
 <223> Identical to residues 1-129 in SEQ ID NO: 1

<220>  
 <221> SIMILAR  
 <222> (131)..(132)  
 <223> Identical to residues 114-115 in SEQ ID NO: 1

<400> 10  
 Ile Pro Thr Glu Ile Pro Thr Ser Ala Leu Val Lys Glu Thr Leu Ala  
           1                  5                  10                  15

Leu Leu Ser Thr His Arg Thr Leu Leu Ile Ala Asn Glu Thr Leu Arg  
                   20                  25                  30

00550016.021502

Ile Pro Val Pro Val His Lys Asn His Gln Leu Cys Thr Glu Glu Ile  
                   35                  40                  45  
 Phe Gln Gly Ile Gly Thr Leu Glu Ser Gln Thr Val Gln Gly Gly Thr  
           50                  55                  60  
 Val Glu Arg Leu Phe Lys Asn Leu Ser Leu Ile Lys Lys Tyr Ile Asp  
   65                  70                  75                  80  
 Gly Gln Lys Lys Lys Cys Gly Glu Glu Arg Arg Arg Val Asn Gln Phe  
                   85                  90                  95  
 Leu Asp Tyr Leu Gln Glu Phe Leu Gly Val Met Asn Thr Phe Asn Asn  
           100                  105                  110  
 Phe Thr Val Ser Phe Trp Leu Arg Val Pro Lys Val Ser Ala Ser His  
           115                  120                  125  
 Leu Glu Glu Ser  
           130

<210> 11  
 <211> 141  
 <212> PRT  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: Human IL5  
       modified by substitution with tetanus toxoid P2  
       and P30 epitopes

<220>  
 <221> MUTAGEN  
 <222> (86)..(100)  
 <223> Tetanus toxoid P2 epitope (SEQ ID NO: 23)

<220>  
 <221> MUTAGEN  
 <222> (119)..(139)  
 <223> Tetanus toxoid P30 epitope (SEQ ID NO: 24)

<220>  
 <221> SIMILAR  
 <222> (1)..(85)  
 <223> Identical to residues 1-85 in SEQ ID NO: 1

<220>  
 <221> SIMILAR  
 <222> (101)..(118)  
 <223> Identical to residues 92-109 in SEQ ID NO: 1

<220>  
 <221> SIMILAR  
 <222> (140)..(141)

2025-10-16 09:55:50

<223> Identical to residues 114-115 in SEQ ID NO: 1

<400> 11

Ile Pro Thr Glu Ile Pro Thr Ser Ala Leu Val Lys Glu Thr Leu Ala  
1 5 10 15

Leu Leu Ser Thr His Arg Thr Leu Leu Ile Ala Asn Glu Thr Leu Arg  
20 25 30

Ile Pro Val Pro Val His Lys Asn His Gln Leu Cys Thr Glu Glu Ile  
35 40 45

Phe Gln Gly Ile Gly Thr Leu Glu Ser Gln Thr Val Gln Gly Gly Thr  
50 55 60

Val Glu Arg Leu Phe Lys Asn Leu Ser Leu Ile Lys Lys Tyr Ile Asp  
65 70 75 80

Gly Gln Lys Lys Lys Gln Tyr Ile Lys Ala Asn Ser Lys Phe Ile Gly  
85 90 95

Ile Thr Glu Leu Arg Val Asn Gln Phe Leu Asp Tyr Leu Gln Glu Phe  
100 105 110

Leu Gly Val Met Asn Thr Phe Asn Asn Phe Thr Val Ser Phe Trp Leu  
115 120 125

Arg Val Pro Lys Val Ser Ala Ser His Leu Glu Glu Ser  
130 135 140

<210> 12

<211> 113

<212> PRT

<213> Mus musculus

<220>

<221> DISULFID

<222> (42)

<223> Interchain disulphide bond to Cys-84 in SEQ ID  
NO:12

<220>

<221> DISULFID

<222> (84)

<223> Interchain disulphide bond to Cys-42 in SEQ ID  
NO:12

<400> 12

Met Glu Ile Pro Met Ser Thr Val Val Lys Glu Thr Leu Ala Leu Leu  
1 5 10 15

Ser Ala His Arg Ala Leu Leu Thr Ser Asn Glu Thr Met Arg Leu Pro  
20 25 30

Val Pro Thr His Lys Asn His Gln Leu Cys Ile Gly Glu Ile Phe Gln

03500916-021902

35                      40                      45  
 Gly Leu Asp Ile Leu Lys Asp Gln Thr Val Arg Gly Gly Thr Val Met  
     50                      55                      60  
 Arg Leu Phe Gln Asn Leu Ser Leu Ile Lys Lys Tyr Ile Asp Arg Gln  
     65                      70                      75                      80  
 Glu Lys Lys Cys Gly Glu Glu Arg Arg Arg Thr Arg Gln Phe Leu Asp  
                     85                      90                      95  
 Tyr Leu Gln Glu Phe Leu Gly Ser Met Asn Thr Ala Ala Ile Ile Glu  
                     100                      105                      110  
 Gly

<210> 13  
 <211> 124  
 <212> PRT  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: Murine IL5  
       modified by substitution with tetanus toxoid P2  
       epitope

<220>  
 <221> MUTAGEN  
 <222> (85)..(99)  
 <223> Tetanus toxoid P2 epitope (SEQ ID NO: 23)

<220>  
 <221> SIMILAR  
 <222> (1)..(84)  
 <223> Identical to residues 1-84 in SEQ ID NO: 12

<220>  
 <221> SIMILAR  
 <222> (100)..(124)  
 <223> Identical to residues 89-113 in SEQ ID NO: 12

<400> 13  
 Met Glu Ile Pro Met Ser Thr Val Val Lys Glu Thr Leu Ala Leu Leu  
     1                      5                      10                      15  
 Ser Ala His Arg Ala Leu Leu Thr Ser Asn Glu Thr Met Arg Leu Pro  
                     20                      25                      30  
 Val Pro Thr His Lys Asn His Gln Leu Cys Ile Gly Glu Ile Phe Gln  
                     35                      40                      45  
 Gly Leu Asp Ile Leu Lys Asp Gln Thr Val Arg Gly Gly Thr Val Met  
     50                      55                      60

"9T509550" 9T509550

Arg Leu Phe Gln Asn Leu Ser Leu Ile Lys Lys Tyr Ile Asp Arg Gln  
65 70 75 80

Glu Lys Lys Cys Gln Tyr Ile Lys Ala Asn Ser Lys Phe Ile Gly Ile  
85 90 95

Thr Glu Leu Arg Arg Thr Arg Gln Phe Leu Asp Tyr Leu Gln Glu Phe  
100 105 110

Leu Gly Ser Met Asn Thr Ala Ala Ile Ile Glu Gly  
115 120

<210> 14  
<211> 116  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Murine IL5  
modified by substitution with tetanus toxoid P2  
epitope

<220>  
<221> MUTAGEN  
<222> (30)..(44)  
<223> Tetanus toxoid P2 epitope (SEQ ID NO: 23)

<220>  
<221> SIMILAR  
<222> (1)..(29)  
<223> Identical to residues 1-29 in SEQ ID NO: 12

<220>  
<221> SIMILAR  
<222> (45)..(116)  
<223> Identical to residues 42-113 in SEQ ID NO: 12

<400> 14  
Met Glu Ile Pro Met Ser Thr Val Val Lys Glu Thr Leu Ala Leu Leu  
1 5 10 15

Ser Ala His Arg Ala Leu Leu Thr Ser Asn Glu Thr Met Gln Tyr Ile  
20 25 30

Lys Ala Asn Ser Lys Phe Ile Gly Ile Thr Glu Leu Cys Ile Gly Glu  
35 40 45

Ile Phe Gln Gly Leu Asp Ile Leu Lys Asp Gln Thr Val Arg Gly Gly  
50 55 60

Thr Val Met Arg Leu Phe Gln Asn Leu Ser Leu Ile Lys Lys Tyr Ile  
65 70 75 80

Asp Arg Gln Glu Lys Lys Cys Gly Glu Glu Arg Arg Arg Thr Arg Gln  
85 90 95

205120"9160650



Phe Leu Asp Tyr Leu Gln Glu Phe Leu Gly Ser Met Asn Thr Ala Ala  
 100 105 110

Ile Ile Glu Gly  
 115

<210> 15  
 <211> 122  
 <212> PRT  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: Murine IL5  
 modified by substitution with tetanus toxoid P2  
 epitope

<220>  
 <221> MUTAGEN  
 <222> (57)..(71)  
 <223> Tetanus toxoid P2 epitope (SEQ ID NO: 23)

<220>  
 <221> SIMILAR  
 <222> (1)..(56)  
 <223> Identical to residues 1-56 in SEQ ID NO: 12

<220>  
 <221> SIMILAR  
 <222> (72)..(122)  
 <223> Identical to residues 63-113 in SEQ ID NO: 12

<400> 15  
 Met Glu Ile Pro Met Ser Thr Val Val Lys Glu Thr Leu Ala Leu Leu  
 1 5 10 15

Ser Ala His Arg Ala Leu Leu Thr Ser Asn Glu Thr Met Arg Leu Pro  
 20 25 30

Val Pro Thr His Lys Asn His Gln Leu Cys Ile Gly Glu Ile Phe Gln  
 35 40 45

Gly Leu Asp Ile Leu Lys Asp Gln Gln Tyr Ile Lys Ala Asn Ser Lys  
 50 55 60

Phe Ile Gly Ile Thr Glu Leu Val Met Arg Leu Phe Gln Asn Leu Ser  
 65 70 75 80

Leu Ile Lys Lys Tyr Ile Asp Arg Gln Glu Lys Lys Cys Gly Glu Glu  
 85 90 95

Arg Arg Arg Thr Arg Gln Phe Leu Asp Tyr Leu Gln Glu Phe Leu Gly  
 100 105 110

Ser Met Asn Thr Ala Ala Ile Ile Glu Gly

05980916 021902  
 2005.12.29 9:09:59

115

120

<210> 16  
 <211> 122  
 <212> PRT  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: Murine IL5  
 modified by substitution with tetanus toxoid P2  
 epitope

<220>  
 <221> MUTAGEN  
 <222> (84)..(98)  
 <223> Tetanus toxoid P2 epitope (SEQ ID NO: 23)

<220>  
 <221> SIMILAR  
 <222> (1)..(83)  
 <223> Identical to residues 1-83 in SEQ ID NO: 12

<220>  
 <221> SIMILAR  
 <222> (99)..(122)  
 <223> Identical to residues 90-113 in SEQ ID NO: 12

<400> 16  
 Met Glu Ile Pro Met Ser Thr Val Val Lys Glu Thr Leu Ala Leu Leu  
   1                  5                  10                  15  
 Ser Ala His Arg Ala Leu Leu Thr Ser Asn Glu Thr Met Arg Leu Pro  
                   20                  25                  30  
 Val Pro Thr His Lys Asn His Gln Leu Cys Ile Gly Glu Ile Phe Gln  
                   35                  40                  45  
 Gly Leu Asp Ile Leu Lys Asp Gln Thr Val Arg Gly Gly Thr Val Met  
                   50                  55                  60  
 Arg Leu Phe Gln Asn Leu Ser Leu Ile Lys Lys Tyr Ile Asp Arg Gln  
                   65                  70                  75                  80  
 Glu Lys Lys Gln Tyr Ile Lys Ala Asn Ser Lys Phe Ile Gly Ile Thr  
                   85                  90                  95  
 Glu Leu Arg Thr Arg Gln Phe Leu Asp Tyr Leu Gln Glu Phe Leu Gly  
                   100                  105                  110  
 Ser Met Asn Thr Ala Ala Ile Ile Glu Gly  
                   115                  120

<210> 17

20250916 09:58:56

<211> 124  
 <212> PRT  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: Murine IL5  
 modified by substitution with tetanus toxoid P2  
 epitope

<220>  
 <221> MUTAGEN  
 <222> (108)..(122)  
 <223> Tetanus toxoid P2 epitope (SEQ ID NO: 23)

<220>  
 <221> SIMILAR  
 <222> (1)..(107)  
 <223> Identical to residues 1-107 in SEQ ID NO: 12

<220>  
 <221> SIMILAR  
 <222> (123)..(124)  
 <223> Identical to residues 112-113 in SEQ ID NO: 12

<400> 17  
 Met Glu Ile Pro Met Ser Thr Val Val Lys Glu Thr Leu Ala Leu Leu  
   1                  5                  10                  15  
 Ser Ala His Arg Ala Leu Leu Thr Ser Asn Glu Thr Met Arg Leu Pro  
                   20                  25                  30  
 Val Pro Thr His Lys Asn His Gln Leu Cys Ile Gly Glu Ile Phe Gln  
           35                  40                  45  
 Gly Leu Asp Ile Leu Lys Asp Gln Thr Val Arg Gly Gly Thr Val Met  
   50                  55                  60  
 Arg Leu Phe Gln Asn Leu Ser Leu Ile Lys Lys Tyr Ile Asp Arg Gln  
   65                  70                  75                  80  
 Glu Lys Lys Cys Gly Glu Glu Arg Arg Arg Thr Arg Gln Phe Leu Asp  
                   85                  90                  95  
 Tyr Leu Gln Glu Phe Leu Gly Ser Met Asn Thr Gln Tyr Ile Lys Ala  
           100                  105                  110  
 Asn Ser Lys Phe Ile Gly Ile Thr Glu Leu Glu Gly  
   115                  120

<210> 18  
 <211> 130  
 <212> PRT  
 <213> Artificial Sequence

<220>

09900916 064500



```

<220>
<221> MUTAGEN
<222> (30)..(50)
<223> Tetanus toxoid P30 epitope (SEQ ID NO: 24)

<220>
<221> SIMILAR
<222> (1)..(29)
<223> Identical to residues 1-29 in SEQ ID NO: 12

<220>
<221> SIGNAL
<222> (51)..(122)
<223> Identical to residues 42-113 in SEQ ID NO: 12

<400> 19
Met Glu Ile Pro Met Ser Thr Val Val Lys Glu Thr Leu Ala Leu Leu
  1             5             10             15
Ser Ala His Arg Ala Leu Leu Thr Ser Asn Glu Thr Met Phe Asn Asn
          20             25             30
Phe Thr Val Ser Phe Trp Leu Arg Val Pro Lys Val Ser Ala Ser His
          35             40             45
Leu Glu Cys Ile Gly Glu Ile Phe Gln Gly Leu Asp Ile Leu Lys Asp
          50             55             60
Gln Thr Val Arg Gly Gly Thr Val Met Arg Leu Phe Gln Asn Leu Ser
          65             70             75             80
Leu Ile Lys Lys Tyr Ile Asp Arg Gln Glu Lys Lys Cys Gly Glu Glu
          85             90             95
Arg Arg Arg Thr Arg Gln Phe Leu Asp Tyr Leu Gln Glu Phe Leu Gly
          100            105            110
Ser Met Asn Thr Ala Ala Ile Ile Glu Gly
          115            120

<210> 20
<211> 128
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:Murine IL5
      modified by substitution with tetanus toxoid P30
      epitope

<220>
<221> MUTAGEN
<222> (57)..(77)

```

<223> Tetanus toxoid P30 epitope (SEQ ID NO: 24)

<220>

<221> SIMILAR

<222> (1)..(56)

<223> Identical to residues 1-56 in SEQ ID NO: 12

<220>

<221> SIMILAR

<222> (78)..(128)

<223> Identical to residues 63-113 in SEQ ID NO: 12

<400> 20

Met Glu Ile Pro Met Ser Thr Val Val Lys Glu Thr Leu Ala Leu Leu  
1 5 10 15

Ser Ala His Arg Ala Leu Leu Thr Ser Asn Glu Thr Met Arg Leu Pro  
20 25 30

Val Pro Thr His Lys Asn His Gln Leu Cys Ile Gly Glu Ile Phe Gln  
35 40 45

Gly Leu Asp Ile Leu Lys Asp Gln Phe Asn Asn Phe Thr Val Ser Phe  
50 55 60

Trp Leu Arg Val Pro Lys Val Ser Ala Ser His Leu Glu Val Met Arg  
65 70 75 80

Leu Phe Gln Asn Leu Ser Leu Ile Lys Lys Tyr Ile Asp Arg Gln Glu  
85 90 95

Lys Lys Cys Gly Glu Glu Arg Arg Arg Thr Arg Gln Phe Leu Asp Tyr  
100 105 110

Leu Gln Glu Phe Leu Gly Ser Met Asn Thr Ala Ala Ile Ile Glu Gly  
115 120 125

<210> 21

<211> 130

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Murine IL5  
modified by substitution with tetanus toxoid P30  
epitope

<220>

<221> MUTAGEN

<222> (108)..(128)

<223> Tetanus toxoid P30 epitope (SEQ ID NO: 24)

205120 "9T68650

<220>  
 <221> SIMILAR  
 <222> (1)..(107)  
 <223> Identical to residues 1-107 in SEQ ID NO: 12

<220>  
 <221> SIMILAR  
 <222> (129)..(130)  
 <223> Identical to residues 112-113 in SEQ ID NO: 12

<400> 21  
 Met Glu Ile Pro Met Ser Thr Val Val Lys Glu Thr Leu Ala Leu Leu  
           1                  5                  10                  15  
 Ser Ala His Arg Ala Leu Leu Thr Ser Asn Glu Thr Met Arg Leu Pro  
                   20                  25                  30  
 Val Pro Thr His Lys Asn His Gln Leu Cys Ile Gly Glu Ile Phe Gln  
                   35                  40                  45  
 Gly Leu Asp Ile Leu Lys Asp Gln Thr Val Arg Gly Gly Thr Val Met  
           50                  55                  60  
 Arg Leu Phe Gln Asn Leu Ser Leu Ile Lys Lys Tyr Ile Asp Arg Gln  
           65                  70                  75                  80  
 Glu Lys Lys Cys Gly Glu Glu Arg Arg Arg Thr Arg Gln Phe Leu Asp  
                   85                  90                  95  
 Tyr Leu Gln Glu Phe Leu Gly Ser Met Asn Thr Phe Asn Asn Phe Thr  
                   100                  105                  110  
 Val Ser Phe Trp Leu Arg Val Pro Lys Val Ser Ala Ser His Leu Glu  
           115                  120                  125  
 Glu Gly  
           130

<210> 22  
 <211> 139  
 <212> PRT  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: Murine IL5  
       modified by substitution with tetanus toxoid P2  
       and P30 epitopes

<220>  
 <221> MUTAGEN  
 <222> (84)..(98)  
 <223> Tetanus toxoid P2 epitope (SEQ ID NO: 23)

<220>  
 <221> MUTAGEN

20250420 16:05:00

<222> (117)..(137)

<223> Tetanus toxoid P30 epitope (SEQ ID NO: 24)

<220>

<221> SIMILAR

<222> (1)..(83)

<223> Identical to residues 1-83 in SEQ ID NO: 12

<220>

<221> SIMILAR

<222> (99)..(116)

<223> Identical to residues 90-109 in SEQ ID NO: 12

<220>

<221> SIMILAR

<222> (138)..(139)

<223> Identical to residues 112-113 in SEQ ID NO: 12

<400> 22

Met Glu Ile Pro Met Ser Thr Val Val Lys Glu Thr Leu Ala Leu Leu  
1 5 10 15

Ser Ala His Arg Ala Leu Leu Thr Ser Asn Glu Thr Met Arg Leu Pro  
20 25 30

Val Pro Thr His Lys Asn His Gln Leu Cys Ile Gly Glu Ile Phe Gln  
35 40 45

Gly Leu Asp Ile Leu Lys Asp Gln Thr Val Arg Gly Gly Thr Val Met  
50 55 60

Arg Leu Phe Gln Asn Leu Ser Leu Ile Lys Lys Tyr Ile Asp Arg Gln  
65 70 75 80

Glu Lys Lys Gln Tyr Ile Lys Ala Asn Ser Lys Phe Ile Gly Ile Thr  
85 90 95

Glu Leu Arg Thr Arg Gln Phe Leu Asp Tyr Leu Gln Glu Phe Leu Gly  
100 105 110

Ser Met Asn Thr Phe Asn Asn Phe Thr Val Ser Phe Trp Leu Arg Val  
115 120 125

Pro Lys Val Ser Ala Ser His Leu Glu Glu Gly  
130 135

<210> 23

<211> 15

<212> PRT

<213> Clostridium tetani

<400> 23

Gln Tyr Ile Lys Ala Asn Ser Lys Phe Ile Gly Ile Thr Glu Leu  
1 5 10 15

20051220 09:50:50



<210> 24  
 <211> 21  
 <212> PRT  
 <213> Clostridium tetani

<400> 24  
 Phe Asn Asn Phe Thr Val Ser Phe Trp Leu Arg Val Pro Lys Val Ser  
   1                  5                  10                  15  
 Ala Ser His Leu Glu  
                   20

<210> 25  
 <211> 45  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <221> CDS  
 <222> (1)..(45)  
 <220>  
 <223> Description of Artificial sequence: DNA encoding His tag

<400> 25  
 atg aaa cac caa cac caa cat caa cat caa cat caa cat caa caa       45  
 Met Lys His Gln His Gln His Gln His Gln His Gln His Gln Gln  
   1                  5                  10                  15

<210> 26  
 <211> 15  
 <212> PRT  
 <213> Artificial Sequence

<220>  
 <223> Artificial Sequence

<400> 26  
 Met Lys His Gln His Gln His Gln His Gln His Gln His Gln Gln  
   1                  5                  10                  15

<210> 27  
 <211> 381  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: Human IL-5  
       modified by substitution with tetanus toxoid  
       epitope

2005-07-20 16:02:16

<220>  
 <221> CDS  
 <222> (1)..(381)

<220>  
 <221> mutation  
 <222> (262)..(306)  
 <223> Tetanus toxoid P2 epitope

<220>  
 <221> misc\_feature  
 <222> (1)..(261)  
 <223> DNA encoding amino acids 1-87 of human IL5

<220>  
 <221> misc\_feature  
 <222> (307)..(378)  
 <223> DNA encoding amino acids 92-115 of human IL5

<400> 27  
 atc ccc aca gaa att ccc aca agt gca ttg gtg aaa gag acc ttg gca 48  
 Ile Pro Thr Glu Ile Pro Thr Ser Ala Leu Val Lys Glu Thr Leu Ala  
     1                    5                    10                    15  
  
 ctg ctt tct act cat cga act ctg ctg ata gcc aat gag act ctc cgg 96  
 Leu Leu Ser Thr His Arg Thr Leu Leu Ile Ala Asn Glu Thr Leu Arg  
                     20                    25                    30  
  
 att cct gtt cct gta cat aaa aat cac caa ctg tgc act gaa gaa atc 144  
 Ile Pro Val Pro Val His Lys Asn His Gln Leu Cys Thr Glu Glu Ile  
                     35                    40                    45  
  
 ttt cag gga ata ggc aca ctc gag agt caa act gtg caa ggg ggt act 192  
 Phe Gln Gly Ile Gly Thr Leu Glu Ser Gln Thr Val Gln Gly Gly Thr  
                     50                    55                    60  
  
 gtg gaa aga cta ttc aaa aac ttg tcc tta ata aag aaa tac atc gat 240  
 Val Glu Arg Leu Phe Lys Asn Leu Ser Leu Ile Lys Lys Tyr Ile Asp  
                     65                    70                    75                    80  
  
 ggc caa aaa aaa aag tgt gga cag tac atc aag gcc aac tcc aag ttc 288  
 Gly Gln Lys Lys Lys Cys Gly Gln Tyr Ile Lys Ala Asn Ser Lys Phe  
                     85                    90                    95  
  
 atc ggc atc acc gag ctg aga gta aac caa ttc cta gac tat ctg cag 336  
 Ile Gly Ile Thr Glu Leu Arg Val Asn Gln Phe Leu Asp Tyr Leu Gln  
                     100                    105                    110  
  
 gag ttt ctt ggt gta atg aac acc gag tgg ata ata gaa agt tga 381  
 Glu Phe Leu Gly Val Met Asn Thr Glu Trp Ile Ile Glu Ser  
                     115                    120                    125

<210> 28  
 <211> 126  
 <212> PRT

20051209 09:06:50

$\langle 220 \rangle$ 

<400> 28

Glu Phe Leu Gly Val Met Asn Thr Glu Trp Ile Ile Glu Ser  
115 120 125

<211> 375

<212> DNA

### <213> Artificial Sequence

 $\langle 220 \rangle$  $\langle 220 \rangle$ 

<221> CDS

<222> (1) .. (375)

 $\langle 220 \rangle$ 

<221> mutation

<222> (94) .. (156)

<223> Tetanus toxoid P30 epitope

 $\langle 220 \rangle$ 

<221> misc feature

<222> (1) .. (93)

<223> DNA encoding amino acids 1-31 of human IL5

<220>  
 <221> misc\_feature  
 <222> (157)..(372)  
 <223> DNA encoding amino acids 44-115 of human IL5

<400> 29

atc	ccc	aca	gaa	att	ccc	aca	agt	gca	ttg	gtg	aaa	gag	acc	ttg	gca	48
Ile	Pro	Thr	Glu	Ile	Pro	Thr	Ser	Ala	Leu	Val	Lys	Glu	Thr	Leu	Ala	
1				5					10					15		
ctg	ctt	tct	act	cat	cga	act	ctg	ctg	ata	gcc	aat	gag	act	ctc	ttc	96
Leu	Leu	Ser	Thr	His	Arg	Thr	Leu	Leu	Ile	Ala	Asn	Glu	Thr	Leu	Phe	
			20					25					30			
aac	aac	ttc	acc	gtg	agc	ttc	tgg	ctg	cgc	gtg	cct	aag	gtg	agc	gcc	144
Asn	Asn	Phe	Thr	Val	Ser	Phe	Trp	Leu	Arg	Val	Pro	Lys	Val	Ser	Ala	
		35					40					45				
agc	cac	ctg	gag	tgc	act	gaa	gaa	atc	ttt	cag	gga	ata	ggc	aca	ctc	192
Ser	His	Leu	Glu	Cys	Thr	Glu	Glu	Ile	Phe	Gln	Gly	Ile	Gly	Thr	Leu	
		50				55					60					
gag	agt	caa	act	gtg	caa	ggg	ggt	act	gtg	gaa	aga	cta	ttc	aaa	aac	240
Glu	Ser	Gln	Thr	Val	Gln	Gly	Gly	Thr	Val	Glu	Arg	Leu	Phe	Lys	Asn	
65					70				75					80		
ttg	tcc	tta	ata	aag	aaa	tac	atc	gat	ggc	caa	aaa	aaa	aag	tgt	gga	288
Leu	Ser	Leu	Ile	Lys	Lys	Tyr	Ile	Asp	Gly	Gln	Lys	Lys	Lys	Cys	Gly	
				85				90						95		
gaa	gaa	aga	cgg	aga	gta	aac	caa	ttc	cta	gac	tat	ctg	cag	gag	ttt	336
Glu	Glu	Arg	Arg	Arg	Val	Asn	Gln	Phe	Leu	Asp	Tyr	Leu	Gln	Glu	Phe	
			100				105						110			
ctt	ggt	gta	atg	aac	acc	gag	tgg	ata	ata	gaa	agt	tga				375
Leu	Gly	Val	Met	Asn	Thr	Glu	Trp	Ile	Ile	Glu	Ser					
		115				120										

<210> 30  
 <211> 124  
 <212> PRT  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: Human Il-5  
 modified by substitution with tetanus toxoid  
 epitope

<400> 30

Ile	Pro	Thr	Glu	Ile	Pro	Thr	Ser	Ala	Leu	Val	Lys	Glu	Thr	Leu	Ala
1				5					10					15	
Leu	Leu	Ser	Thr	His	Arg	Thr	Leu	Leu	Ile	Ala	Asn	Glu	Thr	Leu	Phe
			20				25						30		

20250706 15:00:00

```

Asn Asn Phe Thr Val Ser Phe Trp Leu Arg Val Pro Lys Val Ser Ala
   35           40           45

Ser His Leu Glu Cys Thr Glu Glu Ile Phe Gln Gly Ile Gly Thr Leu
   50           55           60

Glu Ser Gln Thr Val Gln Gly Gly Thr Val Glu Arg Leu Phe Lys Asn
   65           70           75           80

Leu Ser Leu Ile Lys Lys Tyr Ile Asp Gly Gln Lys Lys Lys Cys Gly
           85           90           95

Glu Glu Arg Arg Arg Val Asn Gln Phe Leu Asp Tyr Leu Gln Glu Phe
   100           105           110

Leu Gly Val Met Asn Thr Glu Trp Ile Ile Glu Ser
   115           120

```

```

<210> 31
<211> 393
<212> DNA
<213> Artificial Sequence

```

```

<220>
<223> Description of Artificial Sequence: Human IL-5
      modified by substitution with tetanus toxoid
      epitope

```

```

<220>
<221> CDS
<222> (1)..(393)

```

```

<220>
<221> mutation
<222> (175)..(237)
<223> Tetanus toxoid P30 epitope

```

```

<220>
<221> misc_feature
<222> (1)..(174)
<223> DNA encoding amino acids 1-58 of human IL5

```

```

<220>
<221> misc_feature
<222> (238)..(390)
<223> DNA encoding amino acids 65-115 of human IL5

```

```

<400> 31
atc ccc aca gaa att ccc aca agt gca ttg gtg aaa gag acc ttg gca   48
Ile Pro Thr Glu Ile Pro Thr Ser Ala Leu Val Lys Glu Thr Leu Ala
   1           5           10           15

ctg ctt tct act cat cga act ctg ctg ata gcc aat gag act ctc cgg   96
Leu Leu Ser Thr His Arg Thr Leu Leu Ile Ala Asn Glu Thr Leu Arg
           20           25           30

```

05980916-021902

att cct gtt cct gta cat aaa aat cac caa ctg tgc act gaa gaa atc 144  
 Ile Pro Val Pro Val His Lys Asn His Gln Leu Cys Thr Glu Glu Ile  
           35                          40                          45

ttt cag gga ata ggc aca ctc gag agt caa ttc aac aac ttc acc gtg 192  
 Phe Gln Gly Ile Gly Thr Leu Glu Ser Gln Phe Asn Asn Phe Thr Val  
           50                          55                          60

agc ttc tgg ctg cgc gtg cct aag gtg agc gcc agc cac ctg gag gtg 240  
 Ser Phe Trp Leu Arg Val Pro Lys Val Ser Ala Ser His Leu Glu Val  
           65                          70                          75                          80

gaa aga cta ttc aaa aac ttg tcc tta ata aag aaa tac atc gat ggc 288  
 Glu Arg Leu Phe Lys Asn Leu Ser Leu Ile Lys Lys Tyr Ile Asp Gly  
                           85                          90                          95

caa aaa aaa aag tgt gga gaa gaa aga cgg aga gta aac caa ttc cta 336  
 Gln Lys Lys Lys Cys Gly Glu Glu Arg Arg Arg Val Asn Gln Phe Leu  
                           100                          105                          110

gac tat ctg cag gag ttt ctt ggt gta atg aac acc gag tgg ata ata 384  
 Asp Tyr Leu Gln Glu Phe Leu Gly Val Met Asn Thr Glu Trp Ile Ile  
           115                          120                          125

gaa agt tga 393  
 Glu Ser  
           130

<210> 32  
 <211> 130  
 <212> PRT  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: Human Il-5  
           modified by substitution with tetanus toxoid  
           epitope

<400> 32  
 Ile Pro Thr Glu Ile Pro Thr Ser Ala Leu Val Lys Glu Thr Leu Ala  
   1                          5                          10                          15

Leu Leu Ser Thr His Arg Thr Leu Leu Ile Ala Asn Glu Thr Leu Arg  
           20                          25                          30

Ile Pro Val Pro Val His Lys Asn His Gln Leu Cys Thr Glu Glu Ile  
           35                          40                          45

Phe Gln Gly Ile Gly Thr Leu Glu Ser Gln Phe Asn Asn Phe Thr Val  
           50                          55                          60

Ser Phe Trp Leu Arg Val Pro Lys Val Ser Ala Ser His Leu Glu Val  
           65                          70                          75                          80

Glu Arg Leu Phe Lys Asn Leu Ser Leu Ile Lys Lys Tyr Ile Asp Gly

095805916 021903  
 2005120 97508550

85 90 95

Gln Lys Lys Lys Cys Gly Glu Glu Arg Arg Arg Val Asn Gln Phe Leu  
100 105 110

Asp Tyr Leu Gln Glu Phe Leu Gly Val Met Asn Thr Glu Trp Ile Ile  
115 120 125

Glu Ser  
130

<210> 33  
<211> 375  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Human IL-5  
modified by substitution with tetanus toxoid  
epitope

<220>  
<221> CDS  
<222> (1)..(375)

<220>  
<221> mutation  
<222> (175)..(219)  
<223> Tetanus toxoid P2 epitope

<220>  
<221> misc\_feature  
<222> (1)..(174)  
<223> DNA encoding amino acids 1-58 of human IL5

<220>  
<221> misc\_feature  
<222> (220)..(372)  
<223> DNA encoding amino acids 65-115 of human IL5

<400> 33  
atc ccc aca gaa att ccc aca agt gca ttg gtg aaa gag acc ttg gca 48  
Ile Pro Thr Glu Ile Pro Thr Ser Ala Leu Val Lys Glu Thr Leu Ala  
1 5 10 15

ctg ctt tct act cat cga act ctg ctg ata gcc aat gag act ctc cgg 96  
Leu Leu Ser Thr His Arg Thr Leu Leu Ile Ala Asn Glu Thr Leu Arg  
20 25 30

att cct gtt cct gta cat aaa aat cac caa ctg tgc act gaa gaa atc 144  
Ile Pro Val Pro Val His Lys Asn His Gln Leu Cys Thr Glu Glu Ile  
35 40 45

ttt cag gga ata ggc aca ctc gag agt caa cag tac atc aag gcc aac 192  
Phe Gln Gly Ile Gly Thr Leu Glu Ser Gln Gln Tyr Ile Lys Ala Asn

05980976 DE 1992

50	55	60	
tcc aag ttc atc ggc atc acc gag ctg gtg gaa aga cta ttc aaa aac			240
Ser Lys Phe Ile Gly Ile Thr Glu Leu Val Glu Arg Leu Phe Lys Asn			
65	70	75	80
ttg tcc tta ata aag aaa tac atc gat ggc caa aaa aaa aag tgt gga			288
Leu Ser Leu Ile Lys Lys Tyr Ile Asp Gly Gln Lys Lys Lys Cys Gly			
	85	90	95
gaa gaa aga cgg aga gta aac caa ttc cta gac tat ctg cag gag ttt			336
Glu Glu Arg Arg Arg Val Asn Gln Phe Leu Asp Tyr Leu Gln Glu Phe			
	100	105	110
ctt ggt gta atg aac acc gag tgg ata ata gaa agt tga			375
Leu Gly Val Met Asn Thr Glu Trp Ile Ile Glu Ser			
	115	120	

&lt;210&gt; 34

&lt;211&gt; 124

&lt;212&gt; PRT

&lt;213&gt; Artificial Sequence

&lt;220&gt;

<223> Description of Artificial Sequence: Human Il-5  
modified by substitution with tetanus toxoid  
epitope

&lt;400&gt; 34

Ile Pro Thr Glu Ile Pro Thr Ser Ala Leu Val Lys Glu Thr Leu Ala			
1	5	10	15

Leu Leu Ser Thr His Arg Thr Leu Leu Ile Ala Asn Glu Thr Leu Arg			
	20	25	30

Ile Pro Val Pro Val His Lys Asn His Gln Leu Cys Thr Glu Glu Ile			
	35	40	45

Phe Gln Gly Ile Gly Thr Leu Glu Ser Gln Gln Tyr Ile Lys Ala Asn			
	50	55	60

Ser Lys Phe Ile Gly Ile Thr Glu Leu Val Glu Arg Leu Phe Lys Asn			
	65	70	75

Leu Ser Leu Ile Lys Lys Tyr Ile Asp Gly Gln Lys Lys Lys Cys Gly			
	85	90	95

Glu Glu Arg Arg Arg Val Asn Gln Phe Leu Asp Tyr Leu Gln Glu Phe			
	100	105	110

Leu Gly Val Met Asn Thr Glu Trp Ile Ile Glu Ser		
	115	120

&lt;210&gt; 35

20250909 14:56:56



<211> 357  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: Human Il-5  
 modified by substitution with tetanus toxoid  
 epitope

<220>  
 <221> CDS  
 <222> (1)..(357)

<220>  
 <221> mutation  
 <222> (94)..(138)  
 <223> Tetanus toxoid P2 epitope

<220>  
 <221> misc\_feature  
 <222> (1)..(93)  
 <223> DNA encoding amino acids 1-31 of human IL5

<220>  
 <221> misc\_feature  
 <222> (139)..(354)  
 <223> DNA encoding amino acids 44-115 of human IL5

<400> 35  
 atc ccc aca gaa att ccc aca agt gca ttg gtg aaa gag acc ttg gca 48  
 Ile Pro Thr Glu Ile Pro Thr Ser Ala Leu Val Lys Glu Thr Leu Ala  
 1 5 10 15  
 ctg ctt tct act cat cga act ctg ctg ata gcc aat gag act ctc cag 96  
 Leu Leu Ser Thr His Arg Thr Leu Leu Ile Ala Asn Glu Thr Leu Gln  
 20 25 30  
 tac atc aag gcc aac tcc aag ttc atc ggc atc acc gag ctg tgc act 144  
 Tyr Ile Lys Ala Asn Ser Lys Phe Ile Gly Ile Thr Glu Leu Cys Thr  
 35 40 45  
 gaa gaa atc ttt cag gga ata ggc aca ctc gag agt caa act gtg caa 192  
 Glu Glu Ile Phe Gln Gly Ile Gly Thr Leu Glu Ser Gln Thr Val Gln  
 50 55 60  
 ggg ggt act gtg gaa aga cta ttc aaa aac ttg tcc tta ata aag aaa 240  
 Gly Gly Thr Val Glu Arg Leu Phe Lys Asn Leu Ser Leu Ile Lys Lys  
 65 70 75 80  
 tac atc gat ggc caa aaa aaa aag tgt gga gaa gaa aga cgg aga gta 288  
 Tyr Ile Asp Gly Gln Lys Lys Lys Cys Gly Glu Glu Arg Arg Arg Val  
 85 90 95  
 aac caa ttc cta gac tat ctg cag gag ttt ctt ggt gta atg aac acc 336  
 Asn Gln Phe Leu Asp Tyr Leu Gln Glu Phe Leu Gly Val Met Asn Thr  
 100 105 110

2005 Feb 24 09:56:50

gag tgg ata ata gaa agt tga  
 Glu Trp Ile Ile Glu Ser  
 115

357

<210> 36  
 <211> 118  
 <212> PRT  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: Human Il-5  
 modified by substitution with tetanus toxoid  
 epitope

<400> 36  
 Ile Pro Thr Glu Ile Pro Thr Ser Ala Leu Val Lys Glu Thr Leu Ala  
 1 5 10 15  
 Leu Leu Ser Thr His Arg Thr Leu Leu Ile Ala Asn Glu Thr Leu Gln  
 20 25 30  
 Tyr Ile Lys Ala Asn Ser Lys Phe Ile Gly Ile Thr Glu Leu Cys Thr  
 35 40 45  
 Glu Glu Ile Phe Gln Gly Ile Gly Thr Leu Glu Ser Gln Thr Val Gln  
 50 55 60  
 Gly Gly Thr Val Glu Arg Leu Phe Lys Asn Leu Ser Leu Ile Lys Lys  
 65 70 75 80  
 Tyr Ile Asp Gly Gln Lys Lys Lys Cys Gly Glu Glu Arg Arg Arg Val  
 85 90 95  
 Asn Gln Phe Leu Asp Tyr Leu Gln Glu Phe Leu Gly Val Met Asn Thr  
 100 105 110  
 Glu Trp Ile Ile Glu Ser  
 115

<210> 37  
 <211> 375  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: Human Il-5  
 modified by substitution with tetanus toxoid  
 epitope

<220>  
 <221> CDS  
 <222> (1)..(375)  
 <220>

20250409 15:03:50

<221> mutation  
 <222> (256)..(300)  
 <223> Tetanus toxoid P2 epitope

<220>  
 <221> misc\_feature  
 <222> (1)..(255)  
 <223> DNA encoding amino acids 1-85 of human IL5

<220>  
 <221> misc\_feature  
 <222> (301)..(372)  
 <223> DNA encoding amino acids 92-115 of human IL5

<400> 37  
 atc ccc aca gaa att ccc aca agt gca ttg gtg aaa gag acc ttg gca 48  
 Ile Pro Thr Glu Ile Pro Thr Ser Ala Leu Val Lys Glu Thr Leu Ala  
 1 5 10 15  
 ctg ctt tct act cat cga act ctg ctg ata gcc aat gag act ctc cgg 96  
 Leu Leu Ser Thr His Arg Thr Leu Leu Ile Ala Asn Glu Thr Leu Arg  
 20 25 30  
 att cct gtt cct gta cat aaa aat cac caa ctg tgc act gaa gaa atc 144  
 Ile Pro Val Pro Val His Lys Asn His Gln Leu Cys Thr Glu Glu Ile  
 35 40 45  
 ttt cag gga ata ggc aca ctc gag agt caa act gtg caa ggg ggt act 192  
 Phe Gln Gly Ile Gly Thr Leu Glu Ser Gln Thr Val Gln Gly Gly Thr  
 50 55 60  
 gtg gaa aga cta ttc aaa aac ttg tcc tta ata aag aaa tac atc gat 240  
 Val Glu Arg Leu Phe Lys Asn Leu Ser Leu Ile Lys Lys Tyr Ile Asp  
 65 70 75 80  
 ggc caa aaa aaa aag cag tac atc aag gcc aac tcc aag ttc atc ggc 288  
 Gly Gln Lys Lys Lys Gln Tyr Ile Lys Ala Asn Ser Lys Phe Ile Gly  
 85 90 95  
 atc acc gag ctg aga gta aac caa ttc cta gac tat ctg cag gag ttt 336  
 Ile Thr Glu Leu Arg Val Asn Gln Phe Leu Asp Tyr Leu Gln Glu Phe  
 100 105 110  
 ctt ggt gta atg aac acc gag tgg ata ata gaa agt tga 375  
 Leu Gly Val Met Asn Thr Glu Trp Ile Ile Glu Ser  
 115 120

<210> 38  
 <211> 124  
 <212> PRT  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: Human Il-5  
 modified by substitution with tetanus toxoid  
 epitope

20250916 09:49:02

&lt;400&gt; 38

Ile Pro Thr Glu Ile Pro Thr Ser Ala Leu Val Lys Glu Thr Leu Ala  
 1 5 10 15

Leu Leu Ser Thr His Arg Thr Leu Leu Ile Ala Asn Glu Thr Leu Arg  
 20 25 30

Ile Pro Val Pro Val His Lys Asn His Gln Leu Cys Thr Glu Glu Ile  
 35 40 45

Phe Gln Gly Ile Gly Thr Leu Glu Ser Gln Thr Val Gln Gly Gly Thr  
 50 55 60

Val Glu Arg Leu Phe Lys Asn Leu Ser Leu Ile Lys Lys Tyr Ile Asp  
 65 70 75 80

Gly Gln Lys Lys Lys Gln Tyr Ile Lys Ala Asn Ser Lys Phe Ile Gly  
 85 90 95

Ile Thr Glu Leu Arg Val Asn Gln Phe Leu Asp Tyr Leu Gln Glu Phe  
 100 105 110

Leu Gly Val Met Asn Thr Glu Trp Ile Ile Glu Ser  
 115 120

&lt;210&gt; 39

&lt;211&gt; 399

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

<223> Description of Artificial Sequence: Human IL-5  
 modified by substitution with tetanus toxoid  
 epitope

&lt;220&gt;

&lt;221&gt; CDS

&lt;222&gt; (1)..(399)

&lt;220&gt;

&lt;221&gt; mutation

&lt;222&gt; (262)..(324)

&lt;223&gt; Tetanus toxoid P30 epitope

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;222&gt; (1)..(261)

&lt;223&gt; DNA encoding amino acids 1-87 of human IL5

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;222&gt; (325)..(396)

&lt;223&gt; DNA encoding amino acids 92-115 of human IL5

2005-12-20 09:03:50

&lt;400&gt; 39

atc ccc aca gaa att ccc aca agt gca ttg gtg aaa gag acc ttg gca 48  
 Ile Pro Thr Glu Ile Pro Thr Ser Ala Leu Val Lys Glu Thr Leu Ala  
     1                    5                    10                    15

ctg ctt tct act cat cga act ctg ctg ata gcc aat gag act ctc cgg 96  
 Leu Leu Ser Thr His Arg Thr Leu Leu Ile Ala Asn Glu Thr Leu Arg  
                     20                    25                    30

att cct gtt cct gta cat aaa aat cac caa ctg tgc act gaa gaa atc 144  
 Ile Pro Val Pro Val His Lys Asn His Gln Leu Cys Thr Glu Glu Ile  
                     35                    40                    45

ttt cag gga ata ggc aca ctc gag agt caa act gtg caa ggg ggt act 192  
 Phe Gln Gly Ile Gly Thr Leu Glu Ser Gln Thr Val Gln Gly Gly Thr  
                     50                    55                    60

gtg gaa aga cta ttc aaa aac ttg tcc tta ata aag aaa tac atc gat 240  
 Val Glu Arg Leu Phe Lys Asn Leu Ser Leu Ile Lys Lys Tyr Ile Asp  
                     65                    70                    75                    80

ggc caa aaa aaa aag tgt gga ttc aac aac ttc acc gtg agc ttc tgg 288  
 Gly Gln Lys Lys Lys Cys Gly Phe Asn Asn Phe Thr Val Ser Phe Trp  
                     85                    90                    95

ctg cgc gtg cct aag gtg agc gcc agc cac ctg gag aga gta aac caa 336  
 Leu Arg Val Pro Lys Val Ser Ala Ser His Leu Glu Arg Val Asn Gln  
                     100                    105                    110

ttc cta gac tat ctg cag gag ttt ctt ggt gta atg aac acc gag tgg 384  
 Phe Leu Asp Tyr Leu Gln Glu Phe Leu Gly Val Met Asn Thr Glu Trp  
                     115                    120                    125

ata ata gaa agt tga 399  
 Ile Ile Glu Ser  
                     130

&lt;210&gt; 40

&lt;211&gt; 132

&lt;212&gt; PRT

&lt;213&gt; Artificial Sequence

&lt;220&gt;

<223> Description of Artificial Sequence: Human Il-5  
 modified by substitution with tetanus toxoid  
 epitope

&lt;400&gt; 40

Ile Pro Thr Glu Ile Pro Thr Ser Ala Leu Val Lys Glu Thr Leu Ala  
     1                    5                    10                    15

Leu Leu Ser Thr His Arg Thr Leu Leu Ile Ala Asn Glu Thr Leu Arg  
                     20                    25                    30

Ile Pro Val Pro Val His Lys Asn His Gln Leu Cys Thr Glu Glu Ile  
                     35                    40                    45

005900916"004000

Phe Gln Gly Ile Gly Thr Leu Glu Ser Gln Thr Val Gln Gly Gly Thr  
 50 55 60  
 Val Glu Arg Leu Phe Lys Asn Leu Ser Leu Ile Lys Lys Tyr Ile Asp  
 65 70 75 80  
 Gly Gln Lys Lys Lys Cys Gly Phe Asn Asn Phe Thr Val Ser Phe Trp  
 85 90 95  
 Leu Arg Val Pro Lys Val Ser Ala Ser His Leu Glu Arg Val Asn Gln  
 100 105 110  
 Phe Leu Asp Tyr Leu Gln Glu Phe Leu Gly Val Met Asn Thr Glu Trp  
 115 120 125  
 Ile Ile Glu Ser  
 130

<210> 41  
 <211> 393  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: Human IL-5  
 modified by substitution with tetanus toxoid  
 epitope

<220>  
 <221> CDS  
 <222> (1)..(393)

<220>  
 <221> mutation  
 <222> (256)..(318)  
 <223> Tetanus toxoid P30 epitope

<220>  
 <221> misc\_feature  
 <222> (1)..(255)  
 <223> DNA encoding amino acids 1-85 of human IL5

<220>  
 <221> misc\_feature  
 <222> (319)..(390)  
 <223> DNA encoding amino acids 92-115 of human IL5

<400> 41  
 atc ccc aca gaa att ccc aca agt gca ttg gtg aaa gag acc ttg gca 48  
 Ile Pro Thr Glu Ile Pro Thr Ser Ala Leu Val Lys Glu Thr Leu Ala  
 1 5 10 15  
 ctg ctt tct act cat cga act ctg ctg ata gcc aat gag act ctc cgg 96  
 Leu Leu Ser Thr His Arg Thr Leu Leu Ile Ala Asn Glu Thr Leu Arg

20050920 14:50:50

20 25 30  
 att cct gtt cct gta cat aaa aat cac caa ctg tgc act gaa gaa atc 144  
 Ile Pro Val Pro Val His Lys Asn His Gln Leu Cys Thr Glu Glu Ile  
 35 40 45  
 ttt cag gga ata ggc aca ctc gag agt caa act gtg caa ggg ggt act 192  
 Phe Gln Gly Ile Gly Thr Leu Glu Ser Gln Thr Val Gln Gly Gly Thr  
 50 55 60  
 gtg gaa aga cta ttc aaa aac ttg tcc tta ata aag aaa tac atc gat 240  
 Val Glu Arg Leu Phe Lys Asn Leu Ser Leu Ile Lys Lys Tyr Ile Asp  
 65 70 75 80  
 ggc caa aaa aaa aag ttc aac aac ttc acc gtg agc ttc tgg ctg cgc 288  
 Gly Gln Lys Lys Lys Phe Asn Asn Phe Thr Val Ser Phe Trp Leu Arg  
 85 90 95  
 gtg cct aag gtg agc gcc agc cac ctg gag aga gta aac caa ttc cta 336  
 Val Pro Lys Val Ser Ala Ser His Leu Glu Arg Val Asn Gln Phe Leu  
 100 105 110  
 gac tat ctg cag gag ttt ctt ggt gta atg aac acc gag tgg ata ata 384  
 Asp Tyr Leu Gln Glu Phe Leu Gly Val Met Asn Thr Glu Trp Ile Ile  
 115 120 125  
 gaa agt tga 393  
 Glu Ser  
 130  
 <210> 42  
 <211> 130  
 <212> PRT  
 <213> Artificial Sequence  
 <220>  
 <223> Description of Artificial Sequence: Human Il-5  
 modified by substitution with tetanus toxoid  
 epitope  
 <400> 42  
 Ile Pro Thr Glu Ile Pro Thr Ser Ala Leu Val Lys Glu Thr Leu Ala  
 1 5 10 15  
 Leu Leu Ser Thr His Arg Thr Leu Leu Ile Ala Asn Glu Thr Leu Arg  
 20 25 30  
 Ile Pro Val Pro Val His Lys Asn His Gln Leu Cys Thr Glu Glu Ile  
 35 40 45  
 Phe Gln Gly Ile Gly Thr Leu Glu Ser Gln Thr Val Gln Gly Gly Thr  
 50 55 60  
 Val Glu Arg Leu Phe Lys Asn Leu Ser Leu Ile Lys Lys Tyr Ile Asp  
 65 70 75 80

09980916 "021002"





att	cct	ggt	gta	cat	aaa	aat	cac	caa	ctg	tgc	act	gaa	gaa	atc	144	
Ile	Pro	Val	Pro	Val	His	Lys	Asn	His	Gln	Leu	Cys	Thr	Glu	Glu	Ile	
		35					40					45				
ttt	cag	gga	ata	ggc	aca	ctc	gag	agt	caa	act	gtg	caa	ggg	ggt	act	192
Phe	Gln	Gly	Ile	Gly	Thr	Leu	Glu	Ser	Gln	Thr	Val	Gln	Gly	Gly	Thr	
	50					55					60					
gtg	gaa	aga	cta	ttc	aaa	aac	ttg	tcc	tta	ata	aag	aaa	tac	atc	gat	240
Val	Glu	Arg	Leu	Phe	Lys	Asn	Leu	Ser	Leu	Ile	Lys	Lys	Tyr	Ile	Asp	
	65				70					75					80	
ggc	caa	aaa	aaa	aag	tgt	gga	cag	tac	atc	aag	gcc	aac	tcc	aag	ttc	288
Gly	Gln	Lys	Lys	Lys	Cys	Gly	Gln	Tyr	Ile	Lys	Ala	Asn	Ser	Lys	Phe	
				85					90					95		
atc	ggc	atc	acc	gag	ctg	ttc	aac	aac	ttc	acc	gtg	agc	ttc	tgg	ctg	336
Ile	Gly	Ile	Thr	Glu	Leu	Phe	Asn	Asn	Phe	Thr	Val	Ser	Phe	Trp	Leu	
			100					105					110			
cgc	gtg	cct	aag	gtg	agc	gcc	agc	cac	ctg	gag	aga	gta	aac	caa	ttc	384
Arg	Val	Pro	Lys	Val	Ser	Ala	Ser	His	Leu	Glu	Arg	Val	Asn	Gln	Phe	
		115					120					125				
cta	gac	tat	ctg	cag	gag	ttt	ctt	ggt	gta	atg	aac	acc	gag	tgg	ata	432
Leu	Asp	Tyr	Leu	Gln	Glu	Phe	Leu	Gly	Val	Met	Asn	Thr	Glu	Trp	Ile	
	130					135					140					
ata	gaa	agt	tga													444
Ile	Glu	Ser														
	145															
<210> 44																
<211> 147																
<212> PRT																
<213> Artificial Sequence																
<220>																
<223> Description of Artificial Sequence: Human Il-5																
modified by substitution with tetanus toxoid																
epitopes																
<400> 44																
Ile	Pro	Thr	Glu	Ile	Pro	Thr	Ser	Ala	Leu	Val	Lys	Glu	Thr	Leu	Ala	
	1			5					10					15		
Leu	Leu	Ser	Thr	His	Arg	Thr	Leu	Leu	Ile	Ala	Asn	Glu	Thr	Leu	Arg	
			20					25					30			
Ile	Pro	Val	Pro	Val	His	Lys	Asn	His	Gln	Leu	Cys	Thr	Glu	Glu	Ile	
		35					40					45				
Phe	Gln	Gly	Ile	Gly	Thr	Leu	Glu	Ser	Gln	Thr	Val	Gln	Gly	Gly	Thr	
	50					55					60					
Val	Glu	Arg	Leu	Phe	Lys	Asn	Leu	Ser	Leu	Ile	Lys	Lys	Tyr	Ile	Asp	

```

<210> 45
<211> 375
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Murine IL-5
      modified by substitution with tetanus toxoid
      epitope

<220>
<221> CDS
<222> (1)..(375)

<220>
<221> mutation
<222> (256)..(300)
<223> Tetanus toxoid P2 epitope

<220>
<221> misc_feature
<222> (1)..(255)
<223> DNA encoding amino acids 1-85 in murine IL5

<220>
<221> misc_feature
<222> (301)..(375)
<223> DNA encoding amino acids 90-113 of murine IL5

```

<400> 45																	
atg	gag	att	ccc	atg	agc	aca	gtg	gtg	aaa	gag	acc	ttg	aca	cag	ctg	48	
Met	Glu	Ile	Pro	Met	Ser	Thr	Val	Val	Lys	Glu	Thr	Leu	Thr	Gln	Leu		
1				5					10					15			
tcc	gct	cac	cga	gct	ctg	ttg	aca	agc	aat	gag	acg	atg	agg	ctt	cct	96	
Ser	Ala	His	Arg	Ala	Leu	Leu	Thr	Ser	Asn	Glu	Thr	Met	Arg	Leu	Pro		
			20					25					30				

gtc cct act cat aaa aat cac cag cta tgc att gga gag atc ttt cag 144  
 Val Pro Thr His Lys Asn His Gln Leu Cys Ile Gly Glu Ile Phe Gln  
                   35                                  40                                  45

ggg cta gac ata ctg aag aat caa act gtc cgt ggg ggt acc gtg gaa 192  
 Gly Leu Asp Ile Leu Lys Asn Gln Thr Val Arg Gly Gly Thr Val Glu  
                   50                                  55                                  60

atg cta ttc caa aac ctg tca tta ata aag aaa tac atc gat aga caa 240  
 Met Leu Phe Gln Asn Leu Ser Leu Ile Lys Lys Tyr Ile Asp Arg Gln  
                   65                                  70                                  75                                  80

aaa gag aag tgt ggc cag tac atc aaa gct aac tcc aaa ttc atc ggt 288  
 Lys Glu Lys Cys Gly Gln Tyr Ile Lys Ala Asn Ser Lys Phe Ile Gly  
                                   85                                  90                                  95

atc acc gag ctg agg acg agg cag ttc ctg gat tat ctg cag gag ttc 336  
 Ile Thr Glu Leu Arg Thr Arg Gln Phe Leu Asp Tyr Leu Gln Glu Phe  
                                   100                                  105                                  110

ctt ggt gtg atg agt aca gag tgg gca atg gaa ggc taa 375  
 Leu Gly Val Met Ser Thr Glu Trp Ala Met Glu Gly  
                   115                                  120

<210> 46

<211> 124

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Murine Il-5  
 modified by substitution with tetanus toxoid  
 epitope

<400> 46

Met Glu Ile Pro Met Ser Thr Val Val Lys Glu Thr Leu Thr Gln Leu  
                   1                                  5                                  10                                  15

Ser Ala His Arg Ala Leu Leu Thr Ser Asn Glu Thr Met Arg Leu Pro  
                                   20                                  25                                  30

Val Pro Thr His Lys Asn His Gln Leu Cys Ile Gly Glu Ile Phe Gln  
                   35                                  40                                  45

Gly Leu Asp Ile Leu Lys Asn Gln Thr Val Arg Gly Gly Thr Val Glu  
                   50                                  55                                  60

Met Leu Phe Gln Asn Leu Ser Leu Ile Lys Lys Tyr Ile Asp Arg Gln  
                   65                                  70                                  75                                  80

Lys Glu Lys Cys Gly Gln Tyr Ile Lys Ala Asn Ser Lys Phe Ile Gly  
                                   85                                  90                                  95

Ile Thr Glu Leu Arg Thr Arg Gln Phe Leu Asp Tyr Leu Gln Glu Phe  
                   100                                  105                                  110

09090916 021303

Leu Gly Val Met Ser Thr Glu Trp Ala Met Glu Gly  
 115 120

<210> 47  
 <211> 369  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: Murine IL-5  
 modified by substitution with tetanus toxoid  
 epitope

<220>  
 <221> CDS  
 <222> (1)..(369)

<220>  
 <221> mutation  
 <222> (88)..(150)  
 <223> Tetanus toxoid P30 epitope

<220>  
 <221> misc\_feature  
 <222> (1)..(87)  
 <223> DNA encoding amino acids 1-29 of murine IL5

<220>  
 <221> misc\_feature  
 <222> (151)..(366)  
 <223> DNA encoding amino acids 42-113 of murine IL5

<400> 47  
 atg gag att ccc atg agc aca gtg gtg aaa gag acc ttg aca cag ctg 48  
 Met Glu Ile Pro Met Ser Thr Val Val Lys Glu Thr Leu Thr Gln Leu  
 1 5 10 15  
 tcc gct cac cga gct ctg ttg aca agc aat gag acg atg ttc aac aac 96  
 Ser Ala His Arg Ala Leu Leu Thr Ser Asn Glu Thr Met Phe Asn Asn  
 20 25 30  
 ttc acc gtg agc ttc tgg ctg cgc gtg ccc aag gtg agc gcc agc cac 144  
 Phe Thr Val Ser Phe Trp Leu Arg Val Pro Lys Val Ser Ala Ser His  
 35 40 45  
 ctg gag tgc att gga gag atc ttt cag ggg cta gac ata ctg aag aat 192  
 Leu Glu Cys Ile Gly Glu Ile Phe Gln Gly Leu Asp Ile Leu Lys Asn  
 50 55 60  
 caa act gtc cgt ggg ggt acc gtg gaa atg cta ttc caa aac ctg tca 240  
 Gln Thr Val Arg Gly Gly Thr Val Glu Met Leu Phe Gln Asn Leu Ser  
 65 70 75 80  
 tta ata aag aaa tac atc gat aga caa aaa gag aag tgt ggc gag gag 288  
 Leu Ile Lys Lys Tyr Ile Asp Arg Gln Lys Glu Lys Cys Gly Glu Glu

00980916 004900

85 90 95  
 aga cgg agg acg agg cag ttc ctg gat tat ctg cag gag ttc ctt ggt 336  
 Arg Arg Arg Thr Arg Gln Phe Leu Asp Tyr Leu Gln Glu Phe Leu Gly  
 100 105 110

gtg atg agt aca gag tgg gca atg gaa ggc taa 369  
 Val Met Ser Thr Glu Trp Ala Met Glu Gly  
 115 120

<210> 48  
 <211> 122  
 <212> PRT  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: Murine Il-5  
 modified by substitution with tetanus toxoid  
 epitope

<400> 48  
 Met Glu Ile Pro Met Ser Thr Val Val Lys Glu Thr Leu Thr Gln Leu  
 1 5 10 15  
 Ser Ala His Arg Ala Leu Leu Thr Ser Asn Glu Thr Met Phe Asn Asn  
 20 25 30  
 Phe Thr Val Ser Phe Trp Leu Arg Val Pro Lys Val Ser Ala Ser His  
 35 40 45  
 Leu Glu Cys Ile Gly Glu Ile Phe Gln Gly Leu Asp Ile Leu Lys Asn  
 50 55 60  
 Gln Thr Val Arg Gly Gly Thr Val Glu Met Leu Phe Gln Asn Leu Ser  
 65 70 75 80  
 Leu Ile Lys Lys Tyr Ile Asp Arg Gln Lys Glu Lys Cys Gly Glu Glu  
 85 90 95  
 Arg Arg Arg Thr Arg Gln Phe Leu Asp Tyr Leu Gln Glu Phe Leu Gly  
 100 105 110  
 Val Met Ser Thr Glu Trp Ala Met Glu Gly  
 115 120

<210> 49  
 <211> 387  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: Murine Il-5  
 modified by substitution with tetanus toxoid  
 epitope

00000516-021903

<220>  
 <221> CDS  
 <222> (1)..(387)

<220>  
 <221> mutation  
 <222> (169)..(231)  
 <223> Tetanus toxoid P2 epitope

<220>  
 <221> misc\_feature  
 <222> (1)..(168)  
 <223> DNA encoding amino acids 1-56 of murine IL5

<220>  
 <221> misc\_feature  
 <222> (232)..(384)  
 <223> DNA encoding amino acids 63-113 of murine IL5

<400> 49  
 atg gag att ccc atg agc aca gtg gtg aaa gag acc ttg aca cag ctg 48  
 Met Glu Ile Pro Met Ser Thr Val Val Lys Glu Thr Leu Thr Gln Leu  
 1 5 10 15  
 tcc gct cac cga gct ctg ttg aca agc aat gag acg atg agg ctt cct 96  
 Ser Ala His Arg Ala Leu Leu Thr Ser Asn Glu Thr Met Arg Leu Pro  
 20 25 30  
 gtc cct act cat aaa aat cac cag cta tgc att gga gag atc ttt cag 144  
 Val Pro Thr His Lys Asn His Gln Leu Cys Ile Gly Glu Ile Phe Gln  
 35 40 45  
 ggg cta gac ata ctg aag aat caa ttc aac aac ttc acc gtg agc ttc 192  
 Gly Leu Asp Ile Leu Lys Asn Gln Phe Asn Asn Phe Thr Val Ser Phe  
 50 55 60  
 tgg ctg cgc gtg ccc aag gtg agc gcc agc cac ctg gag gtg gaa atg 240  
 Trp Leu Arg Val Pro Lys Val Ser Ala Ser His Leu Glu Val Glu Met  
 65 70 75 80  
 cta ttc caa aac ctg tca tta ata aag aaa tac atc gat aga caa aaa 288  
 Leu Phe Gln Asn Leu Ser Leu Ile Lys Lys Tyr Ile Asp Arg Gln Lys  
 85 90 95  
 gag aag tgt ggc gag gag aga cgg agg acg agg cag ttc ctg gat tat 336  
 Glu Lys Cys Gly Glu Glu Arg Arg Arg Thr Arg Gln Phe Leu Asp Tyr  
 100 105 110  
 ctg cag gag ttc ctt ggt gtg atg agt aca gag tgg gca atg gaa ggc 384  
 Leu Gln Glu Phe Leu Gly Val Met Ser Thr Glu Trp Ala Met Glu Gly  
 115 120 125  
 taa 387

05950916 0515002

<210> 50  
 <211> 128  
 <212> PRT  
 <213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Murine Il-5  
 modified by substitution with tetanus toxoid  
 epitope

<400> 50

Met Glu Ile Pro Met Ser Thr Val Val Lys Glu Thr Leu Thr Gln Leu  
 1 5 10 15

Ser Ala His Arg Ala Leu Leu Thr Ser Asn Glu Thr Met Arg Leu Pro  
 20 25 30

Val Pro Thr His Lys Asn His Gln Leu Cys Ile Gly Glu Ile Phe Gln  
 35 40 45

Gly Leu Asp Ile Leu Lys Asn Gln Phe Asn Asn Phe Thr Val Ser Phe  
 50 55 60

Trp Leu Arg Val Pro Lys Val Ser Ala Ser His Leu Glu Val Glu Met  
 65 70 75 80

Leu Phe Gln Asn Leu Ser Leu Ile Lys Lys Tyr Ile Asp Arg Gln Lys  
 85 90 95

Glu Lys Cys Gly Glu Glu Arg Arg Arg Thr Arg Gln Phe Leu Asp Tyr  
 100 105 110

Leu Gln Glu Phe Leu Gly Val Met Ser Thr Glu Trp Ala Met Glu Gly  
 115 120 125

<210> 51  
 <211> 351  
 <212> DNA  
 <213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Murine Il-5  
 modified by substitution with tetanus toxoid  
 epitope

<220>

<221> CDS  
 <222> (1)..(351)

<220>

<221> mutation  
 <222> (88)..(132)  
 <223> Tetanus toxoid P2 epitope

<220>

206120"9T603550

<221> misc\_feature

<222> (1)..(87)

<223> DNA encoding amino acids 1-29 of murine IL5

<220>

<221> misc\_feature

<222> (133)..(348)

<223> DNA encoding amino acids 42-113 of murine IL5

<400> 51

```

atg gag att ccc atg agc aca gtg gtg aaa gag acc ttg aca cag ctg      48
Met Glu Ile Pro Met Ser Thr Val Val Lys Glu Thr Leu Thr Gln Leu
  1             5             10             15

tcc gct cac cga gct ctg ttg aca agc aat gag acg atg cag tac atc      96
Ser Ala His Arg Ala Leu Leu Thr Ser Asn Glu Thr Met Gln Tyr Ile
             20             25             30

aaa gct aac tcc aaa ttc atc ggt atc acc gag ctg tgc att gga gag      144
Lys Ala Asn Ser Lys Phe Ile Gly Ile Thr Glu Leu Cys Ile Gly Glu
             35             40             45

atc ttt cag ggg cta gac ata ctg aag aat caa act gtc cgt ggg ggt      192
Ile Phe Gln Gly Leu Asp Ile Leu Lys Asn Gln Thr Val Arg Gly Gly
             50             55             60

acc gtg gaa atg cta ttc caa aac ctg tca tta ata aag aaa tac atc      240
Thr Val Glu Met Leu Phe Gln Asn Leu Ser Leu Ile Lys Lys Tyr Ile
             65             70             75             80

gat aga caa aaa gag aag tgt ggc gag gag aga cgg agg acg agg cag      288
Asp Arg Gln Lys Glu Lys Cys Gly Glu Glu Arg Arg Arg Thr Arg Gln
             85             90             95

ttc ctg gat tat ctg cag gag ttc ctt ggt gtg atg agt aca gag tgg      336
Phe Leu Asp Tyr Leu Gln Glu Phe Leu Gly Val Met Ser Thr Glu Trp
             100             105             110

gca atg gaa ggc taa                                          351
Ala Met Glu Gly
             115

```

<210> 52

<211> 116

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Murine Il-5  
modified by substitution with tetanus toxoid  
epitope

<400> 52

```

Met Glu Ile Pro Met Ser Thr Val Val Lys Glu Thr Leu Thr Gln Leu
  1             5             10             15

```

20250516 09:05:56



Ser Ala His Arg Ala Leu Leu Thr Ser Asn Glu Thr Met Gln Tyr Ile  
                     20                    25                    30  
 Lys Ala Asn Ser Lys Phe Ile Gly Ile Thr Glu Leu Cys Ile Gly Glu  
                     35                    40                    45  
 Ile Phe Gln Gly Leu Asp Ile Leu Lys Asn Gln Thr Val Arg Gly Gly  
                     50                    55                    60  
 Thr Val Glu Met Leu Phe Gln Asn Leu Ser Leu Ile Lys Lys Tyr Ile  
                     65                    70                    75                    80  
 Asp Arg Gln Lys Glu Lys Cys Gly Glu Glu Arg Arg Arg Thr Arg Gln  
                     85                    90                    95  
 Phe Leu Asp Tyr Leu Gln Glu Phe Leu Gly Val Met Ser Thr Glu Trp  
                     100                    105                    110  
 Ala Met Glu Gly  
                     115

<210> 53  
 <211> 369  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: Murine IL-5  
       modified by substitution with tetanus toxoid  
       epitope

<220>  
 <221> CDS  
 <222> (1)..(369)

<220>  
 <221> mutation  
 <222> (250)..(294)  
 <223> Tetanus toxoid P2 epitope

<220>  
 <221> misc\_feature  
 <222> (1)..(249)  
 <223> DNA encoding amino acids 1-83 of murine IL5

<220>  
 <221> misc\_feature  
 <222> (295)..(366)  
 <223> DNA encoding amino acids 90-113 of murine IL5

<400> 53  
 atg gag att ccc atg agc aca gtg gtg aaa gag acc ttg aca cag ctg 48  
 Met Glu Ile Pro Met Ser Thr Val Val Lys Glu Thr Leu Thr Gln Leu  
       1                    5                    10                    15

20250915 09:59:50

```

tcc gct cac cga gct ctg ttg aca agc aat gag acg atg agg ctt cct 96
Ser Ala His Arg Ala Leu Leu Thr Ser Asn Glu Thr Met Arg Leu Pro
          20          25          30

gtc cct act cat aaa aat cac cag cta tgc att gga gag atc ttt cag 144
Val Pro Thr His Lys Asn His Gln Leu Cys Ile Gly Glu Ile Phe Gln
          35          40          45

ggg cta gac ata ctg aag aat caa act gtc cgt ggg ggt acc gtg gaa 192
Gly Leu Asp Ile Leu Lys Asn Gln Thr Val Arg Gly Gly Thr Val Glu
          50          55          60

atg cta ttc caa aac ctg tca tta ata aag aaa tac atc gat aga caa 240
Met Leu Phe Gln Asn Leu Ser Leu Ile Lys Lys Tyr Ile Asp Arg Gln
          65          70          75          80

aaa gag aag cag tac atc aag gcc aac tcc aag ttc atc ggc atc acc 288
Lys Glu Lys Gln Tyr Ile Lys Ala Asn Ser Lys Phe Ile Gly Ile Thr
          85          90          95

gag ctg agg acg agg cag ttc ctg gat tat ctg cag gag ttc ctt ggt 336
Glu Leu Arg Thr Arg Gln Phe Leu Asp Tyr Leu Gln Glu Phe Leu Gly
          100          105          110

gtg atg agt aca gag tgg gca atg gaa ggc taa 369
Val Met Ser Thr Glu Trp Ala Met Glu Gly
          115          120

<210> 54
<211> 122
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Murine Il-5
        modified by substitution with tetanus toxoid
        epitope

<400> 54
Met Glu Ile Pro Met Ser Thr Val Val Lys Glu Thr Leu Thr Gln Leu
  1          5          10          15

Ser Ala His Arg Ala Leu Leu Thr Ser Asn Glu Thr Met Arg Leu Pro
          20          25          30

Val Pro Thr His Lys Asn His Gln Leu Cys Ile Gly Glu Ile Phe Gln
          35          40          45

Gly Leu Asp Ile Leu Lys Asn Gln Thr Val Arg Gly Gly Thr Val Glu
          50          55          60

Met Leu Phe Gln Asn Leu Ser Leu Ile Lys Lys Tyr Ile Asp Arg Gln
          65          70          75          80

Lys Glu Lys Gln Tyr Ile Lys Ala Asn Ser Lys Phe Ile Gly Ile Thr
          85          90          95

```

Glu Leu Arg Thr Arg Gln Phe Leu Asp Tyr Leu Gln Glu Phe Leu Gly  
 100 105 110

Val Met Ser Thr Glu Trp Ala Met Glu Gly  
 115 120

<210> 55  
 <211> 393  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: Murine Il-5  
 modified by substitution with tetanus toxoid  
 epitope

<220>  
 <221> CDS  
 <222> (1)..(393)  
 <220>  
 <221> mutation  
 <222> (256)..(318)  
 <223> Tetanus toxoid P30 epitope

<220>  
 <221> misc\_feature  
 <222> (1)..(255)  
 <223> DNA encoding amino acids 1-85 of murine IL5

<220>  
 <221> misc\_feature  
 <222> (319)..(390)  
 <223> DNA encoding amino acids 90-113 of murine IL5

<400> 55  
 atg gag att ccc atg agc aca gtg gtg aaa gag acc ttg aca cag ctg 48  
 Met Glu Ile Pro Met Ser Thr Val Val Lys Glu Thr Leu Thr Gln Leu  
 1 5 10 15  
 tcc gct cac cga gct ctg ttg aca agc aat gag acg atg agg ctt cct 96  
 Ser Ala His Arg Ala Leu Leu Thr Ser Asn Glu Thr Met Arg Leu Pro  
 20 25 30  
 gtc cct act cat aaa aat cac cag cta tgc att gga gag atc ttt cag 144  
 Val Pro Thr His Lys Asn His Gln Leu Cys Ile Gly Glu Ile Phe Gln  
 35 40 45  
 ggg cta gac ata ctg aag aat caa act gtc cgt ggg ggt acc gtg gaa 192  
 Gly Leu Asp Ile Leu Lys Asn Gln Thr Val Arg Gly Gly Thr Val Glu  
 50 55 60  
 atg cta ttc caa aac ctg tca tta ata aag aaa tac atc gat aga caa 240  
 Met Leu Phe Gln Asn Leu Ser Leu Ile Lys Lys Tyr Ile Asp Arg Gln

20050916 09:50:50

65	70	75	80	
aaa gag aag tgt ggc ttc aac aac ttc acc gtg agc ttc tgg ctg cgc				288
Lys Glu Lys Cys Gly Phe Asn Asn Phe Thr Val Ser Phe Trp Leu Arg				
	85	90	95	
gtg ccc aag gtg agc gcc agc cac ctg gag agg acg agg cag ttc ctg				336
Val Pro Lys Val Ser Ala Ser His Leu Glu Arg Thr Arg Gln Phe Leu				
	100	105	110	
gat tat ctg cag gag ttc ctt ggt gtg atg agt aca gag tgg gca atg				384
Asp Tyr Leu Gln Glu Phe Leu Gly Val Met Ser Thr Glu Trp Ala Met				
	115	120	125	
gaa ggc taa				393
Glu Gly				
130				

<210> 56  
 <211> 130  
 <212> PRT  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: Murine Il-5  
 modified by substitution with tetanus toxoid  
 epitope

<400> 56  
 Met Glu Ile Pro Met Ser Thr Val Val Lys Glu Thr Leu Thr Gln Leu  
 1 5 10 15  
 Ser Ala His Arg Ala Leu Leu Thr Ser Asn Glu Thr Met Arg Leu Pro  
 20 25 30  
 Val Pro Thr His Lys Asn His Gln Leu Cys Ile Gly Glu Ile Phe Gln  
 35 40 45  
 Gly Leu Asp Ile Leu Lys Asn Gln Thr Val Arg Gly Gly Thr Val Glu  
 50 55 60  
 Met Leu Phe Gln Asn Leu Ser Leu Ile Lys Lys Tyr Ile Asp Arg Gln  
 65 70 75 80  
 Lys Glu Lys Cys Gly Phe Asn Asn Phe Thr Val Ser Phe Trp Leu Arg  
 85 90 95  
 Val Pro Lys Val Ser Ala Ser His Leu Glu Arg Thr Arg Gln Phe Leu  
 100 105 110  
 Asp Tyr Leu Gln Glu Phe Leu Gly Val Met Ser Thr Glu Trp Ala Met  
 115 120 125  
 Glu Gly  
 130

20060420 14:03:50

```
<220>
<223> Description of Artificial Sequence: Murine Il-5
modified by substitution with tetanus toxoid
epitope
```

```
<220>
<221> mutation
<222> (250)..(312)
<223> Tetanus toxoid P30 epitope
```

```
<220>
<221> misc_feature
<222> (313)..(384)
<223> DNA encoding amino acids 90-113 of murine IL5
```

<400> 57																
atg	gag	att	ccc	atg	agc	aca	gtg	gtg	aaa	gag	acc	ttg	aca	cag	ctg	48
Met	Glu	Ile	Pro	Met	Ser	Thr	Val	Val	Lys	Glu	Thr	Leu	Thr	Gln	Leu	
1				5					10					15		
tcc	gct	cac	cga	gct	ctg	ttg	aca	agc	aat	gag	acg	atg	agg	ctt	cct	96
Ser	Ala	His	Arg	Ala	Leu	Leu	Thr	Ser	Asn	Glu	Thr	Met	Arg	Leu	Pro	
			20					25					30			
gtc	cct	act	cat	aaa	aat	cac	cag	cta	tgc	att	gga	gag	atc	ttt	cag	144
Val	Pro	Thr	His	Lys	Asn	His	Gln	Leu	Cys	Ile	Gly	Glu	Ile	Phe	Gln	
		35					40					45				
ggg	cta	gac	ata	ctg	aag	aat	caa	act	gtc	cgt	ggg	ggg	acc	gtg	gaa	192
Gly	Leu	Asp	Ile	Leu	Lys	Asn	Gln	Thr	Val	Arg	Gly	Gly	Thr	Val	Glu	
	50					55					60					
atg	cta	ttc	caa	aac	ctg	tca	tta	ata	aag	aaa	tac	atc	gat	aga	caa	240
Met	Leu	Phe	Gln	Asn	Leu	Ser	Leu	Ile	Lys	Lys	Tyr	Ile	Asp	Arg	Gln	
65					70					75					80	
aaa	gag	aag	ttc	aac	aac	ttc	acc	gtg	agc	ttc	tgg	ctg	cgc	gtg	ccc	288
Lys	Glu	Lys	Phe	Asn	Asn	Phe	Thr	Val	Ser	Phe	Trp	Leu	Arg	Val	Pro	
				85					90					95		
aag	gtg	agc	gcc	agc	cac	ctg	gag	agg	acg	agg	cag	ttc	ctg	gat	tat	336

<220>  
<223> Description of Artificial Sequence: Murine Il-5  
modified by substitution with tetanus toxoid  
epitopes

<400> 59																
atg	gag	att	ccc	atg	agc	aca	gtg	gtg	aaa	gag	acc	ttg	aca	cag	ctg	48
Met	Glu	Ile	Pro	Met	Ser	Thr	Val	Val	Lys	Glu	Thr	Leu	Thr	Gln	Leu	
1				5				10				15				
tcc	gct	cac	cga	gct	ctg	ttg	aca	agc	aat	gag	acg	atg	agg	ctt	cct	96
Ser	Ala	His	Arg	Ala	Leu	Leu	Thr	Ser	Asn	Glu	Thr	Met	Arg	Leu	Pro	
				20				25				30				
gtc	cct	act	cat	aaa	aat	cac	cag	cta	tgc	att	gga	gag	atc	ttt	cag	144
Val	Pro	Thr	His	Lys	Asn	His	Gln	Leu	Cys	Ile	Gly	Glu	Ile	Phe	Gln	
35								40				45				
ggg	cta	gac	ata	ctg	aag	aat	caa	act	gtc	cgt	ggg	ggt	acc	gtg	gaa	192
Gly	Leu	Asp	Ile	Leu	Lys	Asn	Gln	Thr	Val	Arg	Gly	Gly	Thr	Val	Glu	
50				55				60								
atg	cta	ttc	caa	aac	ctg	tca	tta	ata	aag	aaa	tac	atc	gat	aga	caa	240
Met	Leu	Phe	Gln	Asn	Leu	Ser	Leu	Ile	Lys	Lys	Tyr	Ile	Asp	Arg	Gln	
65				70				75				80				
aaa	gag	aag	tgt	ggc	cag	tac	atc	aag	gcc	aac	tcc	aag	ttc	atc	ggc	288
Lys	Glu	Lys	Cys	Gly	Gln	Tyr	Ile	Lys	Ala	Asn	Ser	Lys	Phe	Ile	Gly	
				85				90				95				
atc	acc	gag	ctg	ttc	aac	aac	ttc	acc	gtg	agc	ttc	tgg	ctg	cgc	gtg	336
Ile	Thr	Glu	Leu	Phe	Asn	Asn	Phe	Thr	Val	Ser	Phe	Trp	Leu	Arg	Val	
100								105				110				
ccc	aag	gtg	agc	gcc	agc	cac	ctg	gag	agg	acg	agg	cag	ttc	ctg	gat	384
Pro	Lys	Val	Ser	Ala	Ser	His	Leu	Glu	Arg	Thr	Arg	Gln	Phe	Leu	Asp	
115								120				125				

ggc taa 438  
Gly  
145

```
<220>
<223> Description of Artificial Sequence: Murine Il-5
modified by substitution with tetanus toxoid
epitopes
```

```
<210> 61
<211> 57
<212> DNA
<213> Homo sapiens
```



<220>  
 <221> CDS  
 <222> (1)..(57)  
 <223> DNA encoding natural human IL5 leader sequence

<400> 61  
 atg agg atg ctt ctg cat ttg agt ttg ctg gct ctt gga gct gcc tac 48  
 Met Arg Met Leu Leu His Leu Ser Leu Leu Ala Leu Gly Ala Ala Tyr  
           1                  5                  10                  15  
  
 gtg tat gcc 57  
 Val Tyr Ala

<210> 62  
 <211> 19  
 <212> PRT  
 <213> Homo sapiens

<400> 62  
 Met Arg Met Leu Leu His Leu Ser Leu Leu Ala Leu Gly Ala Ala Tyr  
           1                  5                  10                  15  
  
 Val Tyr Ala

<210> 63  
 <211> 60  
 <212> DNA  
 <213> Mus musculus

<220>  
 <221> CDS  
 <222> (1)..(60)  
 <223> DNA encoding natural murine IL5 leader sequence

<400> 63  
 atg aga agg atg ctt ctg cac ttg agt gtt ctg act ctc agc tgt gtc 48  
 Met Arg Arg Met Leu Leu His Leu Ser Val Leu Thr Leu Ser Cys Val  
           1                  5                  10                  15  
  
 tgg gcc act gcc 60  
 Trp Ala Thr Ala  
                   20

<210> 64  
 <211> 20  
 <212> PRT  
 <213> Mus musculus

<400> 64  
 Met Arg Arg Met Leu Leu His Leu Ser Val Leu Thr Leu Ser Cys Val  
           1                  5                  10                  15

0960516.021902

```
<210> 65
<211> 13
<212> PRT
<213> Artificial Sequence
```

<223> Description of Artificial Sequence: Promiscuous T helper epitope

Ala Lys Phe Val Ala Ala Trp Thr Leu Lys Ala Ala Ala  
1 5 10